

Number Theory

Introduction

Vision

The goal of this open-source number theory textbook is to gather up all the core subfields of number theory into one text. By making it open-source, everyone will be able to contribute in terms of adding new material and improving existing material, and tailor it to their own learning or teaching.

It is an era of mass collaboration, in mathematics and many other fields. I aim to follow the example of other successful online textbooks such as CRing and the Stacks Project. Because we all are good at different subfields, we will be able to achieve much more if each person writes notes on an area they have studied well. For instance, I have found that one of the best ways to learn a semi-advanced topic is to find an undergraduate thesis on it: because the author has spent a long time thinking on the subject, had long conversations with advisors, and taken time to lay out the big motivations and deeper connections.

The philosophy behind this textbook is the following. If you'd like to contribute to this work I'd recommend following these guidelines.

1. Take a problem-oriented approach. In other words, **give motivation** behind abstract theory by relating them to interesting, concrete problems.
2. Create a **user-friendly learning resource**: Start each chapter by telling the reader why the material matters, what problems in number theory it solves, and how it fits into the big picture.¹ For instance, class field theory gives a way to understand field extensions through information intrinsic to the field, it gives framework for reciprocity laws (among many other things, see chapter 16), and it is part of the larger Langlands program. Give a summary of the takeaway ideas after the chapter, and exercises that help the reader conceptually grasp the material. Always motivate proofs, especially long hard ones, and tell the reader why the big theorems matter. Make connections between different ways of approaching a particular problems. Highlight recurring techniques.

(These points have not all been implemented in the chapters here, but it is what I'm shooting for.)

¹I learned this style from Patrick Winston. You can see it in action in his textbook on Artificial Intelligence.

3. Be a **self-contained** work: This means that proofs should refer to theorems in the text itself, possibly from a previous part (for instance, refer to the Algebraic Number Theory section in the Class Field Theory section.) (I may eventually move some of the material into appendices, such as the complex analysis background required for analytic number theory.)
4. Connect up elementary with advanced number theory, and offer a road map of the subject.

Some more notes:

1. All material is under a creative commons license.
2. The source is available at <http://github.com/holdenlee/number-theory>.
3. Please contribute! You will be credited. It doesn't have to be finished/polished stuff—it takes much less time to edit material that's already written than to write it myself.
 - (a) If you find mistakes, add missing sections, edit the material, etc., then send me a pull request on github (<http://github.com/holdenlee/number-theory>).
 - (b) Alternatively, you email me corrections, suggestions, and contributions at holdenlee@alum.mit.edu and I'll update the document. In either case, I'll list you as a contributor.
4. Much of this is in very rough shape right now.
5. (*) denotes optional material. (†) denotes theorems that will be made obsolete by stronger theorems later (for example, the $p \equiv 1 \pmod{n}$ case of Dirichlet's theorem).

This file was last updated August 14, 2017.

Contributors: Holden Lee, Oleg Muskarov, Teo Andrica

Thanks for proofreading: Delong Meng, Timo Keller, Dan Elbert

For a list of references, see refs.bib.

Contents

I	Algebraic Number Theory	1
1	Rings of integers	3
1	Integrality	3
2	Norms and Traces	7
3	Discriminant	9
4	Integral bases	11
5	Problems	15
2	Ideal factorization	17
1	Discrete Valuation Rings	17
2	Dedekind Domains	18
3	Primary decomposition*	21
4	Ideal class group	21
5	Factorization in extensions	22
6	Computing factorizations	24
7	Decomposition and inertia groups	26
7.1	Decomposition group	28
7.2	Inertia group	29
7.3	Further properties and applications	31
8	Problems	32
3	The class group	33
1	Norms of ideals	33
2	Minkowski's Theorem	35
3	Finiteness of the class number	36
4	Example: Quadratic extensions	41
4	The algebra of quadratic forms	47
1	Quadratic forms	47
2	Representing integers	48
3	Reduction of quadratic forms	49
3.1	Examples	50
4	Ideals on quadratic rings	54

4.1	Proper and invertible ideals	56
5	Gauss composition	58
6	Ideal class group of an order	62
7	Cube law	63
5	Units in number fields	67
1	Units	67
2	Dirichlet's unit theorem	68
3	S -units	71
4	Examples and algorithms	71
5	Regulator	71
6	Cyclotomic fields	73
1	Cyclotomic polynomials	73
2	Ring of integers	75
3	Subfields of cyclotomic extensions	79
4	Fermat's last theorem: Regular primes	80
5	Exercises	81
7	Valuations and completions	83
1	Case study: p -adic integers	83
1.1	p -adics as an inverse limit	84
1.2	p -adics as completions	84
1.3	Units in \mathbb{Z}_p	84
1.4	Monsky's Theorem*	85
2	Valuations	87
2.1	Equivalent valuations	88
3	Places	89
3.1	Approximation	91
4	Completion	93
4.1	Completions of archimedean fields; Classification of places	93
4.2	Completions of nonarchimedean fields	94
5	Hensel's lemma	96
6	Extending valuations	98
7	Places as Galois orbits	100
8	Krasner's lemma and consequences	101
8	Local and global fields	103
1	Topology of local fields	103
1.1	Open sets and continuity	104
2	Unramified extensions	104
3	Ramified extensions	108
4	Witt vectors*	110

4.1	Frobenius and Transfer maps	114
5	Extending valuations on global fields	114
6	Product formula	115
7	Problems	116
9	Ramification	117
1	Lattices and χ	117
1.1	Filtrations of modules	117
1.2	The function χ_A	118
1.3	χ and localization	119
1.4	Discriminant of bilinear forms	120
2	Discriminant and different	121
2.1	Basic properties	122
3	Discriminant and ramification	123
3.1	Types of ramification	125
3.2	Computation of different	126
4	Ramification groups	127
4.1	$\mathfrak{D}_{L/K}$ and i_G	128
4.2	Filtration of ramification groups	131
4.3	First ramification group	133
5	Herbrand's Theorem	136
5.1	Functions φ and ψ	136
5.2	Transitivity of φ and ψ	136
6	Hasse-Arf Theorem	139
10	Geometric algebraic number theory	141
1	Generalized ideal classes	141
2	Counting lattice points	142
3	Riemann-Roch problem	143
4	Asymptotics of generalized ideal classes	143
II	Class Field Theory	145
11	Class Field Theory: Introduction	147
1	Frobenius elements	147
1.1	Examples	149
1.2	The Frobenius map is a nice homomorphism	151
2	Local reciprocity	151
3	Ray class groups	153
4	Global reciprocity	155
5	Ideles	157
5.1	Ray class groups vs. ideles	159

6	Global reciprocity via ideles	162
6.1	Connecting the two formulations	164
7	Kronecker-Weber Theorem	166
8	Problems	167
12	Group homology and cohomology	169
1	Projectives and injectives	170
2	Complexes	171
3	Homology and cohomology	172
4	Derived functors	173
4.1	Right derived functors and Ext	173
4.1.1	Covariant case	173
4.1.2	Contravariant case	174
4.2	Left derived functors and Tor	175
4.3	Long exact sequences	176
5	Homological and cohomological functors	177
6	Group cohomology	179
7	Bar resolutions	181
8	Group homology	182
8.1	Shapiro's lemma	183
9	Tate groups	185
9.1	Complete resolution*	187
9.2	Dimension shifting	188
10	Cup products	189
10.1	Cup product calculations	191
11	Change of group	192
11.1	Construction of maps	193
11.2	Extending maps to Tate cohomology	194
11.3	Further properties	196
11.4	Inflation-restriction exact sequence	197
11.5	Transfer	198
12	Cohomology of cyclic groups	199
12.1	Herbrand quotient	201
13	Tate's Theorem	202
14	Profinite groups	203
15	Nonabelian cohomology	204
13	Introduction to Galois cohomology	207
1	Basic results	207
2	Kummer theory	209
3	Nonabelian Galois cohomology	212
4	Brauer group	216
4.1	Background from noncommutative algebra	216

4.2	Central simple algebras and the Brauer group	218
4.3	Subfields and splitting of central simple algebras	219
5	Brauer group and cohomology	220
5.1	The Brauer group is a second cohomology group	220
5.2	Exact sequence of Brauer groups	222
6	Problems	223
14	Local class field theory	225
1	Cohomology of the units	226
2	The invariant map	228
2.1	Defining the invariant maps	228
2.2	Compatibility of the invariant maps	229
3	$H^2(\overline{K}/K) \cong H^2(K^{\text{ur}}/K)$	231
3.1	First proof (Brauer group)	231
3.2	Second proof (Herbrand quotient calculation)	232
3.2.1	Herbrand quotient calculation	232
3.2.2	First inequality	234
3.2.3	Second inequality	234
3.2.4	Finishing the proof	235
4	Class formations	235
4.1	Class formations in the abstract	235
4.2	Class formations for local class field theory	242
5	Examples	244
5.1	Unramified case	244
5.2	Ramified case	245
6	Hilbert symbols	246
7	Existence theorem	249
7.1	Existence theorem in the abstract	249
7.2	Existence theorem for local class field theory	251
8	Topology of the local reciprocity map	251
8.1	Uniqueness of the reciprocity map	252
15	Global class field theory	255
1	Basic definitions	255
2	The first inequality	256
2.1	Reduce to finite number of places	257
2.2	Cohomology of \mathbb{I}_L^S and \mathbb{I}_L	258
2.3	Cohomology of lattices and U_L^T	260
2.4	Herbrand quotient of \mathbf{C}_L	262
2.5	The Frobenius map is surjective	262
3	The second inequality	263
3.1	Analytic approach	263
3.2	Algebraic approach	264

3.2.1	Local computations	265
3.2.2	Constructing E	265
3.3	Finishing the proof	266
3.4	Local-to-global principle for algebras	267
4	Proof of the reciprocity law	267
4.1	(A) holds for $\mathbb{Q}(\zeta_n)/\mathbb{Q}$	269
4.2	(B) holds for all cyclomic extensions	270
4.3	(A) for cyclotomic implies (B) for α split by cyclic cyclotomic	270
4.4	(B) for cyclic cyclotomic implies (B) in general	271
4.5	(B) implies (A) for all abelian extensions	272
5	The ideles are a class formation	273
6	Existence theorem	277
16	Applications	279
1	Reciprocity laws	280
1.1	Weak reciprocity and the Legendre symbol	280
1.2	Strong reciprocity and the Hilbert symbol	282
1.3	Quadratic and biquadratic reciprocity	285
1.4	Reciprocity for odd primes	286
2	Hasse-Minkowski Theorem	290
2.1	Hasse norm theorem	290
2.2	Quadratic forms	291
2.2.1	Proof for $n \leq 2$	292
2.2.2	Proof for $n = 3$	293
2.2.3	Proof for $n = 4$	293
2.2.4	Proof for $n \geq 5$	295
3	Chebotarev density theorem	297
3.1	Proof	297
3.2	Applications	299
4	Splitting of primes	301
4.1	Splitting of primes	301
4.2	Roots of polynomials over finite fields	302
5	Hilbert class field	303
6	Primes represented by quadratic forms	304
7	Introduction to the Langlands program	307
7.1	Definitions	308
7.2	Class field theory is 1-dimensional Langlands	309
7.2.1	Galois representations are automorphic representations	309
7.2.2	Artin L -functions are Hecke L -functions	310
7.2.3	Algebraic varieties and Galois representations	311
7.3	Elliptic curves and 2-dimensional Langlands	312
7.3.1	Galois representations and automorphic representations	312
7.3.2	Elliptic curves and Galois representations	313

8	Problems	314
17	Complex multiplication	317
1	Elliptic curves over \mathbb{C}	318
2	Complex multiplication over \mathbb{C}	319
2.1	Embedding the endomorphism ring	319
2.2	The class group parameterizes elliptic curves	320
2.3	Ideals define maps	322
3	Defining CM elliptic curves over $\overline{\mathbb{Q}}$	322
4	Hilbert class field	324
4.1	Motivation: Class field theory for $\mathbb{Q}(\zeta_n)$ and Kronecker-Weber	324
4.1.1	The case of \mathbb{Q}	324
4.1.2	The case of K	325
4.1.3	The case of K : Part 1	325
4.1.4	The case of K : Part 2	325
4.2	The Galois group and class group act compatibly	326
4.3	Hilbert class field	327
5	Maximal abelian extension	330
6	The Main Theorem of Complex Multiplication	335
6.1	The associated Grössencharacter	339
7	L -series of CM elliptic curve	342
7.1	Defining the L -function	342
7.2	Analytic continuation	344
III	Analytic Number Theory	347
18	Elementary estimates for primes	349
1	Chebyshev's Theorem	349
1.1	Comparing the three functions	350
1.2	Upper Bound	350
1.3	Lower Bound	351
1.4	The n th prime	352
19	Crash course in complex analysis	353
1	Holomorphic functions	353
2	Complex integration	354
3	Cauchy's Theorem	355
4	Power series and Laurent series	356
4.1	Cauchy's residue formula	357
5	Convergence	358
6	Series and product developments	358
7	Gamma function	359

20	Dirichlet series	361
1	Dirichlet series, convergence	361
2	Basic properties	362
3	Dirichlet generating functions	364
4	Summing coefficients	365
21	Zeta functions and the prime number theorem	367
1	Prime number theorem: Outline	367
1.1	The big picture	368
1.2	Main steps	368
2	Riemann zeta function	370
3	Zeros of zeta	374
4	Prime number theorem: proof	378
5	The Riemann hypothesis	383
22	L-functions and Dirichlet's theorem	385
1	Outline	385
2	L -functions	386
3	Zeros of L	393
4	Prime number theorem in arithmetic progressions	397
5	Siegel zero	401
5.1	$L'(\beta, \chi)$ is not too large	401
5.2	$L(1, \chi)$ is not too small	402
5.3	Proof of Siegel-Walfisz	406
23	Special values of L-functions	407
0	Introduction	407
0.1	Points of interest	407
0.2	Road map	408
0.3	First explorations	409
1	The analytic class number formula	411
1.1	Quadratic fields	411
1.2	Proof in general	415
1.2.1	Reduce to sum over principal ideals	415
1.2.2	Reduce to sum over numbers	416
1.2.3	Reduce to a volume calculation	416
1.2.4	Calculate the volume	418
2	Special values of L -functions	419
2.1	Evaluating $L(1, \chi)$ for quadratic characters	419
2.2	The class number counts quadratic residues/nonresidues	422
2.2.1	$p \equiv 1 \pmod{4}$	422
2.2.2	$p \equiv 3 \pmod{4}$	423
3	For further study	424

4	Appendix	425
A	Multiplication modulo n	427
1	Introduction	427
2	Order of an element	428
3	Euler's theorem and Fermat's little theorem	430
4	Examples	432
	4.1 Using Euler's theorem	432
	4.2 Computing the order	432
5	Groups	436
6	Primitive roots	436
7	Multiplicative structure of $\mathbb{Z}/n\mathbb{Z}$	439
8	Wilson's theorem	439
9	Problems	442
B	Quadratic residues	443
1	Quadratic residues	443
2	Quadratic reciprocity	445
3	Jacobi symbol	448
C	Unique factorization	449
1	Unique factorization domains	449
	1.1 Step 1: Euclidean domains	451
	1.2 Step 2: Euclidean domain \implies PID	451
	1.3 Step 3: PID \implies UFD	452
2	Example: $x^2 + y^2 = n$	452
3	Problems	453
D	Polynomials	455
1	Gauss's argument	455
	1.1 More Proofs	457
	1.2 Problems	459
2	Main Theorems	459
	2.1 Problems	460
3	Arithmetic Properties	462
	3.1 Problems	465
4	Polynomials in Number Theory	466
	4.1 Problems	468
5	Resultant	469
E	Galois Theory	473
1	Galois groups and Galois extensions	473
2	Fixed fields	473

3	Splitting fields	474
4	Fundamental theorem of Galois theory	475
5	Cubic and quartic equations	477
6	Quintic equations	477
7	Inverse limits and profinite groups	477
	7.1 Limits	477
	7.2 Profinite groups	480
8	Infinite Galois theory	481
F	Arithmetic over Finite Fields	485
1	Characters	485
	1.1 Dirichlet characters	487
	1.2 Characters on finite fields	488
2	Gauss Sums	489
3	Enumerating Solutions	491
4	Applications to Waring's Problem	493

Part I

Algebraic Number Theory

Chapter 1

Rings of integers

When we have a field extension L of \mathbb{Q} , we would like to define a ring of integers for L , with properties similar to the ring $\mathbb{Z} \subseteq \mathbb{Q}$. We will define this ring of integers in a slightly more general context.

1 Integrality

Definition 1.1.1: Let A be an integral domain and L a field containing A . An element of $x \in L$ is **integral** over A if it is the zero of a monic polynomial with coefficients in A :

$$x^n + a_{n-1}x^{n-1} + \cdots + a_1x + a_0 = 0, \quad n \geq 1, \quad a_0, \dots, a_{n-1} \in A.$$

The **integral closure** of A in L is the set of elements of L integral over A .

Example 1.1.2: The integral closure of \mathbb{Z} in \mathbb{Q} is simply \mathbb{Z} itself (we see this more generally in Proposition 1.1.8). Thus, integral closure generalizes the notion of what it means to be an “integer” in other number fields. As we will see in Example 1.4.7, for d squarefree, the integral closure of $\mathbb{Q}(\sqrt{d})$ is $\mathbb{Z}[\sqrt{d}]$ when $d \equiv 3 \pmod{4}$ and $\mathbb{Z}\left[\frac{1+\sqrt{d}}{2}\right]$ when $d \equiv 1 \pmod{4}$. Algebra is much nicer in integral extensions—which is why, for instance, we would study $\mathbb{Z}\left[\frac{1+\sqrt{-3}}{2}\right]$ rather than just $\mathbb{Z}[\sqrt{-3}]$.

Theorem 1.1.3. *Let L be a field containing the ring A . Then the elements of L integral over A form a ring.*

Proof. We give two proofs. We need to show that if a, b are algebraic over A then so are $a + b$ and ab .

Proof 1: Let p, q be the minimal polynomials of a, b , let a_1, \dots, a_k be the conjugates of a and b_1, \dots, b_l be the conjugates of b . The coefficients of

$$\prod_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} (x - (a_i + b_j)), \quad \prod_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} (x - (a_i b_j))$$

are symmetric in the a_i and symmetric in the b_j so by the Fundamental Theorem of Symmetric Polynomials can be written in terms of the elementary symmetric polynomials in the a_i and in the b_j , with coefficients in A . By Vieta's Theorem these are expressible in terms of the coefficients of p, q , which are in A . Hence these polynomials have coefficients in A . They have $a + b, ab$ as roots, as desired.

Proof 2: We use the following lemma.

Lemma 1.1.4 (Criterion for integrality). *An element $\alpha \in L$ is integral over A if and only if there exists a nonzero finitely generated A -submodule of L such that $\alpha M \subseteq M$. If so, then we can take $M = A[\alpha]$.*

Example 1.1.5: For example, $\frac{1}{\sqrt{2}}$ fails this criterion over \mathbb{Z} —multiplying by it has the effect of making M “finer.” $\sqrt{2}$, however, is integral.

In the case $A = \mathbb{Z}$ and $B = \mathbb{Q}$, $a \in \mathbb{Q}$ is integral over \mathbb{Z} iff $a \in \mathbb{Z}$. Indeed, $a \in \mathbb{Z}$ satisfies $x - a$, and if $a \notin \mathbb{Z}$, then powers of a contain arbitrarily large denominators so $\mathbb{Z}[\alpha]$ is not finitely generated.

Proof. \Rightarrow : If α satisfies a monic polynomial of degree n , then $A[\alpha]$ is generated by $1, \alpha, \dots, \alpha^{n-1}$.

\Leftarrow : Suppose M is generated by v_1, \dots, v_n . Then we can find a matrix T with coefficients in A such that

$$\alpha \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} = T \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}.$$

Since $v_1, \dots, v_n \neq 0$, $\alpha I - T$ is singular, and $\det(\alpha I - T) = 0$. This gives a monic polynomial equation satisfied by α . □

Now for $\alpha, \beta \in L$ and let $M = A[\alpha]$ and $N = A[\beta]$. Note

1. if M, N are finitely generated by $\{\alpha_i\}$ and $\{\beta_j\}$, then MN is finitely generated by $\{\alpha_i\beta_j\}$.
2. $\alpha\beta MN \subseteq MN$ and $(\alpha + \beta)MN \subseteq MN$.

Hence $\alpha\beta$ and $\alpha + \beta$ are integral over A by Lemma 1.1.4 as needed. □

For the rest of this chapter, A is an integral domain, K is its fraction field, L is an extension of K , and B is the integral closure of A in L .

$$\begin{array}{ccc} L & \text{---} & B \\ | & & | \\ K & \text{---} & A \end{array} \tag{1.1}$$

Definition 1.1.6: A is **integrally closed** or **normal** if its integral closure in $K = \text{Frac}(A)$ is itself.

Proposition 1.1.7: If L is algebraic over K then every element of L can be written as $\frac{b}{a}$ where $b \in B$ and $a \in A$. Thus $L = \text{Frac}(B)$. In particular, for any extension L/\mathbb{Q} , $\text{Frac}(\mathcal{O}_L) = L$.

Proof. Given $\alpha \in L$, suppose that it satisfies the equation

$$P(x) := a_n x^n + a_{n-1} x^{n-1} + \cdots + a_0 = 0$$

with $a_0, \dots, a_n \in K$ and $a_n \neq 0$. Since $\text{Frac}(A) = K$, by multiplying by an element of A as necessary we may assume $a_0, \dots, a_n \in A$. Then

$$a_n^{n-1} P\left(\frac{x}{d}\right) := x^n + a_{n-1} x^{n-1} + a_n a_{n-2} x^{n-2} + \cdots + a_n^{n-1} a_0.$$

Hence $a_n \alpha$ is integral over A , i.e. $a_n \alpha \in B$. This shows α is in the desired form.

For the last part, take $K = \mathbb{Q}$ and $A = \mathbb{Z}$. □

For short we call (1.1) the “AKLB” setup if we further assume A is integrally closed in K . In the usual case, A is the integral closure of \mathbb{Z} in K . In this case, we write $A = \mathcal{O}_K$.¹

When $F = \overline{\mathbb{Q}}$, the algebraic closure of \mathbb{Q} , $a \in \overline{\mathbb{Q}}$ is called an algebraic number and $a \in \mathcal{O}_{\overline{\mathbb{Q}}}$ is an algebraic integer.

Theorem 1.1.8 (Rational Roots Theorem). *A UFD is integrally closed.*

Proof. Suppose R is a UFD with field of fractions K . Let $x \in K$ be integral over R ; suppose x satisfies

$$x^n + a_{n-1} x^{n-1} + \cdots + a_0 = 0$$

where $a_0, \dots, a_{n-1} \in R$. Write $x = \frac{p}{q}$ where $p, q \in R$ are relatively prime. Then multiplying the above by q^n gives

$$\begin{aligned} p^n + a_{n-1} p^{n-1} q + \cdots + a_1 p q^{n-1} + a_0 q^n &= 0 \\ q(a_{n-1} p^{n-1} + \cdots + a_0 q^{n-1}) &= -p^n \end{aligned}$$

Thus $q \mid p$, possible only if $q = 1$. This shows $x \in R$. □

Note that in the definition of integrality, an element is integral if it is the zero of *any* monic polynomial in $A[x]$. However, it suffices to check that its *minimal* polynomial is in $A[x]$.

Proposition 1.1.9: Let L be an algebraic extension of K and A be integrally closed. Then $\alpha \in L$ is integral over A iff its minimal polynomial f over K has coefficients in A .

¹Later on, when we take K to be an extension of the p -adic field \mathbb{Q}_p , we will use \mathcal{O}_K to denote the integral closure of \mathbb{Z}_p in K .

Proof. The reverse direction is clear. For the forward direction, note all zeros of f are integral over K since they satisfy the same polynomial equation that α satisfies. The coefficients of f are polynomial expressions in the roots so are integral over A , and hence in A (since they are already in K). \square

Proposition 1.1.10 (Finite generation):

1. Let $A \subseteq B \subseteq C$ be rings. If B is finitely generated as an A -module and C is finitely generated as a B -module, then C is finitely generated as an A -module.
2. If B is integral over A and finitely generated as an A -algebra, then it is finitely generated as an A -module.

Proof.

1. Take products of generators.
2. Let algebra generators be β_1, \dots, β_m . Then

$$A \subseteq A[\beta_1] \subseteq \dots \subseteq A[\beta_1, \dots, \beta_m]$$

is a chain of integral extensions, so item 2 follows from 1. \square

Combining this proposition with Lemma 1.1.4 we get the following:

Proposition 1.1.11 (Transitivity of integrality): Let $A \subseteq B \subseteq C$ be integral domains and K, L, M be their fraction fields.

1. If B is integral over A and C is integral over B , then C is integral over A .
2. Let A' be the integral closure of A over B and A'' be the integral closure of A' over C . Let A''' be the integral closure of A in C . Then $A''' = A''$.
3. The integral closure of A is integrally closed.

Proof.

1. For $\gamma \in C$, let b_i be the coefficients of the minimal polynomial of C over B . Then γ is integral over $A[b_0, \dots, b_m]$, so by Proposition 1.1.10, item 2, $A[b_0, \dots, b_m, \gamma]$ is finitely generated over A . Since $\gamma A[b_0, \dots, b_m, \gamma] \subseteq A[b_1, \dots, b_m, \gamma]$, by Lemma 1.1.4, γ is integral over A .
2. By item 1 applied to $A \subseteq A' \subseteq A''$, A'' is integral over A so $A'' \subseteq A'''$. Conversely, any element $a \in A'''$ is integral over A so *a fortiori* integral over A'' ; thus $A''' \subseteq A''$.
3. Follows from item 2 applied to $A = B = C$. \square

2 Norms and Traces

Let B be a free A -module of rank n . Then any element $\beta \in B$ defines an A -linear map m_β (or $[\beta]$), multiplication by β . It is helpful to think of β as a linear map because then we can apply results from linear algebra.

Definition 1.2.1: The trace, determinant, and characteristic polynomial of m_β are called the **trace**, **norm**, and **characteristic polynomial** of β .

These are computed by choosing any basis of e_1, \dots, e_n for B over A , and then computing the action of β on this basis.

Proposition 1.2.2 (Elementary properties): The following hold ($a \in A; \beta, \beta' \in B$):

1. $\text{tr}(\beta + \beta') = \text{tr}(\beta) + \text{tr}(\beta')$
2. $\text{tr}(a\beta) = a \text{tr}(\beta)$
3. $\text{tr}(a) = na$
4. $\text{Nm}(\beta\beta') = \text{Nm}(\beta) \cdot \text{Nm}(\beta')$
5. $\text{Nm}(a) = a^n$

Proposition 1.2.3 (Behavior with respect to field extensions): Suppose L/K is a degree n field extension, M is a finite extension of L , and $\beta \in L$.

1. (Relationship with roots of minimal polynomial) If $f(X)$ is the minimal polynomial of β over K and β_1, \dots, β_m are the roots of $f(X) = 0$ in a Galois closure of K , then letting $r = [L : K(\beta)] = \frac{n}{m}$,
 - (a) $\text{char}_{L/K}(\beta) = f(X)^r$
 - (b) $\text{tr}_{L/K}(\beta) = r(\beta_1 + \dots + \beta_m)$
 - (c) $\text{Nm}_{L/K}(\beta) = (\beta_1 \cdots \beta_m)^r$
2. (Relationship with embeddings) Suppose L is separable over K , M is a Galois extension of K , and $\sigma_1, \dots, \sigma_n$ are the set of distinct embeddings $L \rightarrow M$ fixing K . Then
 - (a) $\text{tr}_{L/K}(\beta) = \sigma_1(\beta) + \dots + \sigma_n(\beta)$
 - (b) $\text{Nm}_{L/K}(\beta) = \sigma_1(\beta) \cdots \sigma_n(\beta)$

In particular, this is true when $L = M$ is a Galois extension of K , and we can think of the σ_k as simply the elements of $G(L/K)$.

3. (Transitivity of trace and norm) Suppose $\beta \in M$ and M/K is separable.² Then

²The last condition is not necessary. TODO: Find a proof of the general case.

- (a) $\text{tr}_{M/K}(\beta) = \text{tr}_{L/K}(\text{tr}_{M/L}(\beta))$
 (b) $\text{Nm}_{M/K}(\beta) = \text{Nm}_{L/K}(\text{Nm}_{M/L}(\beta))$
4. (Integrality) Assume AKLB. If $\beta \in B$, then the coefficients of $\text{char}_{L/K}(\beta)$, and hence $\text{tr}_{L/K}(\beta)$ and $\text{Nm}_{L/K}(\beta)$, are integral over A . In particular, if A is integrally closed in L then they are in A .

Proof.

1. If $r = 1$, i.e. $K[\beta] = L$, then by the Cayley-Hamilton Theorem, $f(m_\beta) = 0$. Since $f(X)$ is irreducible, $f(X) \mid \text{char}_{L/K}(\beta)$. However, these are monic polynomials of the same degree so they are equal.

In the general case, take a basis x_i of $K[\beta]$ over K and a basis y_j of L over $K[\beta]$. Then $x_i y_j$ form a basis of L over K , and the matrix of m_β with respect to this basis is n copies of A . This proves (a), which implies the rest of the statements.

2. Let β_1, \dots, β_m be the conjugates of β . There are m distinct imbeddings $K(\beta) \rightarrow M$; they each take β to a different β_k . Each of these imbeddings extend to $r := [L : K(\beta)] = \frac{n}{m}$ imbeddings $L \rightarrow M$. Now use item 1.
3. Note that for any finite extensions $K \subseteq L \subseteq N$ with N Galois, an imbedding $L \hookrightarrow N$ fixing K can be extended to a K -automorphism on N , and so be considered an element of the set $G(N/K)/G(N/L)$.³

Let N be a Galois extension containing M . By item 2,

$$\begin{aligned} \text{tr}_{M/K}(\beta) &= \sum_{\sigma \in G(N/K)/G(N/M)} \sigma(\beta) \\ \text{tr}_{L/K}(\text{tr}_{M/L}(\beta)) &= \text{tr}_{L/K} \left(\sum_{\sigma \in G(N/L)/G(N/M)} \sigma(\beta) \right) \\ &= \sum_{\tau \in G(N/K)/G(N/L)} \sum_{\sigma \in G(N/L)/G(N/M)} \tau(\sigma(\beta)) \end{aligned}$$

where in the second sum we take arbitrary representatives $\tau \in G(N/K)$ and $\sigma \in G(N/L)$. These are equal because for any choice of these representatives,

$$\{\sigma \in G(N/K)/G(N/M)\} = \{\tau\sigma \mid \tau \in G(N/K)/G(N/L), \sigma \in G(N/L)/G(N/M)\}$$

when considered in $G(N/K)/G(N/M)$ (i.e. as imbeddings $M \hookrightarrow N$ fixing K). The same is true of the norm.

4. The minimal polynomial of α has coefficients in A , by Proposition 1.1.9. Hence the result follows from item 1. □

³Using the primitive element theorem, write $L = K(\beta)$. The imbeddings $L \rightarrow N$ are those taking β to a conjugate; there are $[L : K]$ imbeddings. But we know $G(N/K)/G(N/L) = [L : K]$, so all of the imbeddings must be extendable. We also use this fact (in addition to a counting argument) in the proof of 2.

3 Discriminant

Definition 1.3.1: If B is a ring and free A -module of rank m , and $\beta_1, \dots, \beta_m \in B$, then their **discriminant** is

$$D(\beta_1, \dots, \beta_m) = \det[\text{tr}_{B/A}(\beta_i \beta_j)]_{1 \leq i, j \leq m}.$$

Proposition 1.3.2: If the change of basis matrix from γ_i to β_i is T , then

$$D(\gamma_1, \dots, \gamma_m) = \det(T)^2 \cdot D(\beta_1, \dots, \beta_m).$$

Proof. Let M_1 and M_2 be the matrices of the bilinear form

$$(\alpha, \alpha') = \text{tr}_{B/A}(\alpha \alpha')$$

with respect to the bases $(\beta_1, \dots, \beta_m)$ and $(\gamma_1, \dots, \gamma_m)$, respectively. Then, using the change of basis formula for bilinear forms,

$$\begin{aligned} D(\beta_1, \dots, \beta_m) &= \det(M_1) \\ D(\gamma_1, \dots, \gamma_m) &= \det(M_2) \\ M_2 &= T^t M_1 T \\ \det(M_2) &= \det(T)^2 \cdot \det(M_1) \end{aligned}$$

from which the result follows. □

Consider the discriminant of an arbitrary basis of B over A . By the above fact, this is well-defined up to multiplication by the square of a unit. The residue in $A/(A^\times)^2$ is called the discriminant $\text{disc}(B/A)$. The discriminant also refers to the ideal of A this element generates.

Note $\text{disc}(B/A)$ can be thought of as the determinant of the matrix of the bilinear form $(\beta, \beta') = \text{tr}_{B/A}(\beta \beta')$.

Proposition 1.3.3 (Criterion for integral basis): Let $A \subseteq B$ be integral domains and B be a free A -module of rank m with $\text{disc}(B/A) \neq 0$. Then $\gamma_1, \dots, \gamma_m \in B$ form a basis for B as an A -module iff

$$(D(\gamma_1, \dots, \gamma_m)) = (\text{disc}(B/A))$$

as ideals.

Proof. Let β_i be a basis. If the change of basis matrix from γ_i to β_i is T , then by Proposition 1.3.2,

$$D(\gamma_1, \dots, \gamma_m) = \det(T)^2 \cdot D(\beta_1, \dots, \beta_m) = \det(T)^2 \text{disc}(B/A)$$

Now γ_i is basis iff T is invertible, iff $\det(T)$ is a unit, iff $(D(\gamma_1, \dots, \gamma_m)) = (\text{disc}(B/A))$. □

Proposition 1.3.4 (Discriminants and Field Extensions):

1. (Relationship with embeddings) Let L be separable finite over K of degree m , and $\sigma_1, \dots, \sigma_m$ be the embeddings of L into a Galois extension M fixing K . Then for any basis β_1, \dots, β_m of L over K ,

$$D(\beta_1, \dots, \beta_m) = \det(\sigma_i \beta_j)^2 \neq 0.$$

2. (Nondegeneracy of trace pairing) If B is free of rank m over A (with fraction fields K, L as above), then the pairing

$$(\beta, \beta') \mapsto \text{tr}(\beta\beta')$$

is a perfect K -bilinear pairing, and $\text{disc}(B/A) = \text{disc}(K/L) \neq 0$.

Here *perfect* means that the map $a \mapsto (b \mapsto (a, b))$ is an isomorphism $L \rightarrow L^*$, and similarly for $b \mapsto (a \mapsto (a, b))$. This is equivalent to saying that the bilinear form is nondegenerate.

Proof. (Unfinished) Use Proposition 1.2.3(1b), and that σ_k, \det are both multiplicative. Inequality follows from independence of characters:

Let G be a group, F a field. Then the homomorphisms $G \rightarrow F^\times$ are linearly independent. \square

Thus for K of degree m over \mathbb{Q} , we can talk of $\text{disc}(\mathcal{O}_K/\mathbb{Z})$.

A closely related quantity to the discriminant is the *different*.

Definition 1.3.5: Assume AKLB, and suppose L/K is a finite separable extension. The **codifferent** of B with respect to A is

$$B^* = \{y \in L \mid \text{tr}(xy) \in A \text{ for all } x \in B\}.$$

The **different** of B with respect to A is

$$\mathfrak{D}_{B/A} = (B^*)^{-1}.$$

In other words, it is the largest B -submodule satisfying $\text{tr}(E) \subseteq A$.

Remark 1.3.6: We will define the discriminant in general, when B is not necessarily a free A -module, in Chapter 9. The relationship between the two definitions is the following: Let \mathfrak{p} be an ideal in A . Then $A_{\mathfrak{p}}$ is a principal ideal domain (in fact, a DVR). Let $S = A - \mathfrak{p}$; then $S^{-1}B$ is free over $S^{-1}A$ by the structure theorem for modules. We have $(\text{disc}(S^{-1}B/S^{-1}A)) = (\mathfrak{p}A_{\mathfrak{p}})^{m(\mathfrak{p})}$ for some $m(\mathfrak{p})$. Then

$$\text{disc}(B/A) = \prod_{\mathfrak{p}} \mathfrak{p}^{m(\mathfrak{p})}.$$

4 Integral bases

Proposition 1.4.1 (Finite generation of integral extensions): Let A be integrally closed and L separable of degree m over K . There are free finite A -submodules M and M' of L such that $M \subseteq B \subseteq M'$. B is a finitely generated A -module if A is Noetherian, and free of rank m if A is a PID.⁴

Proof. Let $\{\beta_1, \dots, \beta_m\} \subseteq B$ be a basis for L over K . Take a basis β'_i so that $\text{tr}(\beta_i \beta'_j) = \delta_{ij}$. Then

$$A\beta_1 + \dots + A\beta_m \subseteq B \subseteq A\beta'_1 + \dots + A\beta'_m.$$

The second inclusion follows because if $\beta \in B$, then writing $\beta = \sum_j b_j \beta'_j$, we have that $b_i = \text{tr}(\beta \beta_i) \in A$. (In other words, the β'_i form a basis for the codifferent B^* , which contains B .)

If A is Noetherian, then M' is finitely generated, so its submodule B is finitely generated over A . If A is a PID, then by the Structure Theorem for Modules (over PIDs), M is a direct sum of cyclic modules and a free module. Since it is contained in a free module of rank m and contains a free module of rank m , it must be free of rank m . \square

The following is immediate:

Theorem 1.4.2. *If K is finite over \mathbb{Q} (i.e. a number field), then \mathcal{O}_K is a finitely generated \mathbb{Z} -module. It is the largest subring that is finitely generated over \mathbb{Z} .*

Definition 1.4.3: A basis for \mathcal{O}_K as a \mathbb{Z} -module is called an **integral basis**.

Proposition 1.4.4: Suppose K has characteristic 0 (so L separable over K), $L = K[\beta]$, and f is the minimal polynomial of β over K . Let $f(X) = \prod (X - \beta_i)$ in the Galois closure of L . Then

$$D(1, \beta, \dots, \beta^{m-1}) = \prod_{1 \leq i < j \leq m} (\beta_i - \beta_j)^2 = (-1)^{m(m-1)/2} \cdot \text{Nm}_{L/K}(f'(\beta)).$$

This is called the discriminant of f .⁵

Proof. Note the β_i are conjugates of β ; assume $\beta = \beta_1$.

By Proposition 1.3.4, we have

$$D(1, \beta, \dots, \beta^{m-1}) = \left| \begin{array}{cccc} 1 & \beta_1 & \cdots & \beta_1^{m-1} \\ 1 & \beta_2 & \cdots & \beta_2^{m-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \beta_m & \cdots & \beta_m^{m-1} \end{array} \right|^2 = \prod_{1 \leq i < j \leq m} (\beta_i - \beta_j)^2,$$

⁴Alternative proof: proceed as in 5.8.

⁵This gives an alternative proof of the perfect pairing.

where the last statement follows by evaluating the Vandermonde determinant.

For the second equality, note by Proposition 1.2.3(1c) that

$$\begin{aligned} \text{Nm}_{L/K}(f'(\beta)) &= \text{Nm}_{L/K}((\beta_1 - \beta_2) \cdots (\beta_1 - \beta_m)) = \prod_{1 \leq i \leq m} \prod_{1 \leq j \leq m, j \neq i} (\beta_i - \beta_j) \\ &= (-1)^{\frac{m(m-1)}{2}} \prod_{1 \leq i < j \leq m} (\beta_i - \beta_j)^2. \end{aligned}$$

□

Proposition 1.4.5: If $K = \mathbb{Q}[\alpha]$, $\alpha \in \mathcal{O}_K$, and $D(1, \alpha, \dots, \alpha^{m-1}) = \text{disc}(\mathcal{O}/\mathbb{Z})$ then $\{1, \alpha, \dots, \alpha^{m-1}\}$ is an integral basis.

Proof. Using change-of-basis and the correspondence between index and determinant,

$$D(1, \alpha, \dots, \alpha^{m-1}) = \text{disc}(\mathcal{O}_K/\mathbb{Z}) \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^2.$$

Now $\text{disc}(\mathcal{O}_K/\mathbb{Z}) \in \mathbb{Z}$ so $[\mathcal{O}_K : \mathbb{Z}[\alpha]] = 1$.

□

Theorem 1.4.6 (Stickelberger's Theorem).

1. Let s is the number of complex (nonreal) embeddings $K \rightarrow \mathbb{C}$. Then

$$\text{sign}[\text{disc}(K/\mathbb{Q})] = (-1)^{s/2}.$$

2. $\text{disc}(\mathcal{O}_K/\mathbb{Z}) \equiv 0$ or $1 \pmod{4}$.

Proof. 1. Write $K = \mathbb{Q}[\alpha]$ by the Primitive Element Theorem and $\alpha_1, \dots, \alpha_r$ be the real conjugates and $\beta_1, \overline{\beta}_1, \dots, \beta_s, \overline{\beta}_s$ be the complex conjugates. By Proposition 1.4.4,

$$\text{sign}(D(1, \alpha, \dots, \alpha^{m-1})) = \text{sign} \left(\prod_{1 \leq j \leq s} (\beta_j - \overline{\beta}_j)^2 \right) = \prod_{1 \leq j \leq s} i^2 = (-1)^{s/2}.$$

2. Let $\alpha_1, \dots, \alpha_m$ be an integral basis. Let P and $-N$ be the sum of the terms in the expansion of $\det(\sigma_i \alpha_j)$ corresponding to even and odd permutations, respectively:

$$\begin{aligned} P &= \sum_{\text{even } \pi \in S_m} \prod_{i=1}^m \sigma_i \alpha_{\pi(i)} \\ N &= \sum_{\text{odd } \pi \in S_m} \prod_{i=1}^m \sigma_i \alpha_{\pi(i)}. \end{aligned}$$

Then

$$\begin{aligned} \text{disc}(\mathcal{O}_K/\mathbb{Z}) &= \det(\sigma_i \alpha_j)^2 \\ &= (P - N)^2 \\ &= (P + N)^2 - 4PN. \end{aligned}$$

Take $\sigma \in G(K^{\text{gal}}/\mathbb{Q})$. Note composition by σ permutes the σ_i , say by ν . Then

$$P = \sum_{\text{even } \pi \in S_m} \prod_{i=1}^m \sigma_i \alpha_{\nu^{-1}\pi(i)}$$

$$N = \sum_{\text{odd } \pi \in S_m} \prod_{i=1}^m \sigma_i \alpha_{\nu^{-1}\pi(i)}$$

and hence σ permutes $\{P, N\}$. Hence σ fixes $P + N, PN$ and they are rational. Since they are integral over \mathbb{Z} they are integers. Thus the above is congruent to 0 or 1 modulo 4. □

Example 1.4.7 (Quadratic extensions): Any quadratic extension of \mathbb{Q} is in the form $\mathbb{Q}(\sqrt{m})$ for some squarefree integer m . We find the ring of integers of $\mathbb{Q}(\sqrt{m})$. Consider two cases.

1. $m \equiv 2, 3 \pmod{4}$: The minimal polynomial of \sqrt{m} is $X^2 - m$, so

$$\text{disc}(1, \sqrt{m}) = (\sqrt{m} - (-\sqrt{m}))^2 = 4m.$$

Note $\frac{\text{disc}(1, \sqrt{m})}{\text{disc}(\mathbb{Q}(\sqrt{m})/\mathbb{Q})}$ must be a square by Proposition 1.3.2 so $\text{disc}(\mathbb{Q}(\sqrt{m})/\mathbb{Q})$ equals m or $4m$. However, by Stickelberger's Theorem, $\text{disc}(\mathbb{Q}(\sqrt{m})/\mathbb{Q}) \equiv 0, 1 \pmod{4}$. Hence $\text{disc}(\mathbb{Q}(\sqrt{m})/\mathbb{Q}) \neq m$ and $\text{disc}(\mathbb{Q}(\sqrt{m})/\mathbb{Q}) = 4m$. By Proposition 1.3.3, $1, \sqrt{m}$ is an integral basis.

2. $m \equiv 1 \pmod{4}$: Note $\frac{1+\sqrt{m}}{2}$ is integral with minimal polynomial $X^2 - X - \frac{m-1}{4}$, so

$$\text{disc}\left(1, \frac{1+\sqrt{m}}{2}\right) = \left(\frac{1+\sqrt{m}}{2} - \frac{1-\sqrt{m}}{2}\right)^2 = m.$$

Since m is squarefree, $\text{disc}(\mathbb{Q}(\sqrt{m})/\mathbb{Q}) = m$ and Proposition 1.3.3 says $1, \frac{1+\sqrt{m}}{2}$ is an integral basis.

The following tells us about integral bases for products of fields.

Proposition 1.4.8: Suppose that K, L are field extensions of \mathbb{Q} such that

$$[KL : \mathbb{Q}] = [K : \mathbb{Q}][L : \mathbb{Q}].$$

Let $d = \gcd(\text{disc}(K/\mathbb{Q}), \text{disc}(L/\mathbb{Q}))$. Then

1. $\mathcal{O}_K \subseteq d^{-1}\mathcal{O}_K\mathcal{O}_L$.
2. If $\mathcal{O}_{KL} = \mathcal{O}_K\mathcal{O}_L$, then $\text{disc}(KL/\mathbb{Q}) = \text{disc}(K/\mathbb{Q})^{[L:\mathbb{Q}]} \text{disc}(L/\mathbb{Q})^{[K:\mathbb{Q}]}$.

Proof. Let $\{\alpha_1, \dots, \alpha_m\}$ be an integral basis for K and $\{\beta_1, \dots, \beta_n\}$ be an integral basis for L . By the degree assumption, we know that $\{\alpha_i \beta_j\}$ is a basis for KL over \mathbb{Q} . Any element of KL integral over \mathbb{Q} can be written as

$$\gamma = \sum_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} \frac{a_{ij}}{r} \alpha_i \beta_j \quad (1.2)$$

where $\gcd(r, \gcd(a_{ij})) = 1$.

We need to show that $r \mid d$. Let $x_i = \sum_{j=1}^n \frac{a_{ij}}{r} \beta_j$. We will turn (1.2) into a system of equations by considering all embeddings $K \hookrightarrow \mathbb{C}$, solve for the x_i using Cramer's rule, and in this way show that each x_i is an algebraic integer in L divided by a bounded denominator.

Note given embeddings $\sigma_K : K \hookrightarrow \mathbb{C}$ and $\sigma_L : L \hookrightarrow \mathbb{C}$, there is exactly one embedding $\sigma_{KL} : KL \hookrightarrow \mathbb{C}$ such that restricts to σ_K and σ_L . It is clearly unique if it exists. To show existence, write $K = \mathbb{Q}(\alpha) \cong \mathbb{Q}(x)/(f(x))$ by PET, and note that the characteristic polynomial of f does not change upon passing to L because of the degree assumption. Hence $KL = L(\alpha) = L(x)/(f(x))$, and in extending σ_L to σ_{KL} , we are allowed to send $\alpha = x$ to $\sigma_L(\alpha)$.

Fix an embedding $\sigma : L \hookrightarrow \mathbb{C}$, and let $\sigma_1, \dots, \sigma_m$ be all embeddings $K \hookrightarrow \mathbb{C}$. Then applying σ_k to 1.2 we obtain the system of equations

$$\sum_{i=1}^m \sigma_k(\alpha_i) x_i = \sigma_k(\gamma), \quad 1 \leq k \leq m.$$

By Cramer's rule, letting $D = \det[(\sigma_k(\alpha_i))_{k,i}]$ we get $Dx_i = D_i$ where D_i has the i th column of D replaced by $(\sigma_k(\alpha_i))_{k=1}^m$. Note that D and D_i are both algebraic integers. Using $\text{disc}(\mathcal{O}_K/\mathbb{Z}) = D^2$ (Proposition 1.3.4), we get

$$\text{disc}(\mathcal{O}_K/\mathbb{Z})x_i = DD_i.$$

Hence $\text{disc}(\mathcal{O}_K/\mathbb{Z})x_i$ is an algebraic integer (in \mathcal{O}_L). Since the β_j are an integral basis for \mathcal{O}_L , this forces $r \mid \text{disc}(\mathcal{O}_K/\mathbb{Z})$. Similarly, $r \mid \text{disc}(\mathcal{O}_L/\mathbb{Z})$, as needed.

Now we prove the second part. Choose $(\alpha_1, \dots, \alpha_m)$ a basis for K/\mathbb{Q} and $(\beta_1, \dots, \beta_n)$ a basis for L/\mathbb{Q} . Then $(\alpha_j \beta_k)_{1 \leq j \leq m, 1 \leq k \leq n}$ is a basis for KL/\mathbb{Q} . For $\gamma \in KL$, let $(\gamma)_{j,k}$ denote the coordinate of $\alpha_j \beta_k$ in γ . Then the $mn \times mn$ matrix

$$\begin{aligned} [\text{tr}(\alpha_{i_1} \beta_{i_2} \alpha_{i'_1} \beta_{i'_2})] &= \left[\sum_{1 \leq j \leq m, 1 \leq k \leq n} (\alpha_{i_1} \beta_{i_2} \alpha_{i'_1} \beta_{i'_2} \alpha_j \beta_k)_{j,k} \right] \\ &= \left[\sum_{1 \leq j \leq m, 1 \leq k \leq n} (\alpha_{i_1} \alpha_{i'_1} \alpha_j)_j (\beta_{i_2} \beta_{i'_2} \beta_k)_k \right] \\ &= \left[\sum_{1 \leq j \leq m} \sum_{1 \leq k \leq n} (\alpha_{i_1} \alpha_{i'_1} \alpha_j)_j (\beta_{i_2} \beta_{i'_2} \beta_k)_k \right] \\ &= [\text{tr}(\alpha_{i_1} \alpha_{i'_1})] \otimes [\text{tr}(\beta_{i_2} \beta_{i'_2})]. \end{aligned}$$

Taking determinants and using

$$\det(A \otimes B) = \det(A)^n \det(B)^m, \quad A \in M_{m \times m}, B \in M_{n \times n}$$

we get

$$\text{disc}(KL/\mathbb{Q}) = \text{disc}(K/\mathbb{Q})^{[L:\mathbb{Q}]} \text{disc}(L/\mathbb{Q})^{[M:\mathbb{Q}]}.$$

□

5 Problems

1. Suppose that $f \in \mathbb{Z}[x]$ is irreducible and has a root of absolute value at least $\frac{3}{2}$. Prove that if α is a root of f then $f(\alpha^3 + 1) \neq 0$.
2. Let a_1, \dots, a_n be algebraic integers with degrees d_1, \dots, d_n . Let a'_1, \dots, a'_n be the conjugates of a_1, \dots, a_n with greatest absolute value. Let c_1, \dots, c_n be integers. Prove that if the LHS of the following expression is not zero, then

$$|c_1 a_1 + \dots + c_n a_n| \geq \left(\frac{1}{|c_1 a'_1| + \dots + |c_n a'_n|} \right)^{d_1 d_2 \dots d_n - 1}.$$

For example,

$$|c_1 + c_2 \sqrt{2} + c_3 \sqrt{3}| \geq \left(\frac{1}{|c_1| + |2c_2| + |2c_3|} \right)^3.$$

3. Let p be a prime and consider k p th roots of unity whose sum is not 0. Prove that the absolute value of their sum is at least $\frac{1}{k^{p-2}}$.

Chapter 2

Ideal factorization

1 Discrete Valuation Rings

Definition 2.1.1: Let K be a field. A **discrete valuation** on K is a surjective function $v : K^\times \rightarrow \mathbb{Z}$ such that for every $x, y \in K^\times$,

1. v is a group homomorphism: $v(xy) = v(x)v(y)$.
2. $v(x + y) \geq \min(v(x), v(y))$.

We set $v(0) = \infty$.

A **discrete valuation ring** (over \mathbb{Z}) is a local integral domain R (not a field), whose fraction field has a discrete valuation v .

An element t with $v(t) = 1$ is a **uniformizing parameter**.

Proposition 2.1.2: Suppose R is a DVR with fraction field K . Let v be the valuation on K .

1. The units are exactly the elements with 0 valuation:

$$R^\times = v^{-1}(0).$$

2. Its maximal ideal is the set of elements with positive valuation.

$$\mathfrak{m} = \{x : v(x) > 0\}.$$

3. R is a PID with ideals $\mathfrak{m}^n = \{x : v(x) \geq n\} = (t^n)$ for $n \in \mathbb{N}$.

4. R is a UFD; any element can be written uniquely in the form ut^n where u is a unit.

Lemma 2.1.3. *Let A be a local domain with maximal ideal \mathfrak{m} principal and nonzero. If $\bigcap_{n \geq 0} \mathfrak{m}^n = 0$ then A is a DVR.*

Theorem 2.1.4. *Let (A, \mathfrak{m}) be a Noetherian local domain. The following conditions are equivalent.*

1. A is a DVR.
2. A is a normal domain of dimension 1. (Dimension 1 means that the longest chain of prime ideals is 2: $\mathfrak{p}_0 \subseteq \mathfrak{p}_1$.) (Since A is local this means it has only two prime ideals.)
3. A is a normal domain of depth 1. (There is a nonzero $x \in A$ with $\mathfrak{m} \in \text{Ass}(A/\langle x \rangle)$.)
4. A is a regular local ring of dimension 1. (Regular means its maximal ideal is generated by a number of elements equal to its dimension. So here it means \mathfrak{m} is principal.)
5. \mathfrak{m} is principal and nonzero.

Proof. Note (5) \implies (1) uses Krull Intersection Theorem: For R a Noetherian ring, \mathfrak{a} an ideal, and M a finitely generated module (esp. when $M = R$), then there exists $x \in \mathfrak{a}$ such that

$$(1+x) \bigcap_{n=0}^{\infty} \mathfrak{a}^n M = 0.$$

□

2 Dedekind Domains

Definition 2.2.1: A **Dedekind domain** is a normal Noetherian integral domain A such that every nonzero prime ideal is maximal.

Proposition 2.2.2: A local integral domain is Dedekind iff it is a DVR.

Proposition 2.2.3: For every nonzero prime ideal \mathfrak{p} in a Dedekind domain A , the localization $A_{\mathfrak{p}}$ is a DVR. (*Locally, Dedekind domains are DVR's.*)

(The converse, i.e. if $A_{\mathfrak{p}}$ is a DVR for every \mathfrak{p} , then A is Dedekind, holds using Serre's criterion.)

Theorem 2.2.4 (Unique factorization of prime ideals). *Let A be a Dedekind domain. Every proper nonzero ideal of A can be written uniquely as a product of prime ideals.*

Proof. Let \mathfrak{a} be a proper nonzero ideal of A .

1. If A is Noetherian, then every ideal $\mathfrak{a} \subseteq A$ contains a product $\mathfrak{b} = \prod \mathfrak{p}_k^{r_k}$ of nonzero prime ideals: Otherwise, choose a maximal counterexample \mathfrak{a} (possible since A is Noetherian). Since \mathfrak{a} is not prime, there exist $x, y \notin \mathfrak{a}$ such that $xy \in \mathfrak{a}$. By the maximality assumption both $\mathfrak{a} + (x)$ and $\mathfrak{a} + (y)$ contain a product of prime ideals, and so does $\mathfrak{a} \supseteq (\mathfrak{a} + (x))(\mathfrak{a} + (y))$.

2. By the Chinese Remainder Theorem

$$A/\mathfrak{b} \cong \prod_k A/\mathfrak{p}_k^{r_k}$$

via the natural map.

3. If \mathfrak{p} is a maximal ideal in a ring A , and $\mathfrak{q} = \mathfrak{p}A_{\mathfrak{p}}$, then the natural map $A/\mathfrak{p}^m \rightarrow (A/\mathfrak{p}^m)_{\mathfrak{p}} = A_{\mathfrak{p}}/\mathfrak{q}^m$ is an isomorphism. (Indeed, it is injective because \mathfrak{p} is prime and surjective because any $s \in A - \mathfrak{p}$ is invertible modulo \mathfrak{p}^m , on account of $(s) + \mathfrak{p}^m = A$.) Thus

$$\prod_k A/\mathfrak{p}_k^{r_k} \cong \prod_k A_{\mathfrak{p}_k}/\mathfrak{q}_k^{r_k}.$$

(This is where we use the fact that nonzero prime ideals are maximal.)

4. Combining the above, we get a one-to-one correspondence between ideals in A containing \mathfrak{b} , and ideals in $\prod_k A_{\mathfrak{p}_k}/\mathfrak{q}_k^{r_k}$. All ideals in the last ring are in the form $\prod_k \mathfrak{q}_k^{s_k}/\mathfrak{q}_k^{r_k}$, so \mathfrak{a} is of the form $\prod_k \mathfrak{q}_k^{s_k}$. Moreover, different prime ideals containing \mathfrak{b} correspond to different $\prod_k \mathfrak{q}_k^{s_k}/\mathfrak{q}_k^{r_k}$, which are different for different s_k , giving uniqueness. \square

Corollary 2.2.5. *Let A be a Dedekind domain.*

1. If $\mathfrak{a} = \prod_{\mathfrak{p}} \mathfrak{p}_k^{r_k}$ and $\mathfrak{b} = \prod_{\mathfrak{p}} \mathfrak{p}_k^{s_k}$ are ideals in A and \mathfrak{p} is a nonzero prime ideal then

$$\begin{aligned} \mathfrak{a} \supseteq \mathfrak{b} &\iff r_k \geq s_k \text{ for all } k \\ &\iff \mathfrak{a}A_{\mathfrak{p}} \supseteq \mathfrak{b}A_{\mathfrak{p}} \text{ for all } \mathfrak{p}. \end{aligned}$$

2. If $\mathfrak{a} \supset \mathfrak{b} \neq 0$ are ideals in A then $\mathfrak{a} = \mathfrak{b} + (a)$ for some $a \in A$. In particular, if $b \in \mathfrak{a}$ then there exists $a \in A$ such that $\mathfrak{a} = (a, b)$; i.e. each ideal is generated by at most two elements.

3. (Inverses) Let $\mathfrak{a} \neq 0$ be an ideal of A . There exists a nonzero ideal \mathfrak{a}^* such that $\mathfrak{a}\mathfrak{a}^*$ is principal.

(a) We can choose \mathfrak{a}^* so $\mathfrak{a}\mathfrak{a}^* = (a)$ for given $a \in \mathfrak{a}$.

(b) Alternatively we can choose \mathfrak{a}^* to be relatively prime to a given ideal $\mathfrak{c} \neq 0$.

Proof. 1. The forward direction was shown in the course of the theorem. The reverse directions are easy.

2. Choose any $a \in \mathfrak{a} \setminus \{0\}$. By unique factorization, we can write

$$\begin{aligned} (a) &= \mathfrak{p}_1^{u_1} \cdots \mathfrak{p}_r^{u_r} \\ \mathfrak{a} &= \mathfrak{p}_1^{v_1} \cdots \mathfrak{p}_r^{v_r} \end{aligned}$$

for primes $\mathfrak{p}_1, \dots, \mathfrak{p}_r$ and $u_j \geq v_j \geq 0$. Now choose $b_j \in \mathfrak{p}_j^{v_j} \setminus \mathfrak{p}_j^{v_j+1}$. By the Chinese remainder theorem we can choose b such that $b \equiv b_j \pmod{\mathfrak{p}_j^{v_j+1}}$ for all j . Since

$\text{ord}_{\mathfrak{p}_j}(b_j) = v_j$, by item 1, the highest power of \mathfrak{p}_j dividing (b) is v_j . The highest power of \mathfrak{p}_j dividing (a) is $u_j \geq v_j$, so the highest power of \mathfrak{p}_j dividing (a, b) is v_j . Now for a prime $\mathfrak{q} \notin \{\mathfrak{p}_1, \dots, \mathfrak{p}_r\}$, we have $a \notin \mathfrak{q}$ (else \mathfrak{q} would divide \mathfrak{a}), so \mathfrak{q} does not divide (a, b) . We conclude

$$(a, b) = \mathfrak{p}_1^{v_1} \cdots \mathfrak{p}_r^{v_r},$$

as needed.

3. (a) follows from item 1; for (b), use item 2 and 3(a) to write $\mathfrak{a} = \mathfrak{a}\mathfrak{c} + (a) = \mathfrak{a}\mathfrak{c} + \mathfrak{a}\mathfrak{a}^* = \mathfrak{a}(\mathfrak{c} + \mathfrak{a}^*)$. □

Theorem 2.2.6. *Assume AKLB, and K/L is finite separable. If A is a Dedekind domain, then so is B . In particular, taking $A = \mathbb{Z}$ and $K = \mathbb{Q}$, every ring of integers in a finite separable extension of \mathbb{Q} is Dedekind.*

Proof.

1. B is noetherian: By Proposition 1.4.1, B is a finitely generated A -module, hence a Noetherian A -module, hence Noetherian as a ring.
2. B is integrally closed by Proposition 1.1.11(2).
3. Every nonzero prime ideal \mathfrak{q} of B is maximal: Take a nonzero $\beta \in \mathfrak{q}$ and let its minimal polynomial be $x^n + a_{n-1}x^{n-1} + \cdots + a_n$. Then $a_n = -\beta^n - \cdots - a_1\beta \in \beta B \cap A \subseteq \mathfrak{q} \cap A$. This shows $\mathfrak{q} \cap A \neq 0$; since A is Dedekind and $\mathfrak{q} \cap A$ is prime, $\mathfrak{q} \cap A$ is maximal and A/\mathfrak{q} is a field. Since B is integral over A , B/\mathfrak{q} is integral over A/\mathfrak{q} .

Lemma 2.2.7. *An integral domain B containing a field k and algebraic over k is a field.*

Proof. Let $\beta \in B$ be nonzero. Then $k[\beta]$ is a finite dimensional vector space and the multiplication-by- β map $m_\beta : k[\beta] \rightarrow k[\beta]$ is injective, hence surjective. Thus there exists β' so $\beta\beta' = 1$, i.e. β has an inverse. □

The lemma shows B/\mathfrak{q} is a field. Hence \mathfrak{q} is maximal.

Alternatively, this follows directly from “lying-over” and “going up” for integral extensions. □

Theorem 2.2.8. *Suppose K is a finite extension of \mathbb{Q} . Then unique factorization of ideals holds in \mathcal{O}_K .*

Proof. Combine Theorem 2.2.4 and Theorem 2.2.6. □

3 Primary decomposition*

[ADD: Commutative algebra generalization, and a new proof of unique ideal factorization]

4 Ideal class group

Let A be a Dedekind domain with fraction field K .

Definition 2.4.1: A **fractional ideal** of A is a nonzero A -submodule of K such that $d\mathfrak{a} \in A$ for some $d \in A$.

A principal fractional ideal is one of the form

$$(b) := bA := \{ba \mid a \in A\}.$$

The product of two fractional ideals is

$$\mathfrak{a}\mathfrak{b} = \left\{ \sum a_i b_i \mid a_i \in \mathfrak{a}, b_i \in \mathfrak{b} \right\}.$$

Note that given a nonzero A -submodule of K , it is finitely generated iff it is a fractional ideal. (Take common denominators of the generators.)

We can extend unique factorization to fractional ideals, in the same way that we can extend unique factorization from \mathbb{Z} to \mathbb{Q} .

Theorem 2.4.2. *The set $\text{Id}(A)$ of fractional ideals is a free abelian group on the set of prime ideals. Thus each fraction ideal can be uniquely written in the form*

$$\mathfrak{a} = \prod_{\mathfrak{p}} \mathfrak{p}^{r_{\mathfrak{p}}}.$$

Proof. Freeness follows from unique factorization (Theorem 2.2.4) and existence of inverses follows from Corollary 2.2.5(3a). \square

Now we are ready for the following definition.

Definition 2.4.3: Let $P(A)$ be the group of principal ideals of A . The **ideal class group** $\text{Cl}(A)$ is $\text{Id}(A)/P(A)$. Its order is the **class number**.

The ideal class group and class number of K are defined as the ideal class group and class number of \mathcal{O}_K .

Note that we have an exact sequence

$$0 \rightarrow P(A) \rightarrow \text{Id}(A) \rightarrow \text{Cl}(A) \rightarrow 0.$$

The class number is 1 iff all A is a PID. Thus in some sense it measures how far A is from being a PID.

Alternatively there is an exact sequence

$$1 \rightarrow \mathcal{O}_K^\times \rightarrow K^\times \rightarrow I_K \rightarrow \text{Cl}_K \rightarrow 1$$

where the map $K^\times \rightarrow I_K$ is given by $a \mapsto (a)$.

Theorem 2.4.4 (Approximation Theorem). *Let $x_1, \dots, x_m \in A$, and $\mathfrak{p}_1, \dots, \mathfrak{p}_m$ be distinct prime ideals. There is $x \in A$ such that*

$$\text{ord}_{\mathfrak{p}_i}(x - x_i) > n$$

for all i .

Proof. Immediate from the Chinese Remainder Theorem. □

5 Factorization in extensions

Assume AKLB, with A Dedekind and L/K finite separable. A prime ideal $\mathfrak{p} \subset A$ will factor in B :

$$\mathfrak{p}B = \mathfrak{P}_1^{e_1} \cdots \mathfrak{P}_g^{e_g}.$$

We say e_i is the **ramification index** of \mathfrak{P}_i . For $\mathfrak{P} \mid \mathfrak{p}$, we write $e(\mathfrak{P}/\mathfrak{p})$ for the ramification index and $f(\mathfrak{P}/\mathfrak{p})$ for the **residue class degree** $[B/\mathfrak{P} : A/\mathfrak{p}]$.

1. If $e_k > 1$ for some k , \mathfrak{p} is **ramified** in B .
 - (a) If $g = 1$ and $e_1 > 1$, \mathfrak{p} is **totally ramified**.
 - (b) When $|A/\mathfrak{p}| = p^n$, p prime, and $p \nmid [B/\mathfrak{P} : A/\mathfrak{p}]$, then \mathfrak{p} is **tamely ramified**.
2. If $e_i = f_i = 1$ for all i , \mathfrak{p} **splits completely**.
3. If $\mathfrak{p}B$ stays prime, \mathfrak{p} is **inert**.

Lemma 2.5.1. *A prime ideal \mathfrak{P} divides \mathfrak{p} iff $\mathfrak{P} \cap K = \mathfrak{p}$.*

Theorem 2.5.2 (Degree equation). *Let $m = [L : K]$ and suppose $\mathfrak{p}B = \mathfrak{P}_1^{e_1} \cdots \mathfrak{P}_g^{e_g}$. Then*

$$\sum_{i=1}^g e_i f_i = m.$$

If L/K is Galois, then all the e_i are equal and all the f_i are equal. Letting e and f denote these common values,

$$efg = m.$$

Proof. We show both sides of the equation equal $\dim_{A/\mathfrak{p}}(B/\mathfrak{p}B)$.

For the LHS, by the Chinese Remainder Theorem $B/\mathfrak{p}B \cong \prod_{i=1}^g B/\mathfrak{P}_i^{e_i}$ so

$$\dim_{A/\mathfrak{p}}(B/\mathfrak{p}B) = \sum_{i=1}^g \dim_{A/\mathfrak{p}}(B/\mathfrak{P}_i^{e_i}). \quad (2.1)$$

Consider the filtration

$$B \supset \mathfrak{P}_i \supset \cdots \supset \mathfrak{P}_i^{e_i}.$$

There are no ideals between any two consecutive ideals by Corollary 2.2.5 (the first iff), so there are no proper B/\mathfrak{P}_i -ideals (i.e. subspaces) of $\mathfrak{P}_i^r/\mathfrak{P}_i^{r+1}$. Hence $\dim_{B/\mathfrak{P}_i}(\mathfrak{P}_i^r/\mathfrak{P}_i^{r+1}) = 1$ and $\dim_{A/\mathfrak{p}}(\mathfrak{P}_i^r/\mathfrak{P}_i^{r+1}) = f_i$. Thus

$$\dim_{A/\mathfrak{p}}(B/\mathfrak{P}_i^{e_i}) = e_i f_i. \quad (2.2)$$

Combining (2.1) and (2.2) give

$$\dim_{A/\mathfrak{p}}(B/\mathfrak{p}B) = \sum_{i=1}^g e_i f_i.$$

For the RHS, let $A' = (A - \mathfrak{p})^{-1}A = A_{\mathfrak{p}}$ and $B' = (A - \mathfrak{p})^{-1}B$. First note that

$$A/\mathfrak{p} = \text{Frac}(A/\mathfrak{p}) \cong (A/\mathfrak{p})_{\mathfrak{p}} = A'/\mathfrak{p}A'$$

and

$$B/\mathfrak{p} \stackrel{(*)}{\cong} (A - \mathfrak{p})^{-1}(B/\mathfrak{p}B) = B'/\mathfrak{p}B',$$

where in (*) we use the fact that all elements of $A - \mathfrak{p}$ are invertible modulo $\mathfrak{p}B$, on account of A/\mathfrak{p} being a field. Note A' is a DVR and hence a PID. Since B is finitely generated over A , and localization is exact, B' is finitely generated over A' . Furthermore, B' is A' -torsion free. The previous three statements along with the Structure Theorem for Modules gives that $B' \cong A'^n$ (as A' -modules) for some n . Perform the following operations:

$$\begin{array}{ccc} & B' \cong A'^n & \\ \swarrow \otimes K & & \searrow \bullet/\mathfrak{p}\bullet \\ K \cong L^n & & B'/\mathfrak{p}B' \cong (A'/\mathfrak{p}A')^n \\ & & \parallel \\ & & B/\mathfrak{p}B \cong (A/\mathfrak{p})^n \end{array}$$

Hence

$$[L : K] = n = \dim_{A/\mathfrak{p}} B/\mathfrak{p}B$$

as needed.

Now suppose L/K is Galois. Then $G(L/K)$ permutes the primes \mathfrak{P} dividing \mathfrak{p} . Since $e(\mathfrak{P}/\mathfrak{p}) = e(\sigma\mathfrak{P}/\mathfrak{p})$ and $f(\mathfrak{P}/\mathfrak{p}) = f(\sigma\mathfrak{P}/\mathfrak{p})$, it suffices to show $G(L/K)$ acts transitively.

Suppose by way of contradiction that \mathfrak{P} and \mathfrak{Q} are not in the same orbit. By the Chinese Remainder Theorem there exists $\beta \in \mathfrak{Q} - \{\sigma\mathfrak{P} \mid \sigma \in G(L/K)\}$. Now

$$\text{Nm}_{L/K}(\beta) = \prod_{\sigma \in G(L/K)} \sigma(\beta) \in \mathfrak{Q} \cap A = \mathfrak{p} \subseteq \mathfrak{P},$$

the first because $\beta \in \mathfrak{Q}$ and the second because $\beta \in B$ is integral over A (which is integrally closed in K). But $\sigma(\beta) \notin \mathfrak{P}$ so

$$\prod_{\sigma \in G(L/K)} \sigma(\beta) \notin \mathfrak{P},$$

a contradiction. □

Note that the ramification indices and residue degrees multiply under field extension.

Proposition 2.5.3: Suppose that M/L and L/K are finite separable extensions (with Dedekind ring of integers), and that $\mathfrak{Q} \mid \mathfrak{P} \mid \mathfrak{p}$ are primes in M, L, K respectively. Then

$$\begin{aligned} e(\mathfrak{Q}/\mathfrak{p}) &= e(\mathfrak{Q}/\mathfrak{P})e(\mathfrak{P}/\mathfrak{p}) \\ f(\mathfrak{Q}/\mathfrak{p}) &= f(\mathfrak{Q}/\mathfrak{P})f(\mathfrak{P}/\mathfrak{p}) \end{aligned}$$

Proof. The first comes from substituting the factorization of $\mathfrak{P}\mathcal{O}_M$ in the factorization of $\mathfrak{p}\mathcal{O}_L$. The second comes from multiplicativity of degrees of field extensions. □

6 Computing factorizations

Theorem 2.6.1 (Criterion for ramification). *Assume $AKLB$, with L/K finite, A Dedekind, and B free over A . (The last condition is satisfied when A is a PID.) Then \mathfrak{p} ramifies in L iff $\mathfrak{p} \mid \text{disc}(B/A)$. In particular, only finitely many prime ideals ramify.*

Proof.

1. If A is a ring, B is a ring containing A and admitting a finite basis $\{e_1, \dots, e_m\}$ as an A -module, and \mathfrak{a} is an ideal of A , then $\{\bar{e}_1, \dots, \bar{e}_m\}$ is a basis for $B/\mathfrak{a}B$ as a A/\mathfrak{a} module, and $D(\bar{e}_1, \dots, \bar{e}_m) = D(e_1, \dots, e_m) \bmod \mathfrak{a}$. Hence

$$\text{disc}(B/A) \bmod \mathfrak{p} = \text{disc}((B/\mathfrak{p}B)/(A/\mathfrak{p})).$$

2. **Lemma 2.6.2.** *Let k be a perfect field and B be a k -algebra of finite dimension. Then B is reduced (has no nilpotent elements) iff $\text{disc}(B/k) \neq 0$.*

Proof. First suppose $\beta \neq 0$ is a nilpotent element of B . Choose a basis $e_1 = \beta, e_2, \dots, e_m$ of B . Then βe_i is nilpotent, so has trace 0. The first row of $(\text{tr}(e_i e_j))$ is zero, so $\text{disc}(B/k) = \det(\text{tr}(e_i e_j)) = 0$.

Now suppose B is reduced. By the Scheinnullstellensatz, $\bigcap_{\mathfrak{p} \text{ prime}} \mathfrak{p} = \text{nil}(R) = \{0\}$. Since B/\mathfrak{p} is integral and algebraic over k , Lemma 2.2.7 shows it is a field. Hence \mathfrak{p} is maximal, and different \mathfrak{p} are relatively prime. Let $\mathfrak{p}_1, \dots, \mathfrak{p}_r$ be prime ideals of B . By the Chinese Remainder Theorem, $B/\bigcap_{i=1}^r \mathfrak{p}_i = \prod_{i=1}^r B/\mathfrak{p}_i$ so

$$\dim_k B \geq \dim_k \left(B / \bigcap_{i=1}^r \mathfrak{p}_i \right) = \sum_{i=1}^r \dim_k (B/\mathfrak{p}_i) \geq r.$$

Since $\dim_k B$ is assumed finite, B has only finitely many prime ideals, say $\mathfrak{p}_1, \dots, \mathfrak{p}_g$. Each B/\mathfrak{p}_i is a *finite separable* (as k is perfect) extension of k , so by Proposition 1.3.4(2) (nondegeneracy of trace pairing), $\text{disc}((B/\mathfrak{p}_i)/k) \neq 0$. Since $B = B/\bigcap_{i=1}^g \mathfrak{p}_i = \prod_{i=1}^g B/\mathfrak{p}_i$, by taking the union of the bases for B/\mathfrak{p}_i , we get $\text{disc}(B/k) \neq 0$. \square

3. Let $\mathfrak{p}B = \prod_i \mathfrak{P}_i^{e_i}$. From the lemma, since A/\mathfrak{p} is perfect (as it is a finite field),

$$\text{disc}((B/\mathfrak{p}B)/(A/\mathfrak{p})) = 0$$

iff $B/\mathfrak{p}B$ is not reduced. By the Chinese Remainder Theorem $B/\mathfrak{p}B = \prod_i B/\mathfrak{P}_i^{e_i}$, and this is nonreduced iff some $e_i > 1$, i.e. \mathfrak{p} ramifies. \square

Theorem 2.6.3 (Computing the factorization of $\mathfrak{p}B$). *Assume $AKLB$, A is Dedekind and L/K is separable. Suppose $B = A[\alpha]$ and $f(X)$ is the minimal polynomial of α over K . Let \mathfrak{p} be a prime ideal in A , and suppose $f(X)$ factorizes into irreducible polynomials modulo \mathfrak{p} as*

$$f(X) \equiv \prod_{i=1}^r g_i(X)^{e_i} \pmod{\mathfrak{p}}.$$

Then

$$\mathfrak{p}B = \prod_{i=1}^r (\mathfrak{p}, g_i(\alpha))^{e_i}$$

is the prime factorization of $\mathfrak{p}B$. Moreover, letting $\bar{g}_i = g_i \pmod{\mathfrak{p}}$,

$$\begin{aligned} B/(\mathfrak{p}, g_i(\alpha)) &\cong (A/\mathfrak{p})[X]/(\bar{g}_i) \\ f_i &= \deg g_i. \end{aligned}$$

Proof. The map $X \mapsto \alpha$ gives an isomorphism

$$A[X]/(f(X)) \cong B.$$

Modding out by \mathfrak{p} gives

$$k[X]/(\bar{f}(X)) \cong B/\mathfrak{p}.$$

This gives a correspondence between ideals in $k[X]/(\bar{f}(X))$ and ideals in B containing \mathfrak{p} :

$$\begin{array}{ll} \text{Maximal ideals of } k[X]/(\bar{f}(X)) & (\bar{g}_i) \\ \longleftrightarrow \text{Maximal ideals of } B/\mathfrak{p} & (\bar{g}_i(\alpha)) \\ \longleftrightarrow \text{Maximal ideals of } B \text{ containing } \mathfrak{p} & (\mathfrak{p}, g_i(\alpha)) \end{array}$$

But the maximal ideals of B containing \mathfrak{p} are exactly the prime ideals (since B is Dedekind) dividing \mathfrak{p} (Lemma 2.5.1).

Now $\prod(\bar{g}_i)^{e_i} = 0$ but no power with smaller exponents is 0. Hence $\mathfrak{p}B \supseteq \prod(\mathfrak{p}, \bar{g}_i)^{e_i}$ but does not contain any power with smaller exponents, and equality holds. \square

Note that the condition that \mathfrak{p} be relatively prime to the conductor is somewhat pesky. The problem is that the we may have prime ideals dividing \mathfrak{p} that are in the form $(\mathfrak{p}, g(\alpha))$ where g does has coefficients with elements of \mathfrak{p} in the denominator. So looking at the polynomial modulo \mathfrak{p} fails to capture this behavior. We can't look at them modulo a power of \mathfrak{p} either—because then we would not be in a field. The solution is to pass to the completion with respect to \mathfrak{p} —we will do this in Chapter ??.

Example 2.6.4 (Quadratic extensions):

	Prime p	$x^2 + 1 \pmod p$	(p)
1.	2	$(x + 1)^2$	Ramifies: $(i + 1)^2$
	$p \equiv 1 \pmod{4}$	factors since $\left(\frac{-1}{p}\right) = 1$	Splits
	$p \equiv 3 \pmod{4}$	irreducible since $\left(\frac{-1}{p}\right) = -1$	Remains prime

	Prime p	$x^2 + 2 \pmod p$	(p)
2.	2	x^2	Ramifies: $(\sqrt{-2})^2$
	$p \equiv 1, 3 \pmod{8}$	factors since $\left(\frac{-2}{p}\right) = 1$	Splits
	$p \equiv 5, 7 \pmod{8}$	irreducible since $\left(\frac{-2}{p}\right) = -1$	Remains prime

	Prime p	$x^2 + x + 1 \pmod p$	(p)
3.	3	$(x - 1)^2$	Ramifies: $\left(\frac{-3 + \sqrt{-3}}{2}\right)^2$
	$p \equiv 1 \pmod{3}$	factors since $\left(\frac{-3}{p}\right) = \left(\frac{p}{3}\right) = 1$	Splits
	$p \equiv 2 \pmod{3}$	irreducible since $\left(\frac{-3}{p}\right) = \left(\frac{p}{3}\right) = -1$	Remains prime

Note we used quadratic reciprocity to translate the “square” condition into a modular condition on p . This is true in general for any quadratic ring: whether a prime p splits is entirely determined by a modular condition on p , because of quadratic reciprocity.

7 Decomposition and inertia groups

Let L/K be a finite Galois extension, with residue fields l and k .

For a prime \mathfrak{p} of K , we know that there are three kinds of behavior it could express when we pass to L :

1. It can split into distinct primes $\mathfrak{P}_1, \dots, \mathfrak{P}_g$.
2. The primes have some residue degree $f = [\mathcal{O}_L/\mathfrak{P}_j : \mathcal{O}_K/\mathfrak{p}]$ over \mathfrak{p} .

3. There can be ramification, the primes \mathfrak{P}_j appearing with exponent e .

Moreover, $[L : K] = efg$. We would like to separate these three kinds of behavior by defining two intermediate extensions $L^{D(\mathfrak{P})}$ and $L^{I(\mathfrak{P})}$.

Definition 2.7.1: Let $\mathfrak{P} \mid \mathfrak{p}$ be primes in L and K .

The **decomposition group** of \mathfrak{P} is

$$D_{L/K}(\mathfrak{P}) = \{\sigma \in G(L/K) : \sigma(\mathfrak{P}) = \mathfrak{P}\}.$$

The **inertia group** of \mathfrak{P} is

$$I_{L/K}(\mathfrak{P}) = \{\sigma \in G(L/K) : \sigma(\alpha) - \alpha \in \mathfrak{P} \text{ for all } \alpha \in \mathcal{O}_L\}.$$

Equivalently, letting l, k be the residue fields of L and K , $I_{L/K}(\mathfrak{P})$ is the kernel of the map $\varepsilon : D(\mathfrak{P}) \rightarrow G(l/k)$.

We drop the subscript when there is no confusion. The main theorem is the following.

Theorem 2.7.2. *Let L/K be a finite Galois extension with residue fields l, k , with l/k separable.¹ Let $\mathfrak{P} \mid \mathfrak{p}$ be primes of L and K . Let e, f, g be the ramification index, residue class degree, and number of prime divisors of \mathfrak{p} in L .*

Let $\mathfrak{P}_D = \mathfrak{P} \cap L^{D(\mathfrak{P})}$ and $\mathfrak{P}_I = \mathfrak{P} \cap L^{I(\mathfrak{P})}$ (the fixed fields of the decomposition and inertia groups). Then the following hold.

1. $[L : L^{I(\mathfrak{P})}] = e$ and \mathfrak{P}_I totally ramifies in $L/L^{I(\mathfrak{P})}$.

$$\mathfrak{P}_I \mathcal{O}_L = \mathfrak{P}^e.$$

2. $[L^{I(\mathfrak{P})} : L^{D(\mathfrak{P})}] = f$ and \mathfrak{P}_D remains inert in the extension $L^{I(\mathfrak{P})}/L^{D(\mathfrak{P})}$.

$$\begin{aligned} \mathfrak{P}_D \mathcal{O}_{L^{I(\mathfrak{P})}} &= \mathfrak{P}_I \\ f(\mathfrak{P}_I/\mathfrak{P}_D) &= f. \end{aligned}$$

Moreover, $L^{I(\mathfrak{P})}/K$ is Galois.


3. $[L^{D(\mathfrak{P})} : K] = g$, and \mathfrak{p} splits completely in $L^{D(\mathfrak{P})}$ if $L^{D(\mathfrak{P})}/K$ is Galois²:

$$\mathfrak{p} \mathcal{O}_{L^{D(\mathfrak{P})}} = \mathfrak{P}_{1,D} \cdots \mathfrak{P}_{g,D}.$$

¹If l/k is not assumed separable, then $[L : L^{I(\mathfrak{P})}] = e[l : k]_i$, $[L^{I(\mathfrak{P})} : L^{D(\mathfrak{P})}] = [l : k]_s$, and $[L^{D(\mathfrak{P})} : L] = g$.

²This is actually an iff. Exercise!

We have the following picture. By Galois theory, the groups on the right are the Galois groups acting on each extension; we set $G = G(L/K)$.



Remark 2.7.3: To study ramification, we can define subgroups of $I(\mathfrak{P})$ called ramification groups and get fixed fields in between L and $L^{I(\mathfrak{P})}$. See Chapter 9.

The rest of this section is devoted to the proof of Theorem 2.7.2. We keep the notations and assumptions in the theorem.

7.1 Decomposition group

Proposition 2.7.4: The decomposition group $D(\mathfrak{P})$ has order ef , and for $\sigma \in G(L/K)$,

$$D(\sigma(\mathfrak{P})) = \sigma D(\mathfrak{P}) \sigma^{-1}.$$

Moreover, the following are equivalent:

1. $D(\mathfrak{P})$ is normal in G .
2. The groups $D(\mathfrak{Q})$ are equal for all $\mathfrak{Q} \mid \mathfrak{p}$.
3. $L^{D(\mathfrak{P})}/L$ is Galois.

Proof. Since $D(\mathfrak{P})$ is the stabilizer of \mathfrak{P} under the action of $G := G(L/K)$, $|G/D(\mathfrak{P})|$ is simply the size of the orbit of G . This equals g since G acts transitively on the primes $\mathfrak{P}_1, \dots, \mathfrak{P}_g$ above \mathfrak{p} . Hence

$$|D(\mathfrak{P})| = \frac{|G|}{|G/D(\mathfrak{P})|} = \frac{n}{g} = ef.$$

The second part follows from the fact that if G acts on S and G is the stabilizer of $s \in S$, then tGt^{-1} is the stabilizer of ts .

For the equivalences, use the second part and the fundamental theorem of Galois theory 5.4.1. □

We first show that \mathfrak{P}_D is non-split in L and prove item 3 of Theorem 2.7.2.

By the Fixed Field Theorem, $D(\mathfrak{P}) = G(L/L^{D(\mathfrak{P})})$, and

$$[L : L^{D(\mathfrak{P})}] = |D(\mathfrak{P})| = ef. \quad (2.3)$$

Since $L/L^{D(\mathfrak{P})}$ is Galois, $D(\mathfrak{P})$ acts transitively on the primes of L above \mathfrak{P}_D . However, $D(\mathfrak{P})$ stabilizes \mathfrak{P} ; thus \mathfrak{P} is the only prime above \mathfrak{P}_D .

By the degree equation,

$$ef = [L : L^{D(\mathfrak{P})}] = e(\mathfrak{P}/\mathfrak{P}_D)f(\mathfrak{P}_D/\mathfrak{p}).$$

By Proposition 2.5.3,

$$\begin{aligned} e &= e(\mathfrak{P}/\mathfrak{P}_D)e(\mathfrak{P}_D/\mathfrak{p}) \\ f &= f(\mathfrak{P}/\mathfrak{P}_D)f(\mathfrak{P}_D/\mathfrak{p}). \end{aligned}$$

All equations are satisfied only when $e = e(\mathfrak{P}/\mathfrak{P}_D)$, $f = f(\mathfrak{P}/\mathfrak{P}_D)$, and $e(\mathfrak{P}_D/\mathfrak{p}) = f(\mathfrak{P}_D/\mathfrak{p}) = 1$.

If $L^{D(\mathfrak{P})}$ is Galois, then $e(\mathfrak{P}_D/\mathfrak{p}) = f(\mathfrak{P}_D/\mathfrak{p}) = 1$ are the same as the e and f values for all primes in $L^{D(\mathfrak{P})}$ over L . Thus \mathfrak{p} is totally split over L .

7.2 Inertia group

First we study the homomorphism

$$\varepsilon : D(\mathfrak{P}) \rightarrow G(l/k).$$

Proposition 2.7.5: Suppose $\mathfrak{P} \mid \mathfrak{p}$ are primes in L and K , and let k and l be the residue fields of L and K with respect to \mathfrak{P} and \mathfrak{p} .

1. l/k is normal (and hence Galois if separable).
2. Let ε be the map $D(\mathfrak{P}) \rightarrow G(l/k)$. Then ε is surjective.

Proof. Let $G = G(L/K)$.

1. We need to show that for $\bar{\alpha} \in l$, its minimal polynomial over k splits completely. Let α be a lift to \mathcal{O}_L and let

$$f(X) = \prod_{\sigma \in G} (X - \sigma(\alpha)) \in \mathcal{O}_K[X].$$

Taking this modulo \mathfrak{P} gives a polynomial in $k[X]$ containing $\bar{\alpha}$ as a root and splitting completely.

Thus l/k is normal, and hence Galois if it is separable.

2. First note we may assume l/k is separable. Indeed, we have $G(l/k) \cong G(l^{\text{sep}}/k)$ ³.

It suffices to show that $\varepsilon(D(\mathfrak{P}))$ acts transitively on the conjugates of $\bar{\alpha}$ over k (as then the image has at least $[l : k] = |G(l/k)|$ elements). By the Chinese Remainder Theorem, choose $\alpha \in \mathcal{O}_L$ such that

$$\alpha \equiv \begin{cases} \bar{\alpha} & (\text{mod } \mathfrak{P}) \\ 0 & (\text{mod } \mathfrak{P}'), \quad \mathfrak{P}' \neq \mathfrak{P}, \mathfrak{P}' \mid \mathfrak{p}. \end{cases}$$

Define f as in item 1. Then, noting that for $\sigma \in G \setminus D(\mathfrak{P})$, we have $\alpha \equiv 0 \pmod{\sigma^{-1}(\mathfrak{P})}$ and hence $\sigma(\alpha) \equiv 0 \pmod{\mathfrak{P}}$,

$$\begin{aligned} \bar{f}(X) &= \prod_{\sigma \in D(\mathfrak{P})} (X - \overline{\sigma(\alpha)}) \prod_{\sigma \notin D(\mathfrak{P})} x \\ &= \underbrace{\prod_{\sigma \in D(\mathfrak{P})} (X - \varepsilon(\sigma)(\bar{\alpha}))}_{(*)} \prod_{\sigma \notin D(\mathfrak{P})} x \in k[x] \end{aligned}$$

Now $(*)$ is in $k[x]$, so is divisible by the minimal polynomial of α over k . Given a conjugate $\bar{\alpha}'$ of $\bar{\alpha}$, it divides $(*)$, so equals $(\varepsilon(\sigma))(\bar{\alpha})$ for some σ . \square

Corollary 2.7.6. *There is a short exact sequence*

$$1 \rightarrow I(\mathfrak{P}) \rightarrow D(\mathfrak{P}) \rightarrow G(l/k) \rightarrow 1,$$

i.e. $D(\mathfrak{P})/I(\mathfrak{P}) \cong G(l/k)$.

Note $I(\mathfrak{P})$ is normal in $D(\mathfrak{P})$ as it is a kernel, so $L^{I(\mathfrak{P})}/K$ is Galois.

Now we finish the proof of Theorem 2.7.2. The above corollary gives

$$|D(\mathfrak{P})/I(\mathfrak{P})| = |G(l/k)| = [l : k] = f.$$

Since $G(L^{I(\mathfrak{P})}/L^{D(\mathfrak{P})}) = |D(\mathfrak{P})/I(\mathfrak{P})| = f$, we get $[L^{I(\mathfrak{P})} : L^{D(\mathfrak{P})}] = f$. From (2.3) we get $[L : L^{I(\mathfrak{P})}] = e$.

We will apply Corollary 2.7.6 to $L/L^{I(\mathfrak{P})}$. Note

$$D_{L/L^{I(\mathfrak{P})}}(\mathfrak{P}) = I_{L/L^{I(\mathfrak{P})}}(\mathfrak{P}) = G(L/L^{I(\mathfrak{P})}) = I(\mathfrak{P})$$

since the fact that $I(\mathfrak{P})$ operates trivially on l/k implies that it operates trivially on $l/\kappa(\mathfrak{P}_I)$. Hence the corollary gives

$$G(l/\kappa(\mathfrak{P}_I)) = 1,$$

i.e. $l = \kappa(\mathfrak{P}_I)$ and $f(\mathfrak{P}/\mathfrak{P}_I) = 1$. We know that \mathfrak{P}_D is non-split in L , so

$$\begin{aligned} e &= [L : L^{I(\mathfrak{P})}] = e(\mathfrak{P}/\mathfrak{P}_I) \underbrace{f(\mathfrak{P}/\mathfrak{P}_I)}_{=1} \\ f &= [L^{I(\mathfrak{P})} : L^{D(\mathfrak{P})}] = e(\mathfrak{P}_I/\mathfrak{P}_D) f(\mathfrak{P}_I/\mathfrak{P}_D). \end{aligned}$$

³From the Fixed Field Theorem $l/l^{G(l/l^{\text{sep}})}$ is Galois. But l/l^{sep} is purely inseparable and normal. Thus we must have $l = l^{G(l/l^{\text{sep}})}$, *i.e.* every automorphism of l/k is trivial on l/l^{sep} .

Now

$$\begin{aligned} e &= e(\mathfrak{P}/\mathfrak{P}_D) = e(\mathfrak{P}/\mathfrak{P}_I)e(\mathfrak{P}_I/\mathfrak{P}_D) \\ f &= f(\mathfrak{P}/\mathfrak{P}_D) = f(\mathfrak{P}/\mathfrak{P}_I)f(\mathfrak{P}_I/\mathfrak{P}_D), \end{aligned}$$

so we must have

$$\begin{aligned} e(\mathfrak{P}/\mathfrak{P}_I) &= e, & f(\mathfrak{P}/\mathfrak{P}_I) &= 1 \\ e(\mathfrak{P}_I/\mathfrak{p}) &= 1, & f(\mathfrak{P}_I/\mathfrak{p}) &= f. \end{aligned}$$

This finishes the proof.

7.3 Further properties and applications

Theorem 2.7.7. *Let M/K be a Galois extension and L/K a subextension. Then*

1.

$$\begin{aligned} D_{M/L}(\mathfrak{P}) &= D_{M/K}(\mathfrak{P}) \cap G(M/L) \\ I_{M/L}(\mathfrak{P}) &= I_{M/K}(\mathfrak{P}) \cap G(M/L). \end{aligned}$$

2. *If L/K is Galois, the following commutes and has exact rows and columns.*

$$\begin{array}{ccccccc} & & 1 & & 1 & & 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & I_{M/L} & \longrightarrow & I_{M/K} & \longrightarrow & I_{L/K} \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & D_{M/L} & \longrightarrow & D_{M/K} & \longrightarrow & D_{L/K} \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & G(M/L) & \longrightarrow & G(M/K) & \longrightarrow & G(L/K) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 1 & & 1 & & 1 \end{array}$$

Theorem 2.7.8. *Let L/K and L'/K be finite extensions. Then \mathfrak{p} unramified in L, L' if and only if \mathfrak{p} is unramified in LL' .*

Proof. First we prove the result for L, L' Galois. Note that for any Galois extension M/K , with $\mathfrak{P} \mid \mathfrak{p}$ primes in M and K ,

$$I_{\mathfrak{P}} = 1 \iff \mathfrak{p} \text{ unramified in } M. \tag{2.4}$$

Now there is a injective homomorphism

$$\begin{aligned}\Phi : G(LL'/K) &\hookrightarrow G(L/K) \times G(L'/K) \\ \Phi(\sigma) &= (\sigma|_L, \sigma|_{L'}).\end{aligned}$$

Take $\mathfrak{Q} | \mathfrak{p}$ with \mathfrak{Q} a prime in LL' , and let $\mathfrak{P} = \mathfrak{Q} \cap \mathcal{O}_L$ and $\mathfrak{P}' = \mathfrak{Q} \cap \mathcal{O}_{L'}$. Suppose $\sigma \in I_{\mathfrak{Q}}$. Then $\sigma(\mathfrak{Q}) = \mathfrak{Q}$ and hence, taking the intersections with $\mathcal{O}_L, \mathcal{O}_{L'}$ (which are fixed by σ since L, L' are Galois)

$$\begin{aligned}\sigma|_L(\mathfrak{P}) &= \mathfrak{P} \\ \sigma|_{L'}(\mathfrak{P}') &= \mathfrak{P}'.\end{aligned}$$

This shows $\sigma|_L \in I_{\mathfrak{P}}, \sigma|_{L'} \in I_{\mathfrak{P}'}$; by assumption and (2.4), we get $(\sigma|_L, \sigma|_{L'}) = (1, 1)$. By injectivity of Φ , $\sigma = 1$. This shows $I_{\mathfrak{Q}} = 1$, by (2.4) again, we get \mathfrak{Q} is unramified over \mathfrak{p} , as needed.

Now consider the general case. Given $\mathfrak{P} | \mathfrak{p}$ in L and K , let \mathfrak{Q} be a prime above \mathfrak{P} in the Galois closure L^{gal} . Now $(L^{\text{gal}})^{I_{\mathfrak{Q}}(L^{\text{gal}}/L)}$ is a Galois extension containing L ; since L^{gal} is the Galois closure of L , we get

$$L^{\text{gal}} = (L^{\text{gal}})^{I_{\mathfrak{Q}}(L^{\text{gal}}/L)},$$

But $[L^{\text{gal}} : (L^{\text{gal}})^{I_{\mathfrak{Q}}(L^{\text{gal}}/L)}]$ is the ramification degree of $\mathfrak{Q}/\mathfrak{P}$; we see that it is 1, i.e. \mathfrak{Q} is not ramified over \mathfrak{P} and hence not ramified over \mathfrak{p} . Thus L^{gal}/K is unramified. Similarly, L'^{gal}/K is unramified. By the above, $L^{\text{gal}}L'^{\text{gal}}/K$ is unramified, so LL'/K is unramified. \square

8 Problems

1. A **half-factorial domain** (HFD) A is an integral domain where any given factorization of a has the same length. Prove Carlitz's Theorem:

Theorem 2.8.1 (Carlitz). *The ring of integers \mathcal{O}_K is a HFD iff the class group has order at most 2.*

See AMM, 12/2011, for related results.

2. Show that if \mathfrak{p} splits completely in $L^{D(\mathfrak{P})}$, then $L^{D(\mathfrak{P})}/L$ is Galois.

Conclude that if \mathfrak{p} splits completely in L , then \mathfrak{p} splits completely in the Galois closure L^{gal} .

Chapter 3

The class group

1 Norms of ideals

Assume AKLB, A is Dedekind, and L/K is separable. We generalize the definition of norm to ideals, not just elements, so that it is a map $\text{Id}(B) \rightarrow \text{Id}(A)$ that is consistent with our old condition, i.e.

$$\text{Nm}_{L/K}((a)) = (\text{Nm}_{L/K}(b)).$$

Consider a principal ideal $\mathfrak{p} = (p) \subseteq A$, and suppose it factors in B as $\mathfrak{p}B = \prod_{i=1}^g \mathfrak{P}^{e_i}$. We want the norm to satisfy

$$\text{Nm}_{L/K}(p) = \text{Nm}_{L/K}(\mathfrak{p}B) = \prod_{i=1}^g \text{Nm}_{L/K}(\mathfrak{P})^{e_i}, \quad (3.1)$$

since we want it to be multiplicative. But $\text{Nm}(p) = p^n$ where $n = [L : K]$. By the degree equation, if $\text{Nm}(\mathfrak{P}) = \mathfrak{P}^{f_i}$ where $f_i = [B/\mathfrak{P}_i : A/\mathfrak{p}]$, then (3.1) will be satisfied. Hence we make the following definition.

Definition 3.1.1: For \mathfrak{P} is a prime of B , let $\mathfrak{p} = \mathfrak{P} \cap A$ and $f(\mathfrak{P}/\mathfrak{p}) = [B/\mathfrak{P} : A/\mathfrak{p}]$. Define the norm of \mathfrak{P} to be

$$\text{Nm}_{L/K}(\mathfrak{P}) = \mathfrak{p}^{f(\mathfrak{P}/\mathfrak{p})}.$$

This extends uniquely to a homomorphism $\text{Id}(A) \rightarrow \text{Id}(B)$, since the ideal group is free.

Proposition 3.1.2 (Behavior with respect to field extensions):

1. For an ideal $\mathfrak{a} \subseteq A$,

$$\text{Nm}_{L/K}(\mathfrak{a}B) = \mathfrak{a}^m,$$

where $m = [L : K]$.

2. If L/K is Galois and $\mathfrak{p} \neq 0$ is a prime ideal of A , and $\mathfrak{P} \mid \mathfrak{p}$, then

$$\text{Nm}_{L/K}(\mathfrak{p}) = \prod_{\sigma \in G(L/K)} \sigma \mathfrak{P}.$$

3. For any nonzero $\beta \in B$, $\text{Nm}_{L/K}(\beta B) = \text{Nm}_{L/K}(\beta)A$. (I.e. this is consistent with our previous definition.)

Compare the first two items to Chapter 1, Proposition 1.2.2(5) and Proposition 1.2.3(2b), respectively.

Proof.

1. By the degree equation (Theorem 2.5.2), for \mathfrak{p} prime

$$\text{Nm}_{L/K}(\mathfrak{p}B) = \text{Nm}_{L/K} \left(\prod_i \mathfrak{P}_i^{e_i} \right) = \mathfrak{p}^{\sum_i e_i f_i} = \mathfrak{p}^m.$$

The general statement follows by multiplicativity of $\text{Nm}_{L/K}$.

2. $G(L/K)$ acts transitively on $\{\mathfrak{P}_1, \dots, \mathfrak{P}_g\}$, so each \mathfrak{P}_i occurs $\frac{m}{g} = ef$ times in $\{\sigma\mathfrak{P} \mid \sigma \in G(L/K)\}$.
3. First suppose L/K is Galois. We use the description in terms of Galois conjugates to relate the norms of elements with the norms of ideals. By part 2 and Proposition 1.2.3(2b), we have

$$\text{Nm}_{L/K}(\beta B) \cdot B \stackrel{(2)}{=} \prod_{\sigma \in G(L/K)} \sigma(\beta B) = \left(\prod_{\sigma \in G(L/K)} \sigma(\beta) \right) B \stackrel{1.2.3}{=} \text{Nm}_{L/K}(\beta) \cdot B.$$

Hence, $\text{Nm}_{L/K}(\beta) \cdot A$ and $\text{Nm}_{L/K}(\beta \cdot B)$ determine the same ideal in B . Since $\text{Id}(A) \rightarrow \text{Id}(B)$ is injective, they are equal in A .

Now consider the general case. Let M be the Galois closure of L over K , let $C = \mathcal{O}_M$, and let $d = [M : L]$. Then the above, together with part 1 and Proposition 1.2.2(5), give

$$\text{Nm}_{L/K}(\beta \cdot B)^d \stackrel{(1)}{=} \text{Nm}_{M/K}(\beta \cdot B) = \text{Nm}_{M/K}(\beta) \cdot A \stackrel{1.2.2(5)}{=} \text{Nm}_{L/K}(\beta)^d \cdot A.$$

Since $\text{Id}(B)$ is torsion-free, $\text{Nm}_{L/K}(\beta \cdot B) = \text{Nm}_{L/K}(\beta) \cdot A$. □

Definition 3.1.3: The **numerical norm** of \mathfrak{a} in \mathcal{O}_K is its index in the lattice of integers:

$$\mathfrak{N}\mathfrak{a} = [\mathcal{O}_K : \mathfrak{a}].$$

Note the following comparisons between the ideal and numerical norms.

1. The ideal norm is defined for a field extension K/F while the numerical norm is defined for any number field K/\mathbb{Q} .
2. The ideal norm returns an ideal while the numerical norm returns an integer.

3. However, if we take the base field F to be \mathbb{Q} , and identify integers with the ideals they generate, the two norms are equivalent. This is the content of the following proposition.

Proposition 3.1.4 (Relationship between ideal and numerical norm):

1. For any ideal $\mathfrak{a} \subseteq \mathcal{O}_K$,

$$\mathrm{Nm}_{K/\mathbb{Q}}(\mathfrak{a}) = (\mathfrak{N}(a)).$$

Therefore, $\mathfrak{N}(ab) = \mathfrak{N}(a)\mathfrak{N}(b)$.

2. Let $\mathfrak{b} \subseteq \mathfrak{a} \subseteq K$ be fractional ideals. Then

$$[\mathfrak{a} : \mathfrak{b}] = \mathfrak{N}(\mathfrak{a}^{-1}\mathfrak{b}).$$

In other words, *the norm of an ideal is its index in the ring of integers.*

Proof.

1. Write $\mathfrak{a} = \prod \mathfrak{p}_i^{e_i}$ and let $(p_i) = \mathbb{Z} \cap \mathfrak{p}_i$, $f_i = f(\mathfrak{p}_i/(p_i))$. By the Chinese remainder theorem,

$$\mathcal{O}_K/\mathfrak{a} \cong \prod_i \mathcal{O}_K/\mathfrak{p}_i^{e_i}.$$

Since $\mathcal{O}_K/\mathfrak{p}_i^{e_i}$ is a vector space over \mathbb{F}_{p_i} of dimension $e_i f_i$, we find

$$\mathfrak{N}\mathfrak{a} = |\mathcal{O}_K/\mathfrak{a}| = \prod_i p_i^{e_i f_i} = \mathrm{Nm}_{K/\mathbb{Q}}(\mathfrak{a}).$$

Multiplicativity follows from the same property for the ideal norm.

2. We can multiply by an integer d so that \mathfrak{a} and \mathfrak{b} are integral ideals. Then

$$[\mathfrak{a} : \mathfrak{b}] = [d\mathfrak{a} : d\mathfrak{b}] = \frac{[\mathcal{O}_K : d\mathfrak{b}]}{[\mathcal{O}_K : d\mathfrak{a}]} = \frac{\mathfrak{N}(d\mathfrak{b})}{\mathfrak{N}(d\mathfrak{a})} \stackrel{(1)}{=} \mathfrak{N}(\mathfrak{a}^{-1}\mathfrak{b}). \quad \square$$

2 Minkowski's Theorem

Theorem 3.2.1 (Minkowski). *Let V be a subset of \mathbb{R}^n that is convex and symmetric around the origin (“centrally symmetric”). Let L be a full lattice with fundamental parallelepiped D . If*

$$\mu(T) > 2^n \mu(D)$$

then T contains a point of L other than the origin. If furthermore D is compact, we can weaken the hypothesis to

$$\mu(T) \geq 2^n \mu(D).$$

Proof. First note that if S is a measurable set such that $\mu(S) > \mu(D)$, then S contains two points a, b such that $a - b \in L$. Indeed, we can tile the space with fundamental parallelepipeds, and translate each of them to the origin. We consider the intersections of these parallelepipeds with S . Since the sum of these volumes is $\mu(S) > \mu(D)$, and they are all packed in D , there must be overlap, i.e. unequal $a, b \in S$ that were translated to the same point. This implies $a - b \in L$.

The set $S = \frac{1}{2}T$ has volume $\frac{1}{2^n}T > \mu(D)$. Hence by the above, there exist $\frac{1}{2}a \neq \frac{1}{2}b \in S$ ($a, b \in T$) such that $\frac{1}{2}a - \frac{1}{2}b \in L$. Since T is symmetric, $-b \in T$; since T is convex, $\frac{1}{2}(a - b) \in T$. This is the desired lattice point.

Now suppose instead T is convex and $\mu(T) \geq 2^n \mu(D)$. Let L_n be the set of lattice points in $(1 + \frac{1}{n})T$ other than the origin. By the first part, L_n is nonempty; since T is bounded it must be finite. We have that $L_n \subseteq L_m$ when $n \geq m$. Hence

$$T \cap L = \bigcap_{n=1}^{\infty} \left(1 + \frac{1}{n}\right) T \cap L = \bigcap_{n=1}^{\infty} L_n \neq \phi. \quad \square$$

Theorem 3.2.2 (Sums of four squares). (*A digression, but nice to talk about*)

3 Finiteness of the class number

We now show that the class number is finite (Theorem 3.3.6). The idea of the proof is as follows.

1. Embed K as a \mathbb{Q} -vector space in $\mathbb{R}^r \times \mathbb{C}^s$. Under the \mathbb{R} -vector space isomorphism $K \otimes_{\mathbb{Q}} \mathbb{R} \rightarrow \mathbb{R}^r \times \mathbb{C}^s$, the ideal \mathfrak{a} is realized as a lattice L in $V = \mathbb{R}^r \times \mathbb{C}^s$ (Proposition 3.3.1). The norm on K translates into a “norm” on V .
2. Find an element in \mathfrak{a} of small norm (Theorem 3.3.2): Find a compact, symmetric convex set in V consisting of elements of norm at most R . Choosing R large enough, we can make sure V has large volume. By Minkowski’s Theorem, V contains an element of L .
3. Using step 2, show that every ideal class contains an representative of norm at most a constant (Theorem 3.3.5).
4. Show that there are a finite number of ideals with bounded norm (Lemma 3.3.7).

We first embed \mathfrak{a} as a full lattice using the embeddings of K , and find the volume of the fundamental parallelepiped in terms of the discriminant (the discriminant is related to the embeddings by Proposition 1.3.4).

Let $\{\sigma_1, \dots, \sigma_r\}$ be the real embeddings and $\{\sigma_{r+1}, \bar{\sigma}_{r+1}, \dots, \sigma_{r+s}, \bar{\sigma}_{r+s}\}$ be the complex embeddings of K . This gives an embedding¹

$$\begin{aligned}\sigma : K &\hookrightarrow \mathbb{R}^r \times \mathbb{C}^s \\ \sigma(\alpha) &= (\sigma_1\alpha, \dots, \sigma_{r+s}\alpha).\end{aligned}$$

Identify $V = \mathbb{R}^r \times \mathbb{C}^s$ with \mathbb{R}^n using the basis $\{1, i\}$ for \mathbb{C} .

Proposition 3.3.1: Let \mathfrak{a} be an ideal in \mathcal{O}_K . Then $\sigma(\mathfrak{a})$ is a full lattice in V and the volume of its parallelepiped is $2^{-s} \cdot \mathbb{N}\mathfrak{a} \cdot |\Delta_K|^{\frac{1}{2}}$.

Proof. Let $\alpha_1, \dots, \alpha_n$ be a basis for \mathfrak{a} as a \mathbb{Z} -module. To prove that $\sigma(\mathfrak{a})$ is a lattice, we need to show $\sigma(\alpha_1), \dots, \sigma(\alpha_n)$ are linearly independent, i.e. the following has nonzero determinant:

$$A = \begin{pmatrix} \sigma_1(\alpha_1) & \cdots & \sigma_r(\alpha_1) & \Re(\sigma_{r+1}(\alpha_1)) & \Im(\sigma_{r+1}(\alpha_1)) & \cdots \\ \sigma_1(\alpha_2) & \cdots & \sigma_r(\alpha_2) & \Re(\sigma_{r+1}(\alpha_2)) & \Im(\sigma_{r+1}(\alpha_2)) & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

To do this we relate this to the matrix

$$B = \begin{pmatrix} \sigma_1(\alpha_1) & \cdots & \sigma_r(\alpha_1) & \sigma_{r+1}(\alpha_1) & \overline{\sigma_{r+1}(\alpha_1)} & \cdots \\ \sigma_1(\alpha_2) & \cdots & \sigma_r(\alpha_2) & \sigma_{r+1}(\alpha_2) & \overline{\sigma_{r+1}(\alpha_2)} & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Note $\det(B) = \pm \text{disc}(\alpha_1, \dots, \alpha_n)^{\frac{1}{2}} \neq 0$. Let $J = \begin{pmatrix} \frac{1}{2} & \frac{1}{2i} \\ \frac{1}{2} & -\frac{1}{2i} \end{pmatrix}$. Then

$$A = B \begin{pmatrix} I_r & 0 & 0 & \cdots \\ 0 & J & 0 & \cdots \\ 0 & 0 & J & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Using

$$\text{disc}(\alpha_1, \dots, \alpha_n) = [\mathcal{O}_K : \mathfrak{a}]^2 \cdot |\text{disc}(\mathcal{O}_K/\mathbb{Z})|$$

we get that the volume of a fundamental parallelepiped for D is

$$\mu(D) = |\det(A)| = 2^{-s} |\det(B)| = 2^{-s} |\text{disc}(\alpha_1, \dots, \alpha_n)|^{\frac{1}{2}} = 2^{-s} \cdot \mathbb{N}\mathfrak{a} \cdot |\Delta_K|^{\frac{1}{2}}.$$

(In particular, this is nonzero.) □

¹This is the canonical embedding $K \hookrightarrow K \otimes_{\mathbb{Q}} \mathbb{R}$: Indeed, by Chinese Remainder

$$K \otimes_{\mathbb{Q}} \mathbb{R} = \mathbb{Q}[x]/(f(x)) \otimes_{\mathbb{Q}} \mathbb{R} = \prod_{i=1}^r \mathbb{R}[x]/(x - \sigma_i\alpha) \times \prod_{j=1}^s (\mathbb{R}[x]/(x - \sigma_{r+j}\alpha)(x - \bar{\sigma}_{r+j}\alpha)) \cong \mathbb{R}^r \times \mathbb{C}^s.$$

Theorem 3.3.2. *Let \mathfrak{a} be a nonzero ideal in \mathcal{O}_K . Then \mathfrak{a} contains a nonzero element α of K with*

$$|\mathrm{Nm}(\alpha)| \leq \left(\frac{4}{\pi}\right)^s \frac{n!}{n^n} \mathbb{N}\mathfrak{a} |\Delta_K|^{\frac{1}{2}}.$$

Proof. The norm on K translates into the “norm”

$$\mathrm{Nm}(x_1, \dots, x_r, z_{r+1}, \dots, z_{r+s}) = |x_1| \cdots |x_r| |z_{r+1}|^2 \cdots |z_{r+s}|^2.$$

However, $\mathbb{N}\mathfrak{a} < r$ is by no means a compact convex set. Fortunately, however, we note by the AM-GM inequality that

$$|\mathrm{Nm}(\mathbf{x})| = |x_1| \cdots |x_r| |z_{r+1}|^2 \cdots |z_{r+s}|^2 \leq \left(\frac{\sum_{k=1}^r |x_k| + 2 \sum_{k=1}^s |z_{r+k}|}{n} \right)^n. \quad (3.2)$$

Defining the norm $\|\cdot\|$ on $V = \mathbb{R}^r \times \mathbb{C}^s$ by

$$\|(x_1, \dots, x_r, z_{r+1}, \dots, z_{r+s})\| = \sum_{k=1}^r |x_k| + 2 \sum_{k=r+1}^s |z_k|,$$

and letting $B(t) = \{x \in V : \|x\| < t\}$, $B(\mathrm{Nm}, t) = \{x \in V : |\mathrm{Nm}(x)| < t\}$, we see from (3.2) that

$$B(t) \subseteq B\left(\mathrm{Nm}, \frac{t^n}{n^n}\right). \quad (3.3)$$

To apply Minkowski we need some computations.

Lemma 3.3.3. *The volume of $B(t) = \{x \in V : \|x\| < t\}$ is*

$$\mu(B(t)) = 2^{r-s} \pi^s \frac{t^n}{n!}.$$

Proof. We write the complex variables as $z_k = x_k + y_k i$. Let

$$B'(t) = \{(x_1, \dots, x_r, x_{r+1}, y_{r+1}, \dots, x_{r+s}, y_{r+s}) \in B(t) : x_1, \dots, x_r \geq 0\}.$$

Write $dV = dx_1 \cdots dx_n$. Using symmetry and a polar change of coordinates, we compute

$$\mu(B(t)) = 2^r \int_{B'(t)} dV dx_{r+1} dy_{r+1} \cdots dx_{r+s} dy_{r+s} \quad (3.4)$$

$$= 2^r \int_{x_1, \dots, x_r \geq 0, \sum x_k + 2 \sum \rho_k \leq t} (\rho_{r+1} \cdots \rho_{r+s}) dV d\rho_{r+1} d\theta_{r+1} \cdots d\rho_{r+s} d\theta_{r+s} \quad (3.5)$$

$$= 2^{r-2s} \int_{x_1, \dots, x_r \geq 0, \sum x_k + \sum \rho_k \leq t} (\rho_{r+1} \cdots \rho_{r+s}) dV d\rho_{r+1} d\theta_{r+1} \cdots d\rho_{r+s} d\theta_{r+s}$$

$$= 2^{r-2s} (2\pi)^s \int_{x_1, \dots, x_r \geq 0, \sum x_k + \sum \rho_k \leq t} (\rho_{r+1} \cdots \rho_{r+s}) dV d\rho_{r+1} \cdots d\rho_{r+s}$$

$$= 2^{r-s} \pi^s t^{(r+s)+s} \frac{1}{((r+s)+s)!} \quad (3.6)$$

$$= 2^{r-s} \pi^s \frac{t^n}{n!}.$$

Note (3.4) follows by symmetry, (3.5) follows from polar change of coordinates, and (3.6) follows from the lemma below. \square

Lemma 3.3.4.

$$\int_{x_i \geq 0, \sum x_i \leq t} x_1^{a_1} \cdots x_m^{a_m} dx_1 \cdots dx_m = t^{m + \sum_{i=1}^m a_i} \frac{\Gamma(a_1 + 1) \cdots \Gamma(a_m + 1)}{\Gamma(a_1 + \cdots + a_m + m + 1)}.$$

Proof. Making the substitution $x_i = tx'_i$, $dx_i = t dx'_i$, we find that the integral equals

$$t^{m + \sum_{i=1}^m a_i} \int_{x_i \geq 0, \sum x_i \leq 1} x_1^{a_1} \cdots x_m^{a_m} dx_1 \cdots dx_m.$$

Hence it suffices to prove the lemma for $t = 1$.

For $m = 1$, note

$$\int_0^1 x^a dx = \frac{1}{a+1} = \frac{\Gamma(a+1)}{\Gamma(a+2)}.$$

For $m = 2$, let $B(\alpha, \beta) = \int_0^1 v^{\alpha-1} (1-v)^{\beta-1} dv$. We need to show $B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$. By Fubini,

$$\Gamma(\alpha)\Gamma(\beta) = \int_0^\infty \int_0^\infty s^{\alpha-1} e^{-s} t^{\beta-1} e^{-t} ds dt = \int_0^\infty \int_0^\infty s^{\alpha-1} t^{\beta-1} e^{-(s+t)} ds dt.$$

Note $F : (0, \infty) \times (0, 1) \rightarrow (0, \infty)^2$ with $F(u, v) = (uv, u(1-v))$ is a diffeomorphism. Indeed, it has an inverse $F^{-1}(s, t) = (t + s, \frac{s}{t+s})$ hence is bijective and its Jacobian is $\det \begin{pmatrix} v & -u \\ -v & -u \end{pmatrix} = u \neq 0$. Using the change of variables $(s, t) = F(u, v)$ gives

$$\begin{aligned} \int_0^1 \int_0^\infty (uv)^{\alpha-1} (u(1-v))^{\beta-1} e^{-(uv+u(1-v))} u du dv &= \int_0^1 \int_0^\infty u^{\alpha+\beta-1} e^{-u} v^{\alpha-1} (1-v)^{\beta-1} du dv \\ &= \left(\int_0^\infty u^{\alpha+\beta-1} e^{-u} du \right) \left(\int_0^1 v^{\alpha-1} (1-v)^{\beta-1} dv \right) \\ &= \Gamma(\alpha + \beta) B(\alpha, \beta), \end{aligned}$$

as needed.

Now we use induction; suppose the theorem proved for $m - 1$. We have

$$\begin{aligned} \int_{x_i \geq 0, \sum_{i=1}^m x_i \leq 1} x_1^{a_1} \cdots x_m^{a_m} dx_1 \cdots dx_m &= \int_0^1 x_m^{a_m} \int_{x_i \geq 0, \sum_{i=1}^{m-1} x_i \leq 1-x_m} x_1^{a_1} \cdots x_{m-1}^{a_{m-1}} dx_1 \cdots dx_{m-1} dx_m \\ &= \int_0^1 x_m^{a_m} (1-x_m)^{m-1 + \sum_{i=1}^{m-1} a_i} \frac{\Gamma(a_1 + 1) \cdots \Gamma(a_{m-1} + 1)}{\Gamma(a_1 + \cdots + a_{m-1} + m)} dx_m \\ &= \frac{\Gamma(a_m + 1) \Gamma(\sum_{i=1}^{m-1} a_i + m)}{\Gamma(a_1 + \cdots + a_m + m + 1)} \cdot \frac{\Gamma(a_1 + 1) \cdots \Gamma(a_{m-1} + 1)}{\Gamma(a_1 + \cdots + a_{m-1} + m)} \\ &= \frac{\Gamma(a_1 + 1) \cdots \Gamma(a_m + 1)}{\Gamma(a_1 + \cdots + a_m + m + 1)}, \end{aligned}$$

using the induction hypothesis and the $m = 2$ case. □

Taking

$$t = \sqrt[n]{n! \cdot \frac{2^{n-r}}{\pi^s} \cdot \mathbb{N}\mathfrak{a}|\Delta_K|^{\frac{1}{2}}}$$

we find by Lemma 3.3.3 that

$$\mu(B(t)) = 2^{r-s}\pi^s \frac{t^n}{n!} = 2^n \left(2^{-s}\mathbb{N}\mathfrak{a}|\Delta_K|^{\frac{1}{2}}\right) = 2^n \mu(D)$$

where D is the fundamental parallelepiped. Note that $B(t)$ is a closed ball, and it is convex by the triangle inequality. Hence by Minkowski's Theorem, $B(t)$ contains an element of $\sigma(\mathfrak{a})$. For this element, we have by (3.3) that

$$\mathrm{Nm}_{K/\mathbb{Q}}(a) \leq \frac{t^n}{n^n} = \left(\frac{4}{\pi}\right)^s \frac{n!}{n^n} \mathbb{N}\mathfrak{a}|\Delta_K|^{\frac{1}{2}}. \quad \square$$

Theorem 3.3.5. *Suppose K/\mathbb{Q} is an extension of degree n , and let $\Delta_K = \mathrm{disc}(K/\mathbb{Q})$. Let $2s$ be the number of nonreal complex embeddings of K . Then there exists a set of representatives for the ideal class group $\mathrm{Cl}(K)$ consisting of integral ideals \mathfrak{a} with*

$$\mathbb{N}(\mathfrak{a}) \leq \underbrace{\frac{n!}{n^n} \left(\frac{4}{\pi}\right)^s}_{C_K} |\Delta_K|^{\frac{1}{2}}.$$

Proof. Given a fractional ideal \mathfrak{c} , there exists \mathfrak{b} such that

$$\mathfrak{b}\mathfrak{c} = (d)$$

is principal. By Theorem 3.3.2, there is an element $\beta \in \mathfrak{b}$ of norm at most $\left(\frac{4}{\pi}\right)^s \frac{n!}{n^n} \mathbb{N}\mathfrak{b}|\Delta_K|^{\frac{1}{2}}$. Since $(\beta) \subseteq \mathfrak{b}$ we have

$$\mathfrak{a}\mathfrak{b} = (\beta)$$

for some \mathfrak{a} . Note $\mathfrak{a} \sim \mathfrak{b}^{-1} \sim \mathfrak{c}$, and taking norms of the above equation gives

$$\mathfrak{N}\mathfrak{a}\mathfrak{N}\mathfrak{b} = \mathfrak{N}(\beta) \leq \left(\frac{4}{\pi}\right)^s \frac{n!}{n^n} \mathbb{N}\mathfrak{b}|\Delta_K|^{\frac{1}{2}}.$$

Canceling $\mathfrak{N}\mathfrak{b}$ gives that \mathfrak{a} is the desired representative. □

Theorem 3.3.6. *The class number of K is finite.*

Proof. By Theorem 3.3.5, every ideal class has a representative with norm at most $C_K|\Delta_K|^{\frac{1}{2}}$. Thus it suffices show the following (take $C = C_K|\Delta_K|^{\frac{1}{2}}$).

Lemma 3.3.7. *There are only a finite number of integral ideals \mathfrak{a} with $\mathbb{N}\mathfrak{a} \leq C$ (take $C = C_K|\Delta_K|^{\frac{1}{2}}$).*

Proof. Suppose \mathfrak{a} is an integral ideal. Write $\mathfrak{a} = \prod \mathfrak{p}_i^{r_i}$. Let $(p_i) = \mathfrak{p}_i \cap \mathbb{Z}$ and $f_i = [\mathcal{O}_K/\mathfrak{p}_i : \mathbb{Z}/(p_i)]$. Then

$$\mathbb{N}\mathfrak{a} = \prod_i p_i^{f_i r_i}.$$

Given $\mathbb{N}\mathfrak{a} \leq C$, there are a finite possibilities for the p_i and hence \mathfrak{p}_i , as well as for the r_i . \square

\square

The bound in Theorem 3.3.5 also gives the following corollaries.

Theorem 3.3.8. *Every algebraic extension of \mathbb{Q} ramifies over \mathbb{Q} .*

Proof. It suffices to prove this statement for finite extensions. Let K/\mathbb{Q} be a finite extension. By Theorem 3.3.2, every ideal contains a representative α with

$$1 \leq |\mathrm{Nm}(\alpha)| \leq \left(\frac{4}{\pi}\right)^s \frac{n!}{n^n}.$$

Hence we have

$$|\Delta_K| \geq \frac{n^{2n}}{n!^2} \left(\frac{\pi}{4}\right)^{2s} > 1. \quad (3.7)$$

The last inequality comes from the fact that defining $a_n = \frac{n^{2n}}{n!^2} \left(\frac{\pi}{4}\right)^{2s}$, we have that $a_2 > 1$ and $\frac{a_{n+1}}{a_n} = \left(\frac{\pi}{4}\right)^{\frac{1}{2}} \left(1 + \frac{1}{n}\right)^n > 1$ for $n \geq 2$.

Since $\Delta_K > 1$ and every prime dividing the discriminant ramifies (Theorem 2.6.1), K/\mathbb{Q} is ramified. \square

Corollary 3.3.9. *There does not exist an irreducible monic polynomial $f(X) \in \mathbb{Z}[X]$ of degree greater than 1 with discriminant ± 1 .*

Proof. Let f be an irreducible monic polynomial of degree greater than 1. Let α be a root of f . By Theorem 3.3.8, $\mathbb{Q}[\alpha]$ is ramified over \mathbb{Q} . By (3.7), $|\Delta_K| > 1$. Then

$$\mathrm{disc}(f) = \mathrm{disc}(\mathbb{Z}[\alpha]/\mathbb{Z}) = |\Delta_K| \cdot [\mathcal{O}_K : \mathbb{Z}[\alpha]]^2 > 1. \quad \square$$

4 Example: Quadratic extensions

To compute the class group in quadratic extensions, note the following two facts.

1. The complete description of prime ideals is given by Example ?? (actually put this in!).
2. By Theorem 3.3.2, each ideal class has a representative of norm at most $\frac{4}{\pi} |\Delta_K|^{\frac{1}{2}}$.

In fact, Minkowski's bound can be improved in the quadratic case.

Theorem 3.4.1. (*) Let $K = \mathbb{Q}(\sqrt{d})$ where d is a negative squarefree integer. Let

$$\mu = \begin{cases} \sqrt{\frac{|d|}{3}}, & d \equiv 1 \pmod{4} \\ 2\sqrt{\frac{|d|}{3}}, & d \equiv 2, 3 \pmod{4}. \end{cases}$$

Every ideal class in \mathcal{O}_K has a representative \mathfrak{a} with

$$\mathfrak{N}\mathfrak{a} \leq \mu.$$

Proof. First we show that every ideal \mathfrak{a} has an element $a \neq 0$ with $\text{Nm}_{K/\mathbb{Q}}(a) \leq \mu \mathfrak{N}(\mathfrak{a})$. For a lattice L let $\Delta(L)$ be the area of a fundamental parallelogram.

Note that $\text{Nm}_{K/\mathbb{Q}}(z) = |z|^2$. An ideal \mathfrak{a} of K forms a lattice in \mathbb{C} . Let a be the element of minimal nonzero norm in \mathfrak{a} and b be the element of minimal nonzero norm that is not an integer multiple of a . By the minimality assumption, since $b - a$ cannot be an integer multiple of a , we have

$$|b - a| \geq |b| \geq |a|.$$

Let A, B denote the points a, b and O the origin. Using the fact that in a triangle the side lengths are in the same order least-to-greatest as the opposite angles, we get that in the triangle AOB , the angle at O is largest, in particular at least 60° . Let O' be so that $OAO'B$ is a parallelogram. The minimality assumption similarly forces $OO' \geq AO, AO'$, so we get $\angle OAO' \geq 60^\circ$. Thus

$$60^\circ \leq \angle AOB \leq 120^\circ. \quad (3.8)$$

Furthermore, the parallelogram with sides OA and OB is a fundamental parallelogram: Suppose C is the point $c \in \mathfrak{a}$, and is in the triangle OAB but not any of the vertices. Let OC intersect AB at C' . We have $\angle OC'B > \angle OAB \geq \angle ABO = \angle C'BO$, where the middle inequality is from $OB \geq OA$. Hence looking at $\triangle OC'B$, $OB > OC' \geq OC$, contradicting minimality of b . Similarly, if C is in ABO' , then we have $|a + b - c| < |b|$, also a contradiction.

By (3.8), the area of the fundamental parallelogram is

$$\Delta(\mathcal{O}_K)\mathfrak{N}\mathfrak{a} = \Delta(\mathfrak{a}) = |ab| \sin \angle AOB \geq |a|^2 \frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{2} \text{Nm}_{K/\mathbb{Q}}(a).$$

Solving gives

$$\text{Nm}_{K/\mathbb{Q}}(a) \leq \frac{2}{\sqrt{3}} \Delta(\mathcal{O}_K)\mathfrak{N}\mathfrak{a}.$$

Finally note that for $d \equiv 1 \pmod{4}$, a basis for \mathcal{O}_K is $(1, \frac{1+\sqrt{d}}{2})$ while for $d \equiv 2, 3 \pmod{4}$ the basis is $(1, \sqrt{d})$. The fundamental parallelograms have areas $\frac{\sqrt{d}}{2}$ and \sqrt{d} , respectively, giving

$$\text{Nm}_{K/\mathbb{Q}}(a) \leq \mu \mathfrak{N}\mathfrak{a}.$$

Given a fractional ideal \mathfrak{c} , there exists \mathfrak{b} such that

$$\mathfrak{b}\mathfrak{c} = (d)$$

is principal. By the above, there is an element $b \in \mathfrak{b}$ of norm at most $\mu\mathfrak{N}\mathfrak{b}$. Since $(b) \subseteq \mathfrak{b}$ we have

$$\mathfrak{a}\mathfrak{b} = (b)$$

for some \mathfrak{a} . Note $\mathfrak{a} \sim \mathfrak{b}^{-1} \sim \mathfrak{c}$, and taking norms of the above equation gives

$$\mathfrak{N}\mathfrak{a}\mathfrak{N}\mathfrak{b} = \mathfrak{N}(b) \leq \mu\mathfrak{N}\mathfrak{b}.$$

Canceling $\mathfrak{N}\mathfrak{b}$ gives that \mathfrak{a} is the desired representative. □

We give an example of computing the class group. The general procedure to compute the class group of $A = \mathcal{O}_K$ where $K = \mathbb{Q}(\sqrt{d})$ and d is negative and squarefree is as follows.

1. List the primes $p \leq \lfloor \mu \rfloor$.
2. For each p , determine whether p splits in A by checking whether

$$f(x) := \begin{cases} x^2 - x + \frac{d-1}{4}, & d \equiv 1 \pmod{4} \\ x^2 - d, & d \equiv 2, 3 \pmod{4} \end{cases}$$

is irreducible.

3. If $p = \mathfrak{a}\bar{\mathfrak{a}}$ splits in A , include it in the list of generators.
4. Compute the norm of some small elements (with prime divisors in the list found above), like $k + \delta$ for $k \in \mathbb{N}_0$, $\delta = \sqrt{d}$ or $\frac{1+\sqrt{d}}{2}$ depending on $d \pmod{4}$. Factor $\text{Nm}_{K/\mathbb{Q}}(a)$ to factor

$$(a)(\bar{a}) = (\text{Nm}_{K/\mathbb{Q}}(a));$$

match factors using unique factorization. Note $(a) \sim (\bar{a}) \sim 1$. Repeat until there are enough relations to determine the group.

5. For the prime 2, if $d \equiv 2, 3 \pmod{4}$, 2 ramifies, $(2) = \mathfrak{p}^2$, and \mathfrak{p} has order 2 for $d \neq -1, -2$. (Note $\mathfrak{p} = (2, \delta)$ and $(2, 1 + \delta)$ in these two cases, respectively.)

We first consider the cases when the class group is trivial.

Theorem 3.4.2. *The rings*

$$\mathbb{Z}[\sqrt{-1}], \mathbb{Z}[\sqrt{-2}], \mathbb{Z}\left[\frac{1+\sqrt{d}}{2}\right], d = -3, -7, -19, -43, -67, -163$$

are unique factorization domains.

In fact, they are the only ones (part of Gauss's class number problem).

Proof. Note $\mathbb{Z}[\sqrt{-1}]$, $\mathbb{Z}[\sqrt{-2}]$, and $\mathbb{Z}\left[\frac{1+\sqrt{d}}{2}\right]$ are Euclidean domains and hence unique factorization domains.

The class group of $\mathbb{Z}\left[\frac{1+\sqrt{d}}{2}\right]$ is generated by the classes of prime ideals whose norms are prime integers $p \leq \mu$, which are the factors of (p) when it splits. When $d \equiv 1 \pmod{4}$ as in all the remaining cases, an integer prime p remains prime in $\mathbb{Z}\left[\frac{1+\sqrt{-d}}{2}\right]$ iff $x^2 - x - \frac{1}{4}(1-d)$ is irreducible modulo p , iff $x^2 - x - \frac{1}{4}(1-d)$ has no zero modulo p . We show that for $d = -7, -11, -19, -43, -67, -163$, $x^2 - x - \frac{1}{4}(1-d)$ is irreducible modulo all primes less than μ . Then no prime ideals have norms that are prime integers $p \leq \mu$, and the only ideal class is that of the principal ideals. It follows that $\mathbb{Z}[\sqrt{d}]$ is a principal ideal domain and hence a unique factorization domain.

d	$\lfloor \mu \rfloor, \mu = \sqrt{\frac{ d }{3}}$	$x^2 - x + \frac{1}{4}(1-d)$	Primes $p \leq \lfloor \mu \rfloor, x^2 - x + \frac{1}{4}(1-d) \pmod{p}$
-7	$\lfloor \sqrt{\frac{7}{3}} \rfloor = 1$		None
-11	$\lfloor \sqrt{\frac{11}{3}} \rfloor = 1$		None
-19	$\lfloor \sqrt{\frac{19}{3}} \rfloor = 2$	$x^2 - x + 5$	2: $x^2 + x + 1 = 1$ for $x = 0, 1$
-43	$\lfloor \sqrt{\frac{43}{3}} \rfloor = 3$	$x^2 - x + 11$	2: $x^2 + x + 1 = 1$ for $x = 0, 1$ 3: $x^2 - x - 1 = \begin{cases} -1 & \text{for } x = 0, 1 \\ 1 & \text{for } x = 2 \end{cases}$
-67	$\lfloor \sqrt{\frac{67}{3}} \rfloor = 4$	$x^2 - x + 17$	2: $x^2 + x + 1 = 1$ for $x = 0, 1$ 3: $x^2 - x - 1 = \begin{cases} -1 & \text{for } x = 0, 1 \\ 1 & \text{for } x = 2 \end{cases}$
-163	$\lfloor \sqrt{\frac{163}{3}} \rfloor = 7$	$x^2 - x + 41$	2: $x^2 + x + 1 = 1$ for $x = 0, 1$ 3: $x^2 - x - 1 = \begin{cases} -1 & \text{for } x = 0, 1 \\ 1 & \text{for } x = 2 \end{cases}$ 5: $x^2 - x + 1 = \begin{cases} 1 & \text{for } x = 0, 1 \\ 3 & \text{for } x = 4, 2 \\ 2 & \text{for } x = 3 \end{cases}$ 7: $x^2 - x - 1 = \begin{cases} -1 & \text{for } x = 0, 1 \\ 1 & \text{for } x = 2, 6 \\ 5 & \text{for } x = 3, 5 \\ 4 & \text{for } x = 4 \end{cases}$

□

Example 3.4.3: We compute the class group of $\mathbb{Z}[\sqrt{-41}]$.

For $d = -41$, $\lfloor \mu \rfloor = \lfloor 2\sqrt{\frac{41}{3}} \rfloor = 7$. Modulo 2, 3, 5, and 7, -41 is congruent to 1, 1, 4, and 1, which are all squares. Factor

$$\begin{aligned} (2) &= A\bar{A} \\ (3) &= B\bar{B} \\ (5) &= C\bar{C} \end{aligned}$$

$$(7) = D\bar{D}$$

Then the class group is generated by $\langle A \rangle, \langle B \rangle, \langle C \rangle, \langle D \rangle$. (Note that $\langle \bar{A} \rangle = \langle A \rangle^{-1}$, etc.) We have

$$(1 + \delta)(\overline{1 + \delta}) = (42) = (2)(3)(7) = A\bar{A}B\bar{B}D\bar{D}.$$

If a prime ideal P divides $(1 + \delta)$ then \bar{P} divides $(\overline{1 + \delta})$. Hence the conjugate factors are divided between $(1 + \delta)$ and $(\overline{1 + \delta})$. Without loss of generality, we can suppose

$$(1 + \delta) = ABD.$$

The class of a principal ideal is the identity in the class group, so

$$\langle A \rangle \langle B \rangle \langle D \rangle = 1. \quad (3.9)$$

Next consider

$$(2 + \delta)(\overline{2 + \delta}) = (45) = (3)^2(5) = B^2\bar{B}^2C\bar{C}.$$

Note that 3 does not divide $2 + \delta$ so $B\bar{B} = (3)$ doesn't divide $(2 + \delta)$. Thus B^2, \bar{B}^2 divide $(2 + \delta), (\overline{2 + \delta})$ in some order. Since we haven't distinguished between C and \bar{C} yet, we may assume WLOG that $\langle B \rangle^2 \langle \bar{C} \rangle, \langle \bar{B} \rangle^2 \langle C \rangle$ are equal to $(2 + \delta)$ and $(\overline{2 + \delta})$ in some order, and

$$\langle B \rangle^2 \langle \bar{C} \rangle = \langle B \rangle^2 \langle C \rangle^{-1} = 1$$

or

$$\langle C \rangle = \langle B \rangle^2. \quad (3.10)$$

Similarly, looking at

$$(3 + \delta)(\overline{3 + \delta}) = (50) = (2)(5)^2 = A\bar{A}C^2\bar{C}^2,$$

we get that

$$\langle A \rangle \langle \bar{C} \rangle^2 = 1 \text{ or } \langle \bar{A} \rangle \langle C \rangle^2 = 1.$$

Noting that $A = \bar{A}$ (since $(2) = (2, 1 + \delta)(2, 1 - \delta)$ and $(2, 1 + \delta) = (2, 1 - \delta)$ when $d \equiv 3 \pmod{4}$) by [Artin, 13.8.4],

$$\langle A \rangle = \langle C \rangle^2. \quad (3.11)$$

From (3.10) we may omit $\langle C \rangle$ from the list of generators for the group, from (3.11) we may omit $\langle A \rangle$, and from (3.9) we may omit $\langle D \rangle$. Thus the class group is the cyclic group generated by $\langle B \rangle$. From (3.10) and (3.11), we get

$$\langle A \rangle = \langle B \rangle^4. \quad (3.12)$$

Since A is not principal, $\langle B \rangle^4 \neq 1$. Note $\langle A \rangle = \langle \bar{A} \rangle = \langle A \rangle^{-1}$ implies $\langle A \rangle^2 = 1$. Combining this with (3.12) gives that $\langle B \rangle^8 = 1$. Since $\langle B \rangle^n \neq 1$ for any proper divisor n of 8 (it sufficed to check $n = 4$), the class group is cyclic of order 8, C_8 .

Chapter 4

The algebra of quadratic forms

We follow Cox [Cox89], except for the proof of Gauss composition, when we follow Cassels (add reference). The last section is based on Bhargava's paper [Bha04].

1 Quadratic forms

Definition 4.1.1: Let R be an integral domain. A **quadratic form** on R is a function on R^n , in the form

$$f(x_1, \dots, x_n) = \sum_{1 \leq i < j \leq n} a_{ij} x_i x_j.$$

Supposing R is a UFD, we say f is **primitive** iff $\gcd_{1 \leq i < j \leq n} a_{ij} = 1$.

A quadratic form may be represented by a matrix

$$Q = \begin{bmatrix} a_{11} & \frac{a_{12}}{2} & \cdots & \frac{a_{1,n-1}}{2} & \frac{a_{1,n}}{2} \\ \frac{a_{12}}{2} & a_{22} & \cdots & \frac{a_{2,n-1}}{2} & \frac{a_{2,n}}{2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \frac{a_{1,n-1}}{2} & \frac{a_{2,n-1}}{2} & \cdots & a_{n-1,n-1} & \frac{a_{n-1,n}}{2} \\ \frac{a_{1,n}}{2} & \frac{a_{2,n}}{2} & \cdots & \frac{a_{n-1,n}}{2} & a_{nn} \end{bmatrix}$$

(working in $K = \text{Frac}(R)$ as necessary to allow division by 2); we have

$$f(\mathbf{x}) = \mathbf{x}Q\mathbf{x}^T.$$

Definition 4.1.2: We say two forms f and g are **equivalent** if there is an invertible matrix A (i.e. a matrix with determinant a unit) such that

$$f(\mathbf{x}) = g(\mathbf{x}A^T).$$

We say f and g are **properly equivalent** if $\det(A) = 1$.

Note that the matrices corresponding to f and g are related by

$$Q_f = A^T Q_g A.$$

For the rest of this chapter, we will focus on integral binary quadratic forms, i.e. those in two variables over \mathbb{Z} .

2 Representing integers

Definition 4.2.1: We say that f **represents** n if there exists $\mathbf{x} = (x_1, \dots, x_n)$ such that $f(\mathbf{x}) = n$. We say that f **properly represents** n if we can choose \mathbf{x} so that $\gcd(x_1, \dots, x_n) = 1$.

Lemma 4.2.2. *A form $f(x, y)$ properly represents n if and only if $f(x, y)$ is properly equivalent to the form $nx^2 + b'xy + c'y^2$ for some $b', c' \in \mathbb{Z}$.*

Proof. If $f(p, q) = n$ with (p, q) relatively prime, then by Bézout we can find r, s such that $ps - qr = 1$. Let $f(x, y) = ax^2 + bxy + cy^2$. Then f is equivalent to

$$f(px + ry, qx + sy) = \underbrace{f(p, q)}_n x^2 + (2apr + bps + brq + 2cqs)xy + f(r, s)y^2.$$

For the converse, note that $nx^2 + bxy + cy^2$ properly represents n by taking $(x, y) = (1, 0)$. □

Theorem 4.2.3. *Let $n \neq 0$ and d be integers. Then the following are equivalent.*

1. *There exists a binary quadratic form of discriminant d which properly represents n .*
2. *d is square modulo $4n$.*

Proof. Suppose f is a binary quadratic form of discriminant d properly representing n . Then by Lemma 4.2.2, f is equivalent to some form $nx^2 + bxy + cy^2$. Hence the discriminant is $d = b^2 - 4nc$, and $d \equiv b^2 \pmod{4n}$.

Conversely, suppose $b^2 \equiv d \pmod{4n}$, so $b^2 = d + 4nc$ for some integer n , i.e. $d = b^2 - 4nc$. Then

$$f(x, y) = nx^2 + bxy + cy^2$$

properly represents n , as $f(1, 0) = n$, and $\text{disc}(f) = b^2 - 4nc = d$. □

Corollary 4.2.4. *Let n be an integer and p an odd prime not representing n . Then $\left(\frac{-n}{p}\right) = 1$ iff p is represented by a primitive form of discriminant $-4n$.*

Proof. Note $\left(\frac{-n}{p}\right) = 1$ iff $\left(\frac{-4n}{p}\right) = 1$, and this is equivalent to the second statement by the theorem. □

The results in this section are particularly useful if there are few quadratic forms with determinant d . There is a method to list all these quadratic forms, as we will show in the next section.

3 Reduction of quadratic forms

We would like to have a canonical representative for every equivalence class of binary quadratic forms. We choose the one with “smallest” coefficients. This is made precise by the following definition.

Definition 4.3.1: A positive definite binary quadratic form $ax^2 + bxy + cy^2$ is **reduced** if it is primitive and

$$|b| \leq a \leq c$$

and

$$b \geq 0 \text{ if } |b| = a \text{ or } a = c.$$

Theorem 4.3.2. *Every equivalence class of primitive binary quadratic forms contains exactly one reduced form.*

Proof. Existence, Step 1: We first show there is a form in the class with $|b| \leq a \leq c$.

Take the form $f(x) = ax^2 + bxy + cy^2$ in the equivalence class such that $|b|$ is smallest. Note $a, c > 0$ because the form is positive definite. We claim that $a, c \geq |b|$. Indeed, we have

$$f(x + my, y) = ax^2 + (2am + b)xy + (am^2 + c)y^2,$$

so $-b \leq 2am + b \leq b$ for all $m \in \mathbb{Z}$, giving $a \geq |b|$. Similarly, $c \geq |b|$.

Next, if $a > c$, then replacing (x, y) by $(-y, x)$ we get $c > a \geq |b|$.

Step 2: The form is reduced unless $b < 0$ and $a = -b$ or $a = c$. We tackle these cases next. In these cases $ax^2 - bxy + cy^2$ is reduced, so it suffices to show $ax^2 \pm bxy + cy^2$ are equivalent. In these two cases we make the following substitutions:

$$\begin{aligned} f(x, y) = ax^2 - axy + cy^2 &\implies f(x + y, y) = ax^2 + axy + cy^2 \\ f(x, y) = ax^2 + bxy + ay^2 &\implies f(-y, x) = ax^2 - bxy + ay^2. \end{aligned}$$

Uniqueness, Step 1: We claim that for $(x, y) \in \mathbb{Z}^2$ with $xy \neq 0$, and $f(x, y) = ax^2 + bxy + cy^2$ with $a, c \geq |b|$, we have

$$f(x, y) \geq (a - |b| + c) \min(x^2, y^2).$$

Indeed, without loss of generality assume $x \geq y$. Then

$$f(x, y) \geq (a - |b|)xy + cy^2 \geq (a - |b| + c)y^2.$$

As a corollary, for $xy \neq 0$,

$$ax^2 + bxy + cy^2 \geq a - |b| + c$$

with equality iff $x, y = \pm 1$, $xy = -\text{sign}(b)$.

Step 2: To distinguish between reduced forms, we examine the smallest nonzero values attained by a them, and the number of primitive solutions to them. Note all solutions (x, y) with $xy = 0$ and one of $|x|, |y| \geq 2$ are removed from consideration.

1. If $|b| < a < c$, then the smallest values attained by f primitively are

$$a < c < a - |b| + c$$

with solutions $(\pm 1, 0)$, $(0, \pm 1)$ and $\pm(-1, \text{sign}(b))$ respectively.

2. If $b \geq 0$ and $|b| = a < c$, then the smallest values attained by f primitively are

$$a < c = a - |b| + c;$$

the first has 2 solutions and the latter has 4 primitive solutions.

3. If $b \geq 0$ and $|b| < a = c$, then the smallest values attained by f primitively are

$$a = c < a - |b| + c;$$

the first has 4 solutions and the latter has 2 primitive solutions.

4. If $b \geq 0$ and $|b| = a = c$, then the smallest value attained by f primitively is

$$a = c = a - |b| + c$$

which has 6 primitive solutions.

After examining this data, the only reduced forms that could be equivalent are those falling in the first category with opposite b 's, i.e. $ax^2 \pm bxy + cy^2$. But any change of variables sending one to the other must preserve the solutions $(\pm 1, 0)$ and $(0, \pm 1)$, so must have matrix $\begin{pmatrix} \pm 1 & 0 \\ 0 & \pm 1 \end{pmatrix}$. If this matrix has determinant 1, then it must be $\pm I$ and cannot change between the two forms. \square

Suppose $d < 0$; note that there is an algorithm to list all reduced quadratic forms with discriminant d . The conditions $|b| \leq a \leq c$ and $b^2 - 4ac = d$ give

$$d = b^2 - 4ac \leq a^2 - 4a^2 = -3a^2.$$

Hence

$$a \leq \sqrt{-\frac{d}{3}}.$$

We simply check for solutions to $b^2 - 4ac = d$ for all $0 \leq |b| \leq a \leq \sqrt{-\frac{d}{3}}$.

3.1 Examples

Example 4.3.3: When $n = 1, 2, 3$, the above check gives that the only reduced form of discriminant $-4n$ is $x^2 + ny^2$.

Combining this fact with Theorem 4.2.3, we get that f properly represents m iff $d := -4n$ is a square modulo $4m$, i.e. -1 is a square modulo m . Thus we have the chain of equivalences:

1. f represents m .
2. f properly represents $\frac{m}{k^2}$ for some square factor $k^2 \mid m$.
3. d is a square modulo $\frac{m}{k^2}$ for some m .
4. d is a square modulo $\frac{m}{k^2}$ for the largest square factor $k^2 \mid m$.
5. d is a square modulo p for every $p \mid m$ with $\text{ord}_p(m)$ odd.

By quadratic reciprocity, we have

$$\begin{aligned} \left(\frac{-1}{p}\right) &= (-1)^{\frac{p-1}{2}} = \begin{cases} 1, & p \equiv 1 \pmod{4} \\ -1, & p \equiv 3 \pmod{4} \end{cases} \\ \left(\frac{-2}{p}\right) &= (-1)^{\frac{p-1}{2}} (-1)^{\frac{p^2-1}{8}} = \begin{cases} 1, & p \equiv 1, 3 \pmod{8} \\ -1, & p \equiv 5, 7 \pmod{8} \end{cases} \\ \left(\frac{-3}{p}\right) &= (-1)^{\frac{p-1}{2}} (-1)^{\frac{3-1}{2} \cdot \frac{p-1}{2}} \left(\frac{p}{3}\right) = \begin{cases} 1, & p \equiv 1 \pmod{3} \\ -1, & p \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

Hence we have the following.

m represented by	iff every such prime has even exponent in m
$x^2 + y^2$	$p \equiv 3 \pmod{4}$
$x^2 + 2y^2$	$p \equiv 5, 7 \pmod{8}$
$x^2 + 3y^2$	$p \equiv 2 \pmod{3}$

Compare this with the proof using factorization in $\mathbb{Z}[\sqrt{-d}]$.¹ In particular, note that $\mathbb{Z}[\sqrt{-d}]$ is a UFD when $d = 1, 2$, and in these cases, there is exactly one form of discriminant $-4d$. *This is not a coincidence!*

Next we show the following.

Example 4.3.4: A positive integer n is represented by $x^2 + 5y^2$ iff

1. Any prime $p \equiv 11, 13, 17, 19 \pmod{20}$ appears in n with even exponent.
2. There are an even number of prime divisors that are $p \equiv 2, 3, 7 \pmod{20}$, counting multiplicity.
3. (No restriction on primes $p \equiv 1, 5, 9 \pmod{20}$.)

Note this condition is quite different from the ones before!

¹When $d = 3$ we have to be slightly careful.

Proof 1. This time we have to check $a \leq \sqrt{-\frac{20}{3}} < 3$. The reduced forms of discriminant -20 are

$$\begin{aligned} f(x) &:= x^2 + 5y^2 \\ g(x) &= 2x^2 + 2xy + 3y^2. \end{aligned}$$

We run into trouble already: Theorem 4.2.3 fails to distinguish between these. We still start with the same argument, though.

Step 1: By Corollary 4.2.3, a prime p is represented by f or g iff $\left(\frac{-5}{p}\right) = 1$. By quadratic reciprocity,

$$\left(\frac{-5}{p}\right) = (-1)^{\frac{p-1}{2}} \left(\frac{p}{5}\right) = \begin{cases} 1, & p \equiv 1, 3, 7, 9 \pmod{20} \\ -1, & p \equiv 11, 13, 17, 19 \pmod{20}. \end{cases}$$

Step 2: Now we distinguish between these two cases. By checking modulo 4, we see that f only represents primes $p \equiv 1, 9 \pmod{20}$ (and 5) and g only represents primes $p \equiv 3, 7 \pmod{20}$ (and 2).² By Step 1, f, g must represent all of these respective primes.

Step 3: We have the desired result for primes. How to pass to products of primes? First note that primes $p \equiv 11, 13, 17, 19 \pmod{20}$ have to appear with even exponent (if $x^2 + 5y^2 \equiv 0 \pmod{p}$), since $\left(\frac{-5}{p}\right) = -1$, we must have $p \mid x, y$; now divide x, y by p and repeat).

Now consider the magical identity

$$(2x^2 + 2xy + 3y^2)(2z^2 + 2zw + 3w^2) = (2xy + xw + yz + 3yw)^2 + 5(xw - yz)^2, \quad (4.1)$$

which says that a product of numbers represented by g is represented by f ! This immediately gives the sufficiency condition.

For the necessary condition, note we may divide x, y by 2 until they are not both even. Now take it modulo 8 to see that $n \equiv 1, 4, 5, 6 \pmod{8}$. This gives that item 2 is necessary. \square

Wait a minute. Where does the magical identity come from? Historically this was the way such problems were solved, and in fact the motivation for *composing* quadratic forms: for primitive quadratic forms f, g, h , we say $f \circ g = h$ iff there exist integral bilinear forms B_1, B_2 satisfying certain conditions such that

$$f(\mathbf{x})g(\mathbf{y}) = h(B_1(\mathbf{x}, \mathbf{y}), B_2(\mathbf{x}, \mathbf{y})).$$

We won't go into the historical details, because the modern way of thinking of composition is cleaner (see Section 5). We know we had the "composition law"

$$(a^2 + b^2)(c^2 + d^2) = (ac - bd)^2 + (ad + bc)^2.$$

²These sets are disjoint; we say f, g are unique in their *genus*.

We can view this as coming from the identity

$$\mathrm{Nm}_{K/\mathbb{Q}}(a + bi) \mathrm{Nm}_{K/\mathbb{Q}}(c + di) = \mathrm{Nm}_{K/\mathbb{Q}}((a + bi)(c + di)) \quad (4.2)$$

where $K = \mathbb{Q}(i)$, so $\mathrm{Nm}_{K/\mathbb{Q}}(z) = |z|^2$. We now look at a different proof of Example 4.3.4.

Proof 2. This time the complication comes from that $\mathbb{Z}[\sqrt{-5}]$ is not a UFD, nor PID; its ideal class group has order 2, with representatives

$$\begin{aligned} \mathfrak{a} &= 1 \\ \mathfrak{b} &= (3, 1 + \sqrt{-5}). \end{aligned}$$

Step 1: Let p be prime. As in the proof of Theorem 3.2.1, we factor the equation $x^2 + 5y^2 = p$ in $\mathbb{Z}[\sqrt{-5}]$ to get

$$(x + \sqrt{-5}y)(x - \sqrt{-5}y) = p.$$

Now we know the ideal (p) splits iff $x^2 + 5 \pmod{p}$ splits, i.e. $\left(\frac{-5}{p}\right) = 1$. We calculated that this happens when $p \equiv 1, 3, 5, 7, 9 \pmod{20}$.

Step 2: So if p is of the above form, we know that either p is a product of two principal ideals, or two (conjugate) ideals similar to \mathfrak{b} . In the two cases, we have respectively

$$\begin{aligned} (p) &= (\lambda)(\bar{\lambda}) \\ (p) &= \lambda(3, 1 + \sqrt{-5})\bar{\lambda}(3, 1 + \sqrt{-5}) \end{aligned}$$

for some $\lambda \in \mathbb{Q}(\sqrt{-5})$. Then calculating the norm of the ideal in $K = \mathbb{Q}(\sqrt{-5})$ gives

$$\begin{aligned} p &= \mathrm{Nm}_{K/\mathbb{Q}}(\lambda) \\ p &= \underbrace{\mathfrak{N}((3, 1 + \sqrt{-5}))}_3 \mathrm{Nm}_{K/\mathbb{Q}}(\lambda)^2. \end{aligned}$$

Let $\lambda = x + y\sqrt{-5}$. In the first case, we must have $p = x^2 + 5y^2$, so $p \equiv 1, 5, 9 \pmod{20}$, while in the second case, we must have $p = 3(x^2 + 5y^2)$ ($x, y \in \mathbb{Q}$, here) so when p is odd, $p \equiv 3 \cdot 1, 3 \cdot 9 \pmod{20}$. (We can check that x, y do not have 2 or 5 in the denominator by an infinite descent argument, so we may consider $x, y \in \mathbb{Z}/20\mathbb{Z}$.) $p = 2$ is possible as $(2, 1 + \sqrt{-5})^2 = (2)$. Thus again we've distinguished between the two cases.

Step 3: A prime $p \equiv 1, 5, 9 \pmod{20}$ splits into two principal ideals, a prime $p \equiv 2, 3, 7 \pmod{20}$ splits into two ideals of type \mathfrak{b} , and a prime $p \equiv 11, 13, 17, 19 \pmod{20}$ remains prime. In order for (n) to split into two principal ideals, we must be able to write

$$(n) = \mathfrak{c}\bar{\mathfrak{c}}$$

where \mathfrak{c} is a product of ideals, containing an *even* number of prime ideals of type \mathfrak{b} , and $\bar{\mathfrak{c}}$ contains the conjugates of those ideals. (Two ideals of type \mathfrak{b} multiply to a principal ideal.) The result follows. \square

It seems like the quadratic forms in the first proof are related to the ideals in the second proof. This is indeed the case: we can explain (4.1) similarly to (4.2) by

$$\begin{aligned} & \frac{\text{Nm}_{K/\mathbb{Q}}(2x + (1 + \sqrt{-5})y)}{\mathfrak{N}(2, 1 + \sqrt{-5})} \cdot \frac{\text{Nm}_{K/\mathbb{Q}}(2z + (1 + \sqrt{-5})w)}{\mathfrak{N}(2, 1 + \sqrt{-5})} \\ &= \frac{\text{Nm}_{K/\mathbb{Q}}((2x + (1 + \sqrt{-5})y)(2z + (1 + \sqrt{-5})w))}{\mathfrak{N}((2))} \end{aligned}$$

The two forms on the LHS are exactly those on the LHS of (4.1) while that on the RHS can be written in the form $B_1^2 + 5B_2^2$ because $\frac{1}{2}(2x + (1 + \sqrt{-5})y)(2z + (1 + \sqrt{-5})w)$ is an integral ideal. We will see that in this way the group law on ideal classes translates into a group law on quadratic forms.

After we establish Gauss composition, we will show the equivalence between a quadratic form Q representing a prime p , and (p) splitting into ideals of a certain form (Theorem 4.5.4). The above proof was a specific example of this.

4 Ideals on quadratic rings

Definition 4.4.1: We will be considering rings that are free \mathbb{Z} -modules of finite rank. We call such rings **quadratic**, **cubic**, **quartic**, and **quintic**, if the rank is 2, 3, 4, or 5, respectively.

The rings we are primarily interested are integral domains, which are exactly the rings that can be embedded in field extensions.

Definition 4.4.2: An **order** \mathcal{O} in a finite extension K/\mathbb{Q} is a subring of K containing 1, that is a free \mathbb{Z} -module of rank $[K : \mathbb{Q}]$.

The maximal order of K is simply \mathcal{O}_K , the ring of integers of K .

Definition 4.4.3: Let R be a ring that is a free \mathbb{Z} -module of finite rank. The **conductor** of R is the greatest integer n for which there exists a ring T such that

$$\mathcal{O} = \mathbb{Z} + nT.$$

(Necessarily, T has the same rank.)

If S is a quadratic ring then $S = \langle 1, \tau \rangle$ for some τ satisfying a quadratic equation $\tau^2 + b\tau + c = 0$. If this polynomial is irreducible over \mathbb{Z} , then S can be embedded in a quadratic field extension. Otherwise, S is not an integral domain. We make the following definitions. The first four are equivalent to our previous definitions when S is integrally closed.

1. The discriminant of S is the discriminant of the characteristic polynomial, $b^2 - 4c$.

2. Conjugation is the linear transformation that takes 1 to 1 and switches the zeros of $x^2 + bx + c$.
3. The norm of an element $\alpha \in S$ is $\alpha\bar{\alpha}$.
4. The numerical norm $\mathfrak{N}_R(\mathfrak{a})$ of an ideal $\mathfrak{a} \in R$ to be $[R : I] = |R/I|$.³
5. A basis (α, β) for $\mathfrak{a} \subseteq R$ is positively oriented if

$$\frac{\begin{vmatrix} \alpha & \bar{\alpha} \\ \beta & \bar{\beta} \end{vmatrix}}{\text{disc}(S)} = \frac{\alpha\bar{\beta} - \beta\bar{\alpha}}{d} > 0.$$

We now describe all quadratic rings.

Proposition 4.4.4: There is a bijection between $D = \{d \in \mathbb{Z} : d \equiv 0, 1 \pmod{4}\}$ and quadratic rings (up to isomorphism), given by

$$S : d \mapsto \mathbb{Z}[\tau_d]$$

where τ_d satisfies a monic quadratic equation with discriminant d .

Moreover,

$$d = f^2 d_K,$$

where f is the conductor of $\mathbb{Z}[\tau_d]$ and, when d is nonsquare, d_K is the discriminant of $\mathbb{Q}(\tau_d)$ ($d_K \equiv 0, 1 \pmod{4}$ and $16 \nmid d_K$).

1. An integer $d \in D$ corresponds to a integral domain if and only if d is not a square.
2. If $d = 0$ then $S(d) = \mathbb{Z}[x]/(x^2)$.
3. If d is a nonzero square then $S(d) = \mathbb{Z} \cdot (1, 1) + \sqrt{d}(\mathbb{Z} \oplus \mathbb{Z})$.
4. If $d_K \equiv 1 \pmod{4}$, $d_K \neq 1$, then $S(d) = \mathbb{Z}[f\tau] = \langle 1, f\tau \rangle$ where $\tau = \frac{1+\sqrt{d_K}}{2}$.
5. If $d_K \equiv 0 \pmod{4}$ then $S(d) = \mathbb{Z}[f\tau] = \langle 1, f\tau \rangle$ where $\tau = \frac{\sqrt{d_K}}{2}$ —the root of the nonsquare part of d .

Proof. Note the map is well-defined, because any two quadratic equations with discriminant d , say $x^2 + b_j x + c_j$, $j = 1, 2$, have $b_1 \equiv b_2 \equiv d \pmod{2}$ and hence are related by the change of variable $x \mapsto x + k$ for some k . The map is injective because the discriminant doesn't change under replacing τ with $\tau + k$.

³For fractional ideals \mathfrak{a} , i.e. R -submodules of $R \otimes_{\mathbb{Z}} \mathbb{Q}$, take a fractional ideal \mathfrak{b} containing \mathfrak{a} and R and define $\mathfrak{N}_R(\mathfrak{a}) = \frac{[\mathfrak{b}:\mathfrak{a}]}{[\mathfrak{b}:R]}$.

For item 1, note d is a square iff the characteristic polynomial factors. Item 2 is clear; for item 3 note that we have the homomorphism

$$\begin{aligned} \mathbb{Z}[\tau]/(\tau^2 - d) &\hookrightarrow \mathbb{Z}[\tau]/(\tau - \sqrt{d}) \times \mathbb{Z}[\tau]/(\tau + \sqrt{d}) \cong \mathbb{Z} \times \mathbb{Z} \\ 1 &\mapsto (1, 1) \\ \tau &\mapsto (\sqrt{d}, -\sqrt{d}) \end{aligned}$$

with image $\mathbb{Z} \cdot (1, 1) + \sqrt{d}(\mathbb{Z} \oplus \mathbb{Z})$.

Now write $d = f^2 d_K$; we will show f is the conductor. Choose $b = 0$ or 1 with $b \equiv d \pmod{4}$ and c such that $b^2 - 4c = d$, and let

$$\begin{aligned} S(d_K) &= \mathbb{Z}[\tau]/(\tau^2 + b\tau + c) = \mathbb{Z} \left[\frac{-b + \sqrt{d_K}}{2} \right] \\ S(d) &= \mathbb{Z}[\tau]/(\tau^2 + fb\tau + fc) = \mathbb{Z} \left[\frac{-fb + f\sqrt{d_K}}{2} \right]. \end{aligned}$$

Now $S(d_K)$ is the ring of integers of $S(d)$, so the largest quadratic ring containing $S(d)$; moreover the above representation gives

$$S(d) = \mathbb{Z} + fS(d_K), \tag{4.3}$$

so f must be the conductor.

Items 4 and 5 come from (4.3) and the fact that $\mathbb{Z} \left[\frac{-b + \sqrt{d_K}}{2} \right] = \mathbb{Z}[\tau_K]$. \square

4.1 Proper and invertible ideals

From now on, assume that d is not a square. We create a bijection between the “ideal class group” of a quadratic ring of discriminant d and quadratic forms of discriminant d . To do this we first have to define the “ideal class group” of a quadratic ring. This is more complicated than defining it for a ring of integers, because a general order is not a Dedekind domain. We find that we first have to restrict the ideals under consideration, in order for inverses to exist.⁴ Later we restrict the ideals further so that we have unique factorization.

Definition 4.4.5: A **proper ideal** of \mathcal{O} is an ideal such that

$$\mathcal{O} = \{\beta \in K : \beta\mathfrak{a} \subseteq \mathfrak{a}\}.$$

(In general we only have \subseteq .)

Note that for the maximal order \mathcal{O}_K , all ideals are proper, and for any order, all principal ideals are proper. Furthermore, any ideal is proper for exactly most one order, namely the order $\{\beta \in K : \beta\mathfrak{a} \subseteq \mathfrak{a}\}$. The following tells us exactly which order that is.

⁴Else we only get a semigroup.

Lemma 4.4.6. *Suppose $\mathfrak{a} = (\alpha, \beta)$ is an ideal in a order of a quadratic field.*

Suppose $\tau = \frac{\beta}{\alpha}$ has degree 2 over \mathbb{Q} and satisfies the equation

$$ax^2 + bx + c = 0$$

where $a > 0$, b , and c are integers with $\gcd(a, b, c) = 1$. Let $K = \mathbb{Q}(\tau)$. Then \mathfrak{a} is a proper ideal of $R := (1, a\tau)$, and

$$\mathfrak{N}_R(\mathfrak{a}) = \frac{\text{Nm}_{K/\mathbb{Q}}(\alpha)}{a}.$$

Proof. Let \mathcal{O} be the order. Now $(1, \tau)$ is also a fractional ideal of $\mathcal{O} \subseteq \mathbb{Q}(\tau)$. We know $\mathcal{O} = \{\beta \in K : \beta\mathfrak{a} \subseteq \mathfrak{a}\}$. Now, β is in this set iff

$$\begin{aligned} \beta &\in (1, \tau) \\ \beta\tau &\in (1, \tau), \end{aligned}$$

i.e.

$$\begin{aligned} \beta &= p + q\tau \text{ for some } p, q \in \mathbb{Z} \\ \beta\tau &= (p + q\tau)\tau = p\tau + q \left(-\frac{b}{a}\tau - \frac{c}{a} \right) \in (1, \tau); \end{aligned}$$

since $\gcd(a, b, c) = 1$, this is true iff $a \mid q$. Hence $\mathcal{O} = (1, a\tau)$.

For the second part, note

$$\mathfrak{N}(\mathfrak{a}) = [\mathcal{O} : \mathfrak{a}] = \frac{[\mathcal{O} : (1, \tau)]}{[\mathfrak{a} : (1, \tau)]} = \frac{[\alpha(1, \tau) : (1, \tau)]}{[(1, a\tau) : (1, \tau)]} = \frac{\text{Nm}(\alpha)}{a}.$$

□

Proposition 4.4.7: Let \mathfrak{a} be a fractional \mathcal{O} -ideal. Then \mathfrak{a} is proper iff it is invertible. Hence the proper fractional ideals form a group $I(\mathcal{O})$ under multiplication.

Proof. If \mathfrak{a} is invertible, then $\mathfrak{a}\mathfrak{b} = \mathcal{O}$ for some \mathfrak{b} . If $\beta\mathfrak{a} \subseteq \mathfrak{a}$, then

$$\beta\mathcal{O} = \beta(\mathfrak{a}\mathfrak{b}) = (\beta\mathfrak{a})\mathfrak{b} \subseteq \mathfrak{a}\mathfrak{b} = \mathcal{O}$$

so $\beta \in \mathcal{O}$. This shows \mathfrak{a} is proper.

Conversely, suppose \mathfrak{a} is proper. Write $\mathfrak{a} = \alpha(1, \tau)$. Letting $ax^2 + bx + c$ be the minimal polynomial of τ with integer coefficients, by Lemma 4.4.6, $\mathcal{O} = (1, a\tau)$. We show that

$$\mathfrak{a}\bar{\mathfrak{a}} = \frac{\text{Nm}_{K/\mathbb{Q}}(\alpha)}{a}\mathcal{O};$$

it will follow that $\frac{a}{\text{Nm}_{K/\mathbb{Q}}(\alpha)}\bar{\mathfrak{a}}$ is the inverse of \mathfrak{a} .

First note $\mathcal{O} = \overline{\mathcal{O}}$, since $\mathcal{O} = (1, a\tau) = (1, a\bar{\tau})$ (on account of $a\tau + a\bar{\tau} = -b$). Hence $\bar{\mathfrak{a}}$ is actually an ideal of \mathcal{O} . Next, we calculate

$$\begin{aligned} \mathfrak{a}\bar{\mathfrak{a}} &= \alpha(1, \tau)\bar{\alpha}(1, \bar{\tau}) \\ &= \text{Nm}_{K/\mathbb{Q}}(\alpha)(1, \tau, \bar{\tau}, \tau\bar{\tau}) \\ &= \text{Nm}_{K/\mathbb{Q}}(\alpha) \left(1, \tau + \bar{\tau}, \tau, -\frac{c}{a}\right) \\ &= \text{Nm}_{K/\mathbb{Q}}(\alpha) \left(1, -\frac{b}{a}, -\frac{c}{a}, \tau\right) \\ &= \frac{\text{Nm}_{K/\mathbb{Q}}(\alpha)}{a} (1, a\tau) \end{aligned}$$

as needed (using $\gcd(a, b, c) = 1$ in the last step). \square

Let $P(\mathcal{O})$ be the subgroup of principal ideals in $I(\mathcal{O})$. Define the **class group** of \mathcal{O} to be

$$C(\mathcal{O}) = I(\mathcal{O})/P(\mathcal{O}).$$

Let $P^+(\mathcal{O})$ be the subgroup of principal ideals in the form (α) where α is *totally positive*, i.e. positive under every real embedding. (This is an empty condition if \mathcal{O} is imaginary.) Define the **narrow class group** of \mathcal{O} to be

$$C^+(\mathcal{O}) = I(\mathcal{O})/P^+(\mathcal{O}).$$

(This is an example of what is called a ray class group in class field theory.)

5 Gauss composition

Theorem 4.5.1 (Correspondence between ideals and binary quadratic forms). *There is a bijection between*

1. narrow ideal classes in quadratic rings with given orientation and
2. binary quadratic forms (up to proper equivalence),

given by

$$\begin{aligned} (\mathfrak{a} = (\alpha, \beta), R) &\mapsto \frac{\text{Nm}_{K/\mathbb{Q}}(\alpha x - \beta y)}{\mathfrak{N}_R(\mathfrak{a})} \\ \left(\left(1, \frac{-b + \sqrt{d}}{2a}\right), \mathbb{Z} \left[\frac{-b + \sqrt{d}}{2} \right] \right) &\leftrightarrow Q(x, y) = ax^2 + bxy + cy^2 \end{aligned}$$

where K is the quadratic field containing \mathfrak{a} , (α, β) is a positively oriented basis for \mathfrak{a} , and $d = b^2 - 4ac$. This restricts to a bijection between invertible oriented ideal classes in the order of discriminant d and primitive binary quadratic forms of discriminant d :

$$C^+(\mathcal{O}(d)) \xrightarrow{\cong} C(d).$$

Corollary 4.5.2 (Gauss composition). *There exists a group structure on equivalence classes of binary quadratic forms, induced by the group structure on ideal classes.*

Proof. Step 1: We show the forward map is well-defined. We need to check two things.

1. Change of basis gives an equivalent form: Temporarily write $Q_{a_1, a_2}(x, y) = \frac{\text{Nm}_{K/\mathbb{Q}}(a_1x - a_2y)}{\mathfrak{N}\mathfrak{a}}$. Suppose $\mathfrak{a} = (a_1, a_2) = (b_1, b_2)$ where both bases are positively oriented. We can write

$$\begin{pmatrix} b_1 \\ -b_2 \end{pmatrix} = A \begin{pmatrix} a_1 \\ -a_2 \end{pmatrix}, \quad A \in \text{SL}_2(\mathbb{Z}).$$

Then

$$Q_{b_1, b_2}(x, y) = \frac{\text{Nm}_{K/\mathbb{Q}}\left((x, y) \begin{pmatrix} b_1 \\ -b_2 \end{pmatrix}\right)}{\mathfrak{N}_R(\mathfrak{a})} = \frac{\text{Nm}_{K/\mathbb{Q}}\left((x, y)A \begin{pmatrix} a_1 \\ -a_2 \end{pmatrix}\right)}{\mathfrak{N}_R(\mathfrak{a})} = Q_{a_1, a_2}((x, y)A) \quad (4.4)$$

so the quadratic forms are equivalent.

2. Multiplying by a totally positive element gives an equivalent form: Suppose λ is totally positive. Then $\text{Nm}_{K/\mathbb{Q}}(\lambda) > 0$. First note that $(\lambda a_1, \lambda a_2)$ is also positively oriented:

$$\frac{\begin{vmatrix} \lambda a_1 & \overline{\lambda a_1} \\ \lambda b_1 & \overline{\lambda b_1} \end{vmatrix}}{d} = \text{Nm}_{K/\mathbb{Q}}(\lambda) \frac{\begin{vmatrix} a_1 & \overline{a_1} \\ b_1 & \overline{b_1} \end{vmatrix}}{d} > 0.$$

Then

$$\begin{aligned} Q_{\lambda a_1, \lambda a_2}(x, y) &= \frac{\text{Nm}(\lambda a_1 x - \lambda a_2 y)}{\mathfrak{N}_R(\lambda \mathfrak{a})} \\ &= \frac{\text{Nm}_{K/\mathbb{Q}}(a_1 x - a_2 y)}{\mathfrak{N}_R(\mathfrak{a})} \\ &= Q_{a_1, a_2}(x, y) \end{aligned}$$

as needed.

Step 2: We show this map is injective. First note an alternate characterization for the forward map. Writing $(\alpha, \beta) = \alpha(1, \tau)$, we find that the quadratic form corresponding to (α, β) is

$$\begin{aligned} Q_{\alpha, \beta}(x, y) &= \frac{\text{Nm}_{K/\mathbb{Q}}(\alpha x - \beta y)}{\mathfrak{N}_R(\mathfrak{a})} \\ &= \frac{(\alpha x - \beta y)(\overline{\alpha} x - \overline{\beta} y)}{\mathfrak{N}_R(\mathfrak{a})} \\ &= \frac{\alpha \overline{\alpha} x^2 - (\alpha \overline{\beta} + \overline{\alpha} \beta) xy + \beta \overline{\beta} y^2}{\mathfrak{N}_R(\mathfrak{a})} \\ &= \frac{\text{Nm}_{K/\mathbb{Q}}(\alpha)}{\mathfrak{N}_R(\mathfrak{a})} (x - \tau y)(x - \overline{\tau} y), \quad \tau = \frac{\beta}{\alpha}. \end{aligned} \quad (4.5)$$

Suppose $Q_{a_1, a_2}(x, y) \sim Q_{b_1, b_2}(x, y)$. By changing the basis of $\mathfrak{b} = (b_1, b_2)$, which by (4.4) corresponds to changing the basis of the quadratic form, we may assume $Q_{a_1, a_2}(x, y) = Q_{b_1, b_2}(x, y)$. The above factorization (4.5) says that one of the following holds:

1. $\frac{a_1}{a_2} = \frac{b_1}{b_2}$. Letting $\lambda = \frac{a_1}{b_1} = \frac{a_2}{b_2}$, we find $\mathfrak{a} = \lambda\mathfrak{b}$. Since both bases are positively oriented,

$$0 < \frac{\begin{vmatrix} a_1 & \overline{a_1} \\ a_2 & \overline{a_2} \end{vmatrix}}{\begin{vmatrix} b_1 & \overline{b_1} \\ b_2 & \overline{b_2} \end{vmatrix}} = \text{Nm}_{K/\mathbb{Q}}(\lambda),$$

showing either λ or $-\lambda$ is totally positive.

2. $\frac{a_1}{a_2} = \frac{\overline{b_1}}{\overline{b_2}}$. We show that this kind of “disorientation” is impossible. Let $\lambda = \frac{a_1}{b_1} = \frac{a_2}{b_2}$. Then

$$0 < \frac{\begin{vmatrix} a_1 & \overline{a_1} \\ a_2 & \overline{a_2} \end{vmatrix}}{\begin{vmatrix} b_1 & \overline{b_1} \\ b_2 & \overline{b_2} \end{vmatrix}} = -\frac{\begin{vmatrix} a_1 & \overline{a_1} \\ a_2 & \overline{a_2} \end{vmatrix}}{\begin{vmatrix} \overline{b_1} & b_1 \\ \overline{b_2} & b_2 \end{vmatrix}} = -\text{Nm}_{K/\mathbb{Q}}(\lambda),$$

giving $\text{Nm}_{K/\mathbb{Q}}(\lambda) < 0$. But

$$\begin{aligned} Q_{b_1, b_2}(x, y) &= \frac{(b_1x - b_2y)(\overline{b_1}x - \overline{b_2}y)}{\mathfrak{N}_R(\mathfrak{a})} \\ Q_{a_1, a_2}(x, y) &= \frac{(a_1x - a_2y)(\overline{a_1}x - \overline{a_2}y)}{\mathfrak{N}_R(\mathfrak{b})} = \lambda\overline{\lambda} \frac{(\overline{b_1}x - \overline{b_2}y)(b_1x - b_2y)}{\mathfrak{N}_R(\mathfrak{b})}; \end{aligned}$$

equating gives $\text{Nm}_{K/\mathbb{Q}}(\lambda) > 0$, contradiction.

Step 3: Applying the reverse map and then the forward map gives the identity.

Given $Q(x, y) = ax^2 + bxy + cy^2 = a(x - \tau y)(x - \overline{\tau}y)$, the reverse map takes it to $\mathfrak{a} := (1, \tau)$. Note $\{1, \tau := \frac{-b+\sqrt{d}}{2a}\}$ is in fact a \mathbb{Z} -basis for $(1, \tau)$ over $R := \mathbb{Z}[a\tau] = \mathbb{Z}\left[\frac{-b+\sqrt{d}}{2}\right]$ (not just a generating set over \mathcal{O}). Indeed, $a\tau(\tau) = (-b\tau - c) \in (1, \tau)$. In exactly the same way, $\{1, a\tau\}$ is a \mathbb{Z} -basis for R over R .

By (4.5), the forward map then takes (\mathfrak{a}, R) to

$$\frac{1}{\mathfrak{N}_R(\mathfrak{a})}(x - \tau y)(x - \overline{\tau}y) = [\mathfrak{a} : R](x - \tau y)(x - \overline{\tau}y) = a(x - \tau y)(x - \overline{\tau}y).$$

Step 4: Invertible classes correspond to primitive forms. Suppose $\mathfrak{a} = \alpha(1, \tau)$ is invertible and τ satisfies $ax^2 + bx + c = 0$, where $\gcd(a, b, c) = 1$. Then by Lemma 4.4.6, $a = \frac{\text{Nm}_{K/\mathbb{Q}}(\alpha)}{\mathfrak{N}_R(\mathfrak{a})}$. Hence by (4.5), the quadratic form is $ax^2 + bxy + cy^2$, which is primitive.

Conversely suppose Q is primitive. Then by Proposition 4.4.6, the corresponding ideal $(1, \tau)$ is proper in $R := (1, a\tau)$.

The fact that the discriminant is preserved can be seen from the reverse map. \square

Example 4.5.3: We calculate the binary quadratic form corresponding to the order \mathcal{O} of discriminant d . This will be the identity element in the form class group $C(D)$. We have $\mathcal{O} = (1, \tau)$ where

$$\tau = \begin{cases} \frac{1+\sqrt{d}}{2}, & d \equiv 1 \pmod{4} \\ \frac{\sqrt{d}}{2}, & d \equiv 0 \pmod{4}. \end{cases}$$

Then

$$Q_{\mathcal{O}}(x, y) = \text{Nm}_{K/\mathbb{Q}}(x + y\tau) = \begin{cases} x^2 - \frac{d}{4}y^2, & d \equiv 0 \pmod{4} \\ x^2 + xy - \frac{d-1}{4}y^2, & d \equiv 1 \pmod{4}. \end{cases}$$

This is consistent with the fact that $x^2 - \frac{d}{4}$ and $x^2 + x - \frac{d-1}{4}$ are the minimal polynomials of τ in the two cases, respectively.

Theorem 4.5.4. *Let \mathfrak{a} be an invertible ideal in the quadratic ring \mathcal{O} and f its associated quadratic binary form. Let m be a nonzero integer. Then the following are equivalent.*

1. *There exists \mathfrak{a}' in the same ideal class as \mathfrak{a} with*

$$\mathfrak{a}'\overline{\mathfrak{a}'} = (m).$$

2. *There exists \mathfrak{a}' in the same ideal class as \mathfrak{a} with $\mathfrak{N}_{\mathcal{O}}(\mathfrak{a}') = m$.*

3. *f represents m .*

Proof. Equivalence of the first two items is clear. We show (2) \iff (3).

Suppose f represents m . Suppose $m = d^2a$, and f represents a primitively. By Proposition 4.2.2, f is equivalent to a form $ax^2 + bxy + cy^2$. By Gauss composition, this form corresponds to an ideal $\mathfrak{a}' = a(1, \tau)$ with $a\tau^2 + b\tau + c = 0$ inside $\mathcal{O} = (1, a\tau)$. Hence $\mathfrak{N}_{\mathcal{O}}(\mathfrak{a}') = a$. Then

$$\mathfrak{N}_{\mathcal{O}}(d\mathfrak{a}') = d^2a,$$

as needed.

Conversely, suppose $\mathfrak{N}_{\mathcal{O}}(\mathfrak{a}) = m$. Write $\mathfrak{a} = \alpha(1, \tau)$ with $\text{Nm}_{K/\mathbb{Q}}(\alpha) > 0$. Suppose $a\tau^2 + b\tau + c = 0$ with $\gcd(a, b, c) = 1$, so $\mathcal{O} = (1, a\tau)$ and $\mathfrak{N}_{\mathcal{O}}((1, \tau)) = \frac{1}{a}$. The corresponding quadratic form is

$$g(x, y) = \frac{\text{Nm}_{K/\mathbb{Q}}(x - \tau y)}{\mathfrak{N}_{\mathcal{O}}((1, \tau))} = a \text{Nm}_{K/\mathbb{Q}}(x - \tau y).$$

Since $\alpha \in \mathcal{O} = (1, a\tau)$, we have $\alpha = p - qa\tau$ for some $p, q \in \mathbb{Z}$. We have $\alpha\tau = p\tau - q(-b\tau - c) =$

$(p + qb)\tau + cq$; since $\alpha\tau \in \mathcal{O} = (1, a\tau)$ as well, we get $\frac{p+qb}{a} \in \mathbb{Z}$. Now by Lemma 4.4.6,

$$\begin{aligned} m = \mathfrak{N}_{\mathcal{O}}(\mathfrak{a}) &= \frac{\text{Nm}_{K/\mathbb{Q}}(\alpha)}{a} \\ &= \frac{1}{a^2} \cdot a \text{Nm}_{K/\mathbb{Q}}(p - qa\tau) \\ &= \frac{1}{a^2} g(p, aq) \\ &= g\left(\frac{p}{a}, q\right) \\ &= g\left(\frac{-bq - p}{a}, q\right) \end{aligned} \quad g(x, y) = g\left(-\frac{b}{a}y - x, y\right).$$

We showed above that $\frac{-bq-p}{a} \in \mathbb{Z}$, as needed. (Think of the last step as “root flipping.”) \square

6 Ideal class group of an order

Suppose \mathcal{O} is an order in the field K , and \mathcal{O}_K is the ring of integers (the maximal order). We want to relate $C(\mathcal{O})$ to $C(\mathcal{O}_K)$, because the latter is the most “natural” class group for K . In reality, we will relate $C(\mathcal{O})$ to a quotient of a subgroup of $I(\mathcal{O}_K)$, a generalized ideal class group of \mathcal{O}_K .

After learning class field theory, which relates generalized class ideal class groups to extensions of K , we will see that the primes represented by the quadratic form corresponding to \mathcal{O} can be characterized in terms of a certain field extensions L/K .

Definition 4.6.1: Define

$$\begin{aligned} I_K(f) &= \{\mathfrak{a} \in I_K : \mathfrak{a} \text{ relatively prime to } f\mathcal{O}_K\} \\ P_K(\mathbb{Z}, f) &= \{\alpha\mathcal{O}_K : \alpha \equiv a \pmod{f\mathcal{O}_K} \text{ for some } a \in \mathbb{Z}\} \\ I_K(\mathcal{O}, f) &= \{\mathfrak{a} \in I(\mathcal{O}) : \mathfrak{a} \text{ relatively prime to } f\mathcal{O}\}. \end{aligned}$$

Theorem 4.6.2. *Let f be the conductor of \mathcal{O} , i.e. $\mathcal{O} = \mathbb{Z} + f\mathcal{O}_K$. There is an isomorphism*

$$I_K(f)/P_K(\mathbb{Z}, f) \rightarrow I(\mathcal{O})/P(\mathcal{O}) = C(\mathcal{O})$$

induced by the map $g : I_K(f) \rightarrow I(\mathcal{O})$,

$$g(\mathfrak{a}) = \mathfrak{a} \cap \mathcal{O}.$$

First, a preliminary lemma.

Lemma 4.6.3. *Let \mathcal{O} be an order of conductor f . Then every \mathcal{O} -ideal prime to f is proper.*

Proof. Cox, Prop. 7.20. Suppose \mathfrak{a} is prime to f . Then $\mathfrak{a} + f\mathcal{O} = \mathcal{O}$. Suppose $\beta\mathfrak{a} \subseteq \mathfrak{a}$. Then

$$\beta\mathcal{O} = \beta(\mathfrak{a} + f\mathcal{O}) = \beta\mathfrak{a} + \beta f\mathcal{O} \subseteq \mathfrak{a} + f\mathcal{O}_K \subseteq \mathcal{O}$$

so $\beta \in \mathcal{O}$. Thus \mathfrak{a} is proper. \square

Proof of Theorem 4.6.2. Step 1: We show there is a norm-preserving isomorphism

$$\begin{aligned} I_K(f) &\rightarrow I(\mathcal{O}, f) \\ \mathfrak{a} &\mapsto \mathfrak{a} \cap \mathcal{O} \\ \mathfrak{b}\mathcal{O}_K &\leftarrow \mathfrak{b}. \end{aligned}$$

Step 2: The map above induces an isomorphism $I_K(f)/P_K(\mathbb{Z}, f) \rightarrow I(\mathcal{O}, f)/P(\mathcal{O}, f)$

Step 3: The inclusion $I(\mathcal{O}, f) \hookrightarrow I(\mathcal{O})$ induces an isomorphism $I(\mathcal{O}, f)/P(\mathcal{O}, f) \rightarrow I(\mathcal{O})/P(\mathcal{O})$. This follows from Theorem 10.1.1. \square

7 Cube law

We now derive quadratic composition in a different way. We will associate a “cube” of integers with three quadratic forms. In order to identify equivalent binary quadratic forms, we mod out by $\mathrm{SL}_2(\mathbb{Z})^3$. After decreeing that the sum of forms making up any cube is 0, we find that we have


1. identified quadratic forms up to equivalence, and
2. recovered our original composition law.

Later we will see that these ideas generalize to composition laws for other polynomial forms and associated ideals/rings.

Let $\mathcal{C}_2 = \mathbb{Z}^2 \otimes \mathbb{Z}^2 \otimes \mathbb{Z}^2$. Choosing a basis (v_1, v_2) for \mathbb{Z}^2 , every element of \mathcal{C}^2 can be written in the form

$$\begin{aligned} &av_1 \otimes v_1 \otimes v_1 + bv_1 \otimes v_2 \otimes v_1 + cv_2 \otimes v_1 \otimes v_1 + dv_2 \otimes v_2 \otimes v_1 \\ &+ ev_1 \otimes v_1 \otimes v_2 + fv_1 \otimes v_2 \otimes v_2 + gv_2 \otimes v_1 \otimes v_2 + hv_2 \otimes v_2 \otimes v_2. \end{aligned}$$

We represent this graphically as a **cube**.



Think of this as a higher-dimensional analogue of a matrix. Let M_i, N_i for $i = 1, 2, 3$ be the two matrices obtained by slicing the cube along the 3 possible directions.

$$\begin{aligned} M_1 &= \begin{pmatrix} a & b \\ c & d \end{pmatrix}, & N_1 &= \begin{pmatrix} e & f \\ g & h \end{pmatrix} \\ M_2 &= \begin{pmatrix} a & c \\ e & g \end{pmatrix}, & N_2 &= \begin{pmatrix} b & d \\ f & h \end{pmatrix} \\ M_3 &= \begin{pmatrix} a & e \\ b & f \end{pmatrix}, & N_3 &= \begin{pmatrix} c & g \\ d & h \end{pmatrix}. \end{aligned}$$

Define an action of $\Gamma = \mathrm{SL}_2(\mathbb{Z}) \times \mathrm{SL}_2(\mathbb{Z}) \times \mathrm{SL}_2(\mathbb{Z})$ on \mathcal{C}_2 by letting $\begin{pmatrix} r & s \\ t & u \end{pmatrix}$ in the i th factor of $\mathrm{SL}_2(\mathbb{Z})^3$ act on A by sending

$$\begin{pmatrix} M_i \\ N_i \end{pmatrix} \mapsto \begin{pmatrix} r & s \\ t & u \end{pmatrix} \begin{pmatrix} M_i \\ N_i \end{pmatrix} = \begin{pmatrix} rM_i + sN_i \\ tM_i + uN_i \end{pmatrix}.$$

Note that the actions of the 3 factors of $\mathrm{SL}_2(\mathbb{Z})$ commute, in the same way that row and column operations commute for a matrix.

Now associate a cube A with three binary quadratic forms Q_1^A, Q_2^A, Q_3^A by letting

$$Q_i^A(x, y) = -\det(M_i x - N_i y).$$

We call A **projective** if Q_1^A, Q_2^A, Q_3^A are all primitive.

Invariant theory gives the following result.

Proposition 4.7.1: The ring of invariants of \mathcal{C}_2 under $\mathrm{SL}_2(\mathbb{Z})^3$ is

$$(\mathcal{C}_2)^{\mathrm{SL}_2(\mathbb{Z})^3} = \mathbb{Z}[\mathrm{disc}(A)]$$

where

$$\begin{aligned} \mathrm{disc}(A) &:= \mathrm{disc}(Q_1) = \mathrm{disc}(Q_2) = \mathrm{disc}(Q_3) \\ &= \sum_{s,t \text{ long diagonal}} s^2 t^2 - 2 \sum_{s,t,u,v \text{ face}} stuv + 4 \sum_{s,t,u,v \text{ regular tetrahedron}} stuv. \end{aligned}$$

(The fact that $\mathrm{disc}(A)$ is invariant is easy to see; we shall not need the opposite implication.)

We now prove the bijection in Theorem 4.5.1 and Gauss composition (Corollary 4.5.2) in a different way, using cubes. The idea is to associate triples of ideals multiplying to 1 with triples of quadratic forms in the same cube (which we will deem to add up to 0), and in this way transfer the group structure from narrow ideal classes to classes of quadratic forms.

Definition 4.7.2: We say that three oriented fractional ideals I_1, I_2, I_3 in a quadratic ring S form a **balanced triple** if

$$\begin{aligned} I_1 I_2 I_3 &\subseteq S \text{ and} \\ \mathbb{N}(I_1) \mathbb{N}(I_2) \mathbb{N}(I_3) &= 1. \end{aligned}$$

We say two balanced triples (I_1, I_2, I_3) and (I'_1, I'_2, I'_3) are equivalent if there are $\lambda_1, \lambda_2, \lambda_3$ such that

$$\begin{aligned} I_1 &= \lambda_1 I'_1 \\ I_2 &= \lambda_2 I'_2 \\ I_3 &= \lambda_3 I'_3. \end{aligned}$$

Theorem 4.7.3. *There is a bijection between equivalence classes of cubes, and ordered pairs $(S, (I_1, I_2, I_3))$ where S is a quadratic ring and (I_1, I_2, I_3) is a balanced triple modulo equivalence.*

$$\mathbb{Z}^2 \otimes \mathbb{Z}^2 \otimes \mathbb{Z}^2 / \mathrm{SL}_2(\mathbb{Z})^3 \leftrightarrow \{(S, (I_1, I_2, I_3))\}$$

If (α_1, α_2) , (β_1, β_2) and (γ_1, γ_2) are correctly oriented bases for I_1 , I_2 , and I_3 , then the cube is given by $(a_{ijk})_{1 \leq i, j, k \leq 2}$ where

$$\alpha_i \beta_j \gamma_k = c_{ijk} + a_{ijk} \tau$$

and τ is such that

$$\begin{aligned} \tau^2 - \frac{d}{4} &= 0, & d &\equiv 0 \pmod{4} \\ \tau^2 - \tau - \frac{d-1}{4} &= 0, & d &\equiv 1 \pmod{4}. \end{aligned}$$

Chapter 5

Units in number fields

1 Units

Any finitely generated abelian group is isomorphic to $A_{\text{tors}} \oplus \mathbb{Z}^t$ where A_{tors} consists of all torsion elements, i.e. elements of finite order. The number t is called the **rank** of A .

The main theorem of this chapter is the following.

Theorem 5.1.1 (Dirichlet’s unit theorem). *Let K be a number field with r real embeddings and $2s$ nonreal complex embeddings. Then the group of units in K is finitely generated with rank equal to $r + s - 1$.*

The idea of the proof is as follows.

1. Following the idea of the proof that the class number is finite (Section 3), we embed the set of units as a lattice in $\mathbb{R}^r \times \mathbb{R}^s$. Since we want to send a group (under multiplication) to a lattice (under addition), we take logarithms of the norm to define our embedding. In actuality, the homomorphism L is not injective, but the kernel will be finite, which is good enough. (See Proposition 5.2.2.)
2. Construct independent units from elements generating the same ideal. We do this by finding α, γ generating the same principal ideal and taking $\alpha\gamma^{-1}$. Consider a fixed large symmetric convex compact set T of $\mathbb{R}^r \times \mathbb{C}^s$, which will contain elements $\sigma(\alpha)$ by Minkowski. For α such that $L(\alpha) \in T$, (α) is one of a finite number of principal ideals (γ_k) . Then $\alpha\gamma_k^{-1}$ is a unit.

However, since we want independent units, we look not for points in the form $L(\alpha)$ but rather of the form $\mathbf{x}L(\alpha)$ where x has norm 1. Think of this as “rotating” or “twisting” the unit that we get.

First, a basic criterion for being a unit.

Proposition 5.1.2: Let K/\mathbb{Q} be a finite extension. An element $\alpha \in K$ is a unit if and only if $\text{Nm}(\alpha) = \pm 1$.

Proof. Suppose α is a unit. Then $\alpha^{-1} \in K$ and

$$\text{Nm}(\alpha) \text{Nm}(\alpha^{-1}) = \text{Nm}(\alpha\alpha^{-1}) = 1$$

so $\text{Nm}(\alpha) = \pm 1$.

Conversely, suppose $\text{Nm}(\alpha) = \pm 1$. Then by Theorem 1.2.3, letting $\sigma_1 = I, \dots, \sigma_n$ be the distinct embeddings of K to the Galois closure, we have

$$\alpha \cdot \prod_{k=2}^n \sigma_k(\alpha) = \text{Nm}_{L/K}(\alpha) = \pm 1.$$

Hence $\alpha^{-1} = \pm \prod_{k=2}^n \sigma_k(\alpha) \in \mathcal{O}_K$. □

2 Dirichlet's unit theorem

We now prove Dirichlet's unit theorem.

Lemma 5.2.1. *There are a finite number of algebraic integers α such that*

$$\begin{aligned} [\mathbb{Q}(\alpha) : \mathbb{Q}] &\leq m \\ |\alpha'| &\leq M \text{ for all conjugates } \alpha'. \end{aligned}$$

Proof. The second condition means that the coefficients of the minimal polynomial f are bounded. Since the degree of f is at most m , there are a finite number of possibilities for the f and hence α .¹ □

Let $\{\sigma_1, \dots, \sigma_r\}$ be the real embeddings and $\{\sigma_{r+1}, \bar{\sigma}_{r+1}, \dots, \sigma_{r+s}, \bar{\sigma}_{r+s}\}$ be the complex embeddings of K . Since

$$\text{Nm}(\alpha) = |\sigma_1(\alpha)| \cdots |\sigma_r(\alpha)| |\sigma_{r+1}(\alpha)|^2 \cdots |\sigma_{r+s}(\alpha)|^2,$$

we define the homomorphism

$$\begin{aligned} L : K^\times &\rightarrow \mathbb{R}^{r+s} \\ L(\alpha) &= (\ln |\sigma_1(\alpha)|, \dots, \ln |\sigma_r(\alpha)|, 2 \ln |\sigma_{r+1}(\alpha)| \cdots, 2 \ln |\sigma_{r+s}(\alpha)|). \end{aligned}$$

This is the composition of our previous embedding σ with f :

$$\begin{aligned} \sigma : K &\rightarrow \mathbb{R}^r \times \mathbb{C}^s & \sigma(\alpha) &= (\sigma_1(\alpha_1), \dots, \sigma_r(\alpha_r)) \\ f : \mathbb{R}^r \times \mathbb{R}^s &\rightarrow \mathbb{R}^{r+s} & f(x_1, \dots, x_r, z_{r+1}, \dots, z_{r+s}) &= (\ln |x_1|, \dots, \ln |x_r|, 2 \ln |z_{r+1}|, \dots, 2 \ln |z_{r+s}|). \end{aligned}$$

Proposition 5.2.2: The image $L(U_K)$ is a lattice contained in the hyperplane

$$H := \{(x_1, \dots, x_{r+s}) : x_1 + \cdots + x_{r+s} = 0\}.$$

Moreover, L has finite kernel.

¹See Chapter ?? for...

Proof. If $\alpha \in U_K$, $L(\alpha) = (x_1, \dots, x_{r+s})$, then

$$\begin{aligned} x_1 + \dots + x_{r+s} &= \ln |\sigma_1(\alpha)| + \dots + \ln |\sigma_r(\alpha)| + 2 \ln |\sigma_{r+1}(\alpha)| + \dots + 2 \ln |\sigma_{r+s}(\alpha)| \\ &= \ln |\text{Nm}(\alpha)| = 0. \end{aligned}$$

To show $L(U_K)$ is a lattice it suffices to show it is discrete. To this end, we show the base elements

$$B(r) = \{(x_1, \dots, x_{r+s}) : |x_j| \leq C\}$$

centered at the origin contain finitely many points of $L(U_K)$. Indeed, if $\sigma(\alpha) \in B(r)$, then $|\sigma_k(\alpha)| < C$ for every embedding σ_k . By Proposition 5.2.1, there are a finite number of possibilities for α .

If $\alpha \in \ker L$, then $|\sigma_k(\alpha)| = 1$ for all k . Again by Proposition 5.2.1 there are a finite number of possibilities for α . \square

Since U_K is abelian, we now know

$$U_K \cong \underbrace{\ker(L)}_{U_K^{\text{tors}}} \oplus \underbrace{L(U_K)}_{\text{lattice of } H}.$$

It remains to show the following.

Lemma 5.2.3. *$L(U_K)$ is a full lattice in H . Therefore it has rank $r + s - 1$.*

Proof. Let $\mathbf{x} \in \mathbb{R}^r \times \mathbb{C}^s$. By Proposition 3.3.1, the volume of the fundamental parallelepiped of $\sigma(\mathbf{a})$ is $2^{-s} \cdot \mathbb{N}\mathbf{a} \cdot |\Delta_K|^{\frac{1}{2}}$. Note that multiplication by \mathbf{x} multiplies the norm by $\text{Nm}(\mathbf{x})$ so the volume of the fundamental parallelepiped of $\sigma(\mathbf{a})$ is $\text{Nm}(\mathbf{x})2^{-s}\mathbb{N}\mathbf{a} \cdot |\Delta_K|^{\frac{1}{2}}$.

Now suppose \mathbf{x} is any element such that $\text{Nm}(\mathbf{x}) = 1$. Let $V = 2^{-s}\mathbb{N}\mathbf{a} \cdot |\Delta_K|^{\frac{1}{2}}$. Let T be any compact convex symmetric set with volume at least $2^{r+s}V$. We note the following.

1. By Minkowski's Theorem, there is point of T in the lattice $\mathbf{x} \cdot \sigma(\mathcal{O}_K)$.
2. Since T is bounded, all elements of T have norm bounded by a constant C . If $\sigma(\alpha) \in T$, then α has norm bounded by C . By Lemma 3.3.7 there are a finite number of principal ideals with norm bounded by C , say $(\gamma_1), \dots, (\gamma_m)$. Then if $\sigma(\alpha) \in T$, we have $(\alpha) = (\gamma_k)$, i.e. $\alpha = u\gamma_k$ for some unit u , and some k .

In conclusion, for each \mathbf{x} we find α such that

$$T \ni \mathbf{x}\sigma(\alpha) = \mathbf{x}\sigma(u\gamma_k) \text{ for some } k,$$

i.e.

$$\mathbf{x}\sigma(u) \in \bigcup_{k=1}^m \sigma(\gamma_k^{-1})T. \quad (5.1)$$

Since T is bounded, so is $\bigcup_{k=1}^m \sigma(\gamma_k^{-1})T$. There exists C' so that every coordinate of $\mathbf{x}\sigma(u)$ is less than C' :

$$(\mathbf{x}\sigma(u))_k < C'. \quad (5.2)$$

The idea is that this places a large constraint on the possibilities for ε , so as we vary \mathbf{x} between “extreme” values, we will have to get linearly independent u .

Take

$$\mathbf{x}_k = \left(C', \dots, C', \underbrace{\frac{1}{C'^{r+s-1}}}_k, C', \dots, C' \right)$$

Then letting u_k be such that (5.1) holds for x_k, u_k , we get by (5.2) that, componentwise,

$$\sigma(u_k) < (1, \dots, 1, C'^{r+s}, 1, \dots, 1),$$

i.e.

$$L(u_k) = f(\sigma(u_k)) < (0, \dots, 0, \ln(C'^{r+s}), 0, \dots, 0).$$

Note the following.

1. Every entry of $L(u_k)$ is negative except for the k th one, which must be positive because the entries sum up to 0.
2. The sum of entries of $L(u_k)$, omitting the last term, is positive.

The following lemma will show that $L(u_1), \dots, L(u_{r+s-1})$ are linearly independent. It will follow that u_1, \dots, u_{r+s-1} generate a free abelian group. This means $\text{rank}(U_K) \geq r + s - 1$; we have equality by Proposition 5.2.2 since $\dim H = r + s - 1$.

Lemma 5.2.4. *Suppose that A is a $n \times n$ matrix such that*

1. $a_{i,j} < 0$ for $i \neq j$ and $a_{i,i} > 0$.
2. $\sum_{j=1}^n a_{i,j} > 0$.

Then A is invertible.

Proof. Suppose $v = \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix}$ is a nonzero vector. Suppose i is such that $|a_i|$ is greatest. Then looking at the i th component gives $\sum_{j=1}^n a_{ij}v_j = 0$. Then

$$\sum_{j=1}^n a_{ij}v_j > a_{ij}v_i + \sum_{j \neq i} a_{ij}v_i > 0,$$

so $Av \neq 0$. Thus A is invertible. □

□

This finishes the proof of Dirichlet’s Unit Theorem.

3 S -units

Definition 5.3.1: Let S be a finite set of prime ideals of K . The **ring of S -integers** is

$$\mathcal{O}_K(S) = \bigcap_{\mathfrak{p} \notin S} (\mathcal{O}_K)_{\mathfrak{p}} = \{\alpha \in K : \text{ord}_{\mathfrak{p}}(\alpha) \geq 0 \text{ for all } \mathfrak{p} \notin S\}.$$

I.e. we allow dividing by elements whose “only prime factors” are in S . The group of S -units is the group of units in $\mathcal{O}_K(S)$:

$$U(S) = \mathcal{O}_K(S)^{\times} = \{\alpha \in K \mid \text{ord}_{\mathfrak{p}}(\alpha) = 0 \text{ for all } \mathfrak{p} \notin S\}.$$

There are more units in $U(S)$ than in U_K ; the following generalization of Dirichlet’s theorem says that we get an “extra” unit for every prime in S .

Theorem 5.3.2 (Dirichlet’s S -unit theorem). *The group of S -units is finitely generated with rank $r + s + |S| - 1$.*

Proof. Let $S = \{\mathfrak{p}_1, \dots, \mathfrak{p}_t\}$. Consider the maps

$$U_K \hookrightarrow U(S) \xrightarrow{\varphi} \mathbb{Z}^m$$

where

$$\varphi(x) = (\text{ord}_{\mathfrak{p}_1}(x), \dots, \text{ord}_{\mathfrak{p}_m}(x)).$$

Its kernel is U_K , as the elements of U_K are exactly those x with order 0 for every prime \mathfrak{p} , and by definition $\text{ord}_{\mathfrak{p}}(x) = 0$ for $x \in U(S)$ and \mathfrak{p} outside of S . Let h be the class number of K . Then $\mathfrak{p}_k^h = (\alpha_k)$ for some α_k . We have

$$\varphi(x) = (0, \dots, 0, \underbrace{h}_k, 0, \dots, 0).$$

Hence $\varphi(U(S))$ is a full lattice in \mathbb{Z}^m . Since U_K has rank $r + s - 1$ by Dirichlet’s Unit Theorem (5.1.1), $U(S)$ has rank $r + s - 1 + m$. \square

4 Examples and algorithms

5 Regulator

Chapter 6

Cyclotomic fields

1 Cyclotomic polynomials

Definition 6.1.1: A **cyclotomic extension** of \mathbb{Q} is a field $\mathbb{Q}[\zeta]$ where ζ is a root of unity. We call ζ a primitive n th root of unity if $\zeta^n = 1$ but $\zeta^m \neq 1$ for $0 < m < n$.

We will use ζ_n to denote a primitive n th root of unity.

The n th cyclotomic polynomial is defined by

$$\Phi_n(x) = \prod_{0 \leq j < n, \gcd(j,n)=1} (x - e^{\frac{2\pi ij}{n}})$$

Equivalently, it can be defined by the recurrence $\Phi_0(x) = 1$ and

$$\Phi_n(x) = \frac{x^n - 1}{\prod_{m|n, m < n} \Phi_m(x)}.$$

Hence, it has integer coefficients.

Theorem 6.1.2. *The cyclotomic polynomials are irreducible over $\mathbb{Q}[x]$.*

Proof. We need the following lemma:

Suppose ω is a primitive n th root of unity, and that its minimal polynomial is $g(x)$. Let p be a prime not dividing n . Then ω^p is a root of $g(x) = 0$.

Since $\Phi_n(\omega) = 0$, we can write $\Phi_n = fg$. If $g(\omega^p) \neq 0$ then $f(\omega^p) = 0$. Since ω is a zero of $f(x^p)$, $f(x^p)$ factors as

$$f(x^p) = g(x)h(x)$$

for some polynomial $h \in \mathbb{Z}[x]$.

Now, in $\mathbb{Z}/p\mathbb{Z}[x]$ note $(f_1 + \dots + f_k)^p = f_1^p + \dots + f_k^p$ since the p th power map is an homomorphism. Hence

$$g(x)h(x) \equiv f(x^p) \equiv f(x)^p \pmod{p}.$$

Hence $f(x)$ and $g(x)$ share a factor modulo p . However, the derivative of $x^n - 1$ modulo p is $nx^{n-1} \not\equiv 0$, showing that $x^n - 1$ has no repeated irreducible factor modulo p ; hence Φ_n has no repeated factor modulo p . Since $\Phi_n = fg$, this produces a contradiction.

Therefore $g(\omega^p) = 0$, as needed.

Any primitive n th root is in the form ω^k for k relatively prime to n . Writing the prime factorization of k as $p_1 \cdots p_m$, we get by the lemma that $\omega^{p_1}, \omega^{p_1 p_2}, \dots, \omega^{p_1 \cdots p_m}$ are all roots of g . Hence g contains all primitive n th roots of unity as roots, and $\Phi_n = g$ is irreducible. \square

Theorem 6.1.3.

$$[\mathbb{Q}(\zeta_n) : \mathbb{Q}] = \varphi(n).$$

Proof. The minimal polynomial of ζ_n equals the cyclotomic polynomial by Theorem 6.1.2; the latter has degree $\varphi(n)$. \square

We use cyclotomic polynomials to prove a special case of Dirichlet's theorem.

Theorem 6.1.4 (Dirichlet's theorem for $p \equiv 1 \pmod{n}$). (†) *Let n be a positive integer. There are infinitely many primes p with $p \equiv 1 \pmod{n}$.*

Lemma 6.1.5. *For any integer m , all divisors of $\Phi_n(m)$ either divide n or are $1 \pmod{n}$.*

Proof. Suppose p is prime and $p \mid \Phi_n(m)$. Then $p \mid m^n - 1$, i.e.

$$m^n \equiv 1 \pmod{p}$$

so $r := \text{ord}_p(m) \mid n$. Since $m^{p-1} \equiv 1 \pmod{p}$ by Fermat's little theorem, $r \mid p - 1$.

If $r = n$, then $n \mid p - 1$, i.e. $p \equiv 1 \pmod{n}$. Suppose that $r < n$. Then

$$p \mid \Phi_n(m) \mid \frac{m^n - 1}{m^r - 1} = m^{r(\frac{n}{r}-1)} + \cdots + m^r + 1.$$

However, $m^r \equiv 1 \pmod{p}$ so

$$m^{r(\frac{n}{r}-1)} + \cdots + m^r + 1 \equiv \frac{n}{r} \pmod{p},$$

so $p \mid \frac{n}{r} \mid n$. \square

Proof of Theorem 6.1.4. Suppose by way of contradiction that only finitely many primes are $1 \pmod{n}$. Let their product be P (if there are no such primes, $P = 1$). Consider $\Phi_n(knP)$, $k \in \mathbb{Z}$. Since it divides $(nP)^n - 1$, it can't have prime divisors in common with n or P . With appropriate choice of k we can be sure $\Phi_n(knP) \neq 0, \pm 1$. By the claim all prime divisors of $\Phi_n(knP)$ are $1 \pmod{n}$, but they don't divide P , contradiction. \square

2 Ring of integers

Our next two propositions will give us information about the ring of integers of $\mathbb{Q}[\zeta]$, as well as some other useful facts. In the process we will rederive Theorem 6.1.3.

Proposition 6.2.1: Suppose ζ and ζ' are primitive n th roots of unity. Then $\frac{1-\zeta'}{1-\zeta}$ is a unit in $\mathbb{Z}[\zeta] = \mathbb{Z}[\zeta']$.

Proof. Then we have $\zeta' = \zeta^s$ and $\zeta = \zeta'^t$ for some s, t , so $\mathbb{Z}[\zeta] = \mathbb{Z}[\zeta']$ and

$$\begin{aligned} \frac{1-\zeta'}{1-\zeta} &= 1 + \zeta + \cdots + \zeta^{s-1} \in \mathbb{Z}[\zeta] \\ \frac{1-\zeta}{1-\zeta'} &= 1 + \zeta' + \cdots + \zeta'^{t-1} \in \mathbb{Z}[\zeta]. \end{aligned}$$

Therefore $\frac{1-\zeta'}{1-\zeta}$ is a unit in $\mathbb{Z}[\zeta]$. □

Proposition 6.2.2: Let p be prime and $r \in \mathbb{N}$. Suppose $p^r > 2$, let ζ_{p^r} be a primitive p^r -th root of unity, and let $K = \mathbb{Q}[\zeta_{p^r}]$. Then

1. $[\mathbb{Q}[\zeta_{p^r}] : \mathbb{Q}] = \varphi(p^r) = p^{r-1}(p-1)$.
2. The element $\pi = 1 - \zeta_{p^r}$ is prime in \mathcal{O}_K , and $(p) = (\pi)^{\varphi(p^r)}$.
3. $\mathcal{O}_K = \mathbb{Z}[\zeta_{p^r}]$.
4. $\text{disc}(\mathcal{O}_K/\mathbb{Z}) = (-1)^{\frac{\varphi(p^r)}{2}} p^{p^r-1(p^r-r-1)}$. Thus p is the only prime ramifying in $\mathbb{Q}[\zeta_{p^r}]$.

Proof. By Proposition 6.4.1,

$$\begin{aligned} p &= 1 + X^{p^{r-1}} + \cdots + X^{(p-1)p^{r-1}} \Big|_{X=1} \\ &= \Phi_{p^r}(1) \\ &= \prod_{\zeta' \text{ primitive } p^r \text{th root of unity}} (1 - \zeta') \\ &= \prod_{\zeta' \text{ primitive } p^r \text{th root of unity}} \frac{1-\zeta'}{1-\zeta_{p^r}} (1 - \zeta_{p^r}) \\ &= u(1 - \zeta)^{\varphi(p^r)} \end{aligned}$$

where $u = \prod_{\zeta' \text{ primitive } p^r \text{th root of unity}} \frac{1-\zeta'}{1-\zeta_{p^r}}$ is a unit by Proposition 6.2.1. Thus $(p) = (\pi)^{\varphi(p^r)}$.

From the degree equation (Theorem 2.5.2), we get that $[\mathbb{Q}[\zeta] : \mathbb{Q}] \geq \varphi(p^r)$ with strict inequality when π factors further. On the other hand $[\mathbb{Q}[\zeta] : \mathbb{Q}] \leq \varphi(p^r)$ since the cyclotomic polynomial has degree $\varphi(p^r)$. Hence equality must hold, and π must be prime, giving (1) and (2).

The degree equation for $(p) = (\pi)^{\varphi(p^r)}$ reads

$$\varphi(p^r) = f((\pi)/(p)) \cdot \varphi(p^r)$$

so we must have $f((\pi)/(p)) = 1$, i.e. the natural map

$$\mathbb{Z}/(p) \xrightarrow{\cong} \mathcal{O}_K/(\pi) \quad (6.1)$$

is an isomorphism.

We first calculate $\text{disc}(\mathbb{Z}[\zeta_p]/\mathbb{Z})$. By Proposition 1.4.4,

$$\begin{aligned} \text{disc}(\mathbb{Z}[\zeta_{p^r}]/\mathbb{Z}) &= \pm \text{Nm}_{\mathbb{Q}(\zeta_{p^r})/\mathbb{Q}}(\Phi'_{p^r}(\zeta)) \\ \Phi'_{p^r}(\zeta) &= \left(\frac{X^{p^r} - 1}{X^{p^{r-1}} - 1} \right)' \Big|_{x=\zeta} \\ &= \frac{p^r X^{p^r-1} (X^{p^{r-1}} - 1) - (X^{p^r} - 1) p^{r-1} X^{p^{r-1}-1}}{(X^{p^{r-1}} - 1)^2} \Big|_{X=\zeta_{p^r}} \\ &= \frac{p^r \zeta_{p^r}^{p^r-1}}{\zeta_{p^r}^{p^{r-1}} - 1} = \frac{p^r \zeta_p^{-1}}{\zeta_p - 1} \end{aligned}$$

where we set $\zeta_p = \zeta^{p^{r-1}}$; this is a primitive p th root of unity. We calculate the norm of each factor.

1. $\text{Nm}_{\mathbb{Q}(\zeta_p)/\mathbb{Q}}(p^r) = (p^r)^{[\mathbb{Q}(\zeta_p):\mathbb{Q}]} = p^{rp^{r-1}(p-1)}$.
2. $\text{Nm}_{\mathbb{Q}(\zeta_p)/\mathbb{Q}}(\zeta_p^{-1}) = \pm 1$ since ζ_p^{-1} is a unit.
3. $\text{Nm}_{\mathbb{Q}(\zeta_p)/\mathbb{Q}}(\zeta_p - 1) = p^{p^{r-1}}$: The minimal polynomial of $\zeta_p - 1$ over \mathbb{Q} is $\Phi_p(X + 1)$, whose constant term is $\Phi_p(1) = X^{p^{r-1}(p-1)} + \dots + X^{p^{r-1}} + 1|_{X=1} = p$. Hence by Proposition 1.2.3(1c), we have

$$\text{Nm}_{\mathbb{Q}(\zeta_p)/\mathbb{Q}}(\zeta_p - 1) = (\pm p)^{[\mathbb{Q}(\zeta_{p^r}):\mathbb{Q}(\zeta_p)]} = \pm p^{\frac{\varphi(p^r)}{\varphi(p)}} = \pm p^{p^{r-1}}.$$

Combining these we get

$$\text{disc}(\mathbb{Z}[\zeta_{p^r}]/\mathbb{Z}) = \text{Nm}_{\mathbb{Q}(\zeta_{p^r})/\mathbb{Q}} \frac{p^r \zeta_p^{-1}}{\zeta_p - 1} = \frac{p^{r(p-1)p^{r-1}} \cdot \pm 1}{\pm p^{p^{r-1}}} = \pm p^{p^{r-1}(pr-r-1)}. \quad (6.2)$$

By Proposition 1.3.2 (fix this a bit), we have

$$\pm p^{p^{r-1}(pr-r-1)} = \text{disc}(\mathcal{O}_K/\mathbb{Z}) = (\mathcal{O}_K : \mathbb{Z}[\zeta_{p^r}])^2 \text{disc}(\mathbb{Z}[\zeta]/\mathbb{Z}).$$

Hence both factors are powers of p up to sign. Since $(\mathcal{O}_K : \mathbb{Z}[\zeta_{p^r}])$ is a power of p , the quotient module is annihilated by a power of p , i.e. then

$$p^m \mathcal{O}_K \subseteq \mathbb{Z}[\zeta_{p^r}] \quad (6.3)$$

for some m . Note surjectivity in (6.1) gives $\mathcal{O}_K = \mathbb{Z} + \pi\mathcal{O}_K$ and hence

$$\mathcal{O}_K = \mathbb{Z}[\zeta_{p^r}] + \pi\mathcal{O}_K. \quad (6.4)$$

Suppose $\mathcal{O}_K = \mathbb{Z}[\zeta_{p^r}] + \pi^n\mathcal{O}_K$. Then substitution into (6.4) gives

$$\mathcal{O}_K = \mathbb{Z}[\zeta_{p^r}] + \pi\mathcal{O}_K = \mathbb{Z}[\zeta_{p^r}] + \pi(\mathbb{Z}[\zeta_{p^r}] + \pi^n\mathcal{O}_K) = \mathbb{Z}[\zeta_{p^r}] + \pi^{n+1}\mathcal{O}_K.$$

Hence by induction, $\mathcal{O}_K = \mathbb{Z}[\zeta_{p^r}] + \pi^n\mathcal{O}_K$ for all n . However, $(p) = (\pi)^{\varphi(p^r)}$ so this means $\mathcal{O}_K = \mathbb{Z}[\zeta_{p^r}] + p^n\mathcal{O}_K$ for all n . Taking $n = m$, (6.3) gives $\mathcal{O}_K = \mathbb{Z}[\zeta_{p^r}]$, proving (3). Together with (6.2), this gives (4). The second part of (4) now follows from Theorem 2.6.1 (A prime ramifies if and only if it divides the discriminant).

All embeddings of $\mathbb{Q}(\zeta_n)$ are complex, and there are $\varphi(n) = [\mathbb{Q}(\zeta_n) : \mathbb{Q}]$ of them. By Theorem 1.4.6(1), the sign is $(-1)^{\varphi(p^r)}$. \square

Now we prove the analogous result for $\mathbb{Q}(\zeta_n)$, for any $n \in \mathbb{N}$, by taking compositums of fields of the form $\mathbb{Q}(\zeta_{p^r})$.

Theorem 6.2.3. *Let $n, r \in \mathbb{N}$ with $n \not\equiv 2 \pmod{4}$ ¹. Let ζ_n be a primitive n th root of unity.*

1. $[\mathbb{Q}(\zeta_n) : \mathbb{Q}] = \varphi(n)$.
2. $\mathcal{O}_K = \mathbb{Z}[\zeta_n]$.
- 3.

$$\text{disc}(\mathcal{O}_K/\mathbb{Z}) = \frac{(-1)^{\frac{\varphi(n)}{2}} n^{\varphi(n)}}{\prod_{p|n} p^{\frac{\varphi(n)}{p-1}}}.$$

Moreover,

1. If $p \neq 2$, then p ramifies iff $p \mid n$.
2. If $p = 2$, then p ramifies iff $4 \mid n$.

Proof. Let $K = \mathbb{Q}(\zeta_n)$. Along with the theorem statement, we will show that if $n = p^r m$, $p \nmid m$, then


$$(p) = \left(\prod \mathfrak{P}_i \right)^{\varphi(p^r)} \quad (6.5)$$

for distinct primes \mathfrak{P}_i .

We induct on the number of prime factors of n . The case when n is a prime power is treated by Proposition 6.2.2. Suppose the theorem true for m and $p \nmid m$; consider $n = p^r m$.

¹If $n \equiv 2 \pmod{4}$, note $\mathbb{Q}(\zeta_n) = \mathbb{Q}(\zeta_{n/2})$.

Writing $\zeta_{p^r} = \zeta_n^m$ and $\zeta_m = \zeta_n^{p^r}$, we consider



By Proposition 6.2.2(2), $(p) = \mathfrak{p}^{\varphi(p^r)}$ in $\mathbb{Q}[\zeta_{p^r}]$, while by part 2, p splits into distinct factors. Matching factorizations in $\mathbb{Q}[\zeta_{p^r m}]$, we get that each $\mathfrak{p}_i \mathcal{O}_K$ must be a perfect $\varphi(p^r)$ th power. Hence $[\mathbb{Q}(\zeta_n) : \mathbb{Q}] \geq \varphi(p^r)$, and equality must hold. Then $[\mathbb{Q}(\zeta_n) : \mathbb{Q}] = \varphi(p^r)\varphi(m) = \varphi(n)$ showing (1).

Item (2) follows from Proposition 1.4.8 since by (3), $\text{disc}(\mathbb{Q}(\zeta_{p^r})/\mathbb{Q})$ and $\text{disc}(\mathbb{Q}(\zeta_m)/\mathbb{Q})$ are relatively prime. Item (3) follows from Proposition 1.4.8 as well. The factorization comes from the fact that since $[\mathbb{Q}(\zeta_{p^r m}) : \mathbb{Q}(\zeta_m)] = \varphi(p^r)$ and each \mathfrak{p}_i is the $\varphi(p^r)$ th power of an ideal, the degree equation says each \mathfrak{p}_i must actually be the $\varphi(p^r)$ th power of a *prime* ideal. \square

We now show a more precise version of (6.5), using Theorem 2.6.3.

Theorem 6.2.4. *Suppose that $n = p^r m$, where $p \nmid m$. Let*

$$f = \text{ord}_m(p).$$

Then the prime factorization of (p) in $\mathbb{Q}(\zeta_n)$ is

$$(p) = (\mathfrak{P}_1 \cdots \mathfrak{P}_g)^{\varphi(p^r)}$$

where \mathfrak{P}_j are distinct primes, each with residue degree f over \mathbb{Q} , and $g = \frac{\varphi(m)}{f}$.

Proof. $(\dagger)^2$ To use Theorem 2.6.3, we find the factorization of $\Phi_n(X)$ modulo p . We have

$$\Phi_n(X) = \prod_{j \pmod{\times n}} (X - \zeta_n^j) = \prod_{j \pmod{\times m}} \prod k \pmod{\times p^r} (X - \zeta_m^j \zeta_{p^r}^k). \quad (6.6)$$

Now note that

$$X - \zeta_m^j \zeta_{p^r}^k \equiv X - \zeta_m^j \pmod{\zeta_{p^r} - 1}.$$

Hence (6.6) gives

$$\Phi_n(X) \equiv \prod_{j \pmod{\times m}} (X - \zeta_m^j)^{\varphi(p^r)} \equiv \Phi_m(X)^{\varphi(p^r)} \pmod{\zeta_{p^r} - 1}.$$

²For an alternate proof see Example 11.1.6.

But both sides are in $\mathbb{Z}[X]$ so this congruence holds modulo $(\zeta_{p^r} - 1) \cap \mathbb{Z} = (p)$.

Now consider $\Phi_m(X) \pmod{p}$. Note that modulo p , $P(X) := X^m - 1$ has no repeated factors since it is relatively prime to $P'(X) = mX^{m-1} \neq 0$; hence its divisor $\Phi_m(X)$ has no repeated factors either. Note $\mathbb{F}_{p^r}^\times$ consists exactly of elements with $x^{p^r-1} = 1$, any root α of $\Phi_m(X)$ satisfies $\alpha^m = 1$ (but not $\alpha^{m'} = 1$ for $0 < m' < m$). Thus the smallest field extension \mathbb{F}_{p^r} containing α is hence the smallest r such that $m \mid p^r - 1$, i.e. $r = \text{ord}_m(p)$. The irreducible factors of $\Phi_m(X)$ have degree f , so f is the residue degree. The number of factors equals $\frac{\varphi(m)}{f}$, and this is the number of distinct prime divisors of (p) . \square

3 Subfields of cyclotomic extensions

Proposition 6.3.1: The Galois group of $\mathbb{Q}(\zeta_n)/\mathbb{Q}$ is

$$G(\mathbb{Q}(\zeta_n)/\mathbb{Q}) = (\mathbb{Z}/n\mathbb{Z})^\times.$$

Proof. The conjugates of ζ_n over \mathbb{Q} are ζ_n^k with $k \in (\mathbb{Z}/n\mathbb{Z})^\times$, the roots of Φ_n . The Galois group acts transitively on the conjugates, so for every $k \in (\mathbb{Z}/n\mathbb{Z})^\times$, there is a automorphism σ_k sending $\zeta_n \rightarrow \zeta_n^k$, and these are all the automorphisms (look at the degree). Since ζ_n generates $\mathbb{Q}(\zeta_n)$, the action of an automorphism on ζ_n determines it completely. It is clear that $k \rightarrow \sigma_k$ is an isomorphism $(\mathbb{Z}/n\mathbb{Z})^\times \rightarrow G(\mathbb{Q}(\zeta_n)/\mathbb{Q})$. \square

Proposition 6.3.2: The unique quadratic extension of \mathbb{Q} contained in $\mathbb{Q}(\zeta_p)$ is

$$\mathbb{Q}\left(\sqrt{(-1)^{\frac{p-1}{2}}p}\right).$$

Proof. By the fundamental theorem of Galois theory, a quadratic extension corresponds to a subgroup of index 2 in $(\mathbb{Z}/p\mathbb{Z})^\times \cong \mathbb{Z}/(p-1)\mathbb{Z}$, and there is exactly one such subgroup. If it equals $\mathbb{Q}(\sqrt{d})$, then the only primes ramifying are those dividing d ; since the only prime ramifying in $\mathbb{Q}(\zeta_p)$ is p , we must have $d = \pm p$.

To determine the sign, we explicitly find express a generator for $\mathbb{Q}(\sqrt{d})$ in terms of ζ_p . Define τ by the Gauss sum

$$\tau = \sum_{k=1}^{p-1} \left(\frac{k}{p}\right) \zeta_p^k.$$

An automorphism $\sigma \in G(L/K)$ is described by $\sigma(\zeta_p) = \zeta_p^j$ for some j ; we have

$$\sigma\tau = \sum_{k=1}^{p-1} \left(\frac{k}{p}\right) \zeta_p^{jk} = \sum_{k=1}^{p-1} \left(\frac{j^{-1}k}{p}\right) \zeta_p^k$$

so $\sigma\tau = \tau$ iff $\left(\frac{j}{k}\right) = 1$, which happens for exactly half the elements of $G(L/K)$. Hence τ indeed generates a quadratic field.³

³This gives motivation for the Gauss sum appearing in the proof of quadratic reciprocity.

Now, if $p \equiv 1 \pmod{4}$, we have $\left(\frac{-1}{p}\right) = 1$ and we can pair $\left(\frac{k}{p}\right) \zeta_p^k + \left(\frac{-k}{p}\right) \zeta_p^{-k} \in \mathbb{R}$, while if $p \equiv 3 \pmod{4}$, we have $\left(\frac{-1}{p}\right) = -1$ and $\left(\frac{k}{p}\right) \zeta_p^k + \left(\frac{-k}{p}\right) \zeta_p^{-k} \in \mathbb{R}i$. This gives the sign of d .

Alternatively, we can calculate τ explicitly as in (BLAH). \square

Proposition 6.3.3: For $n > 2$, $\mathbb{Q}(\zeta_n)$ is a CM-field with totally real subfield

$$\mathbb{Q}(\zeta_n + \zeta_n^{-1}) = \mathbb{Q}\left(\cos \frac{2\pi}{n}\right).$$

Proof. \square

4 Fermat's last theorem: Regular primes

Theorem 6.4.1. Any unit $u \in \mathbb{Z}[\zeta_n]$ can be written in the form

$$u = \zeta_n^k v$$

where v is totally positive, i.e. $\sigma(v) \in \mathbb{R}$ for any embedding $\sigma : \mathbb{Q}[\zeta_n] \rightarrow \mathbb{C}$.

Definition 6.4.2: A prime p is **regular** if p does not divide the class number of $\mathbb{Z}[\zeta_p]$.

Theorem 6.4.3 (First case of Fermat's last theorem for regular primes). *Suppose that $p > 2$ is a regular prime. Then any integer solution to*

$$x^p + y^p = z^p$$

satisfies $p \mid xyz$.

Proof. For $p = 3$, note that any cube must be congruent to 0 or ± 1 modulo 9. Hence in order for $x^3 + y^3 \equiv z^3 \pmod{9}$, one of x, y, z is divisible by 3, as needed.

Now assume $p > 3$. By dividing by $\gcd(x, y, z)$ we may assume x, y, z are relatively prime.

Step 1: Factor the equation as

$$\prod_{j=0}^{p-1} (x + \zeta_p^j y) = z^p. \tag{6.7}$$

(Note p is odd.) We show that if $p \nmid xyz$, then the factors on the left are relatively prime. Take $j \neq k$ and consider $\mathfrak{a} := \gcd((x + \zeta_p^j y), (x + \zeta_p^k y))$. We have

$$\mathfrak{a} \mid (x + \zeta_p^j y - x - \zeta_p^k y) = (\zeta_p^j - \zeta_p^k)(y).$$

Now x, y have no common factor in \mathbb{Z} , so (x) and (y) have no common factor in $\mathbb{Z}[\zeta_p]$, and $(x + \zeta_p^j y)$ and (y) have no common factor. This shows

$$\mathfrak{a} \mid (\zeta_p^j - \zeta_p^k).$$

The RHS is prime, so either $\mathfrak{a} = (\zeta_p^j - \zeta_p^k) = (1 - \mathfrak{p})$ or $\mathfrak{a} = (1)$. In the first case, we get $(1 - \mathfrak{p}) \mid \prod_{j=0}^{p-1} (x + \zeta_p^j y) = z^n$ so $p \mid z^n$, contradiction.

Step 2: By uniqueness of ideal factorization, each factor of (6.7) is a perfect p th power.

$$(x + \zeta_p^j y) = \mathfrak{a}_j^p$$

However, $p \nmid |C(\mathbb{Z}[\zeta_p])|$ so $C(\mathbb{Z}[\zeta_p])$ has no p -torsion. Since $(x + \zeta_p^j y)$ is a principal ideal, \mathfrak{a}_j must also be a principal ideal (a_j) . By Theorem 6.4.1, we can write

$$a_j = \zeta_p^{r_j} v_j, \quad v_j \in \mathbb{Q}[\zeta_p]^+.$$

□

5 Exercises

Problems

- 1.1 Let p be a prime. Prove that any equiangular p -gon with rational side lengths is regular.
- 1.2 (Komal) Prove that there exists a positive integer n so that any prime divisor of $2^n - 1$ is smaller than $2^{\frac{n}{1993}} - 1$.
- 1.3 Find all rational $p \in [0, 1]$ such that $\cos p\pi$ is...
 - (a) rational
 - (b) the root of a quadratic polynomial with rational coefficients
- 1.4 (China) Prove that there are no solutions to $2 \cos p\pi = \sqrt{n+1} - \sqrt{n}$ for rational p and positive integer n .
- 1.5 (TST 2007/3) Let θ be an angle in the interval $(0, \pi/2)$. Given that $\cos \theta$ is irrational and that $\cos k\theta$ and $\cos[(k+1)\theta]$ are both rational for some positive integer k , show that $\theta = \pi/6$.
- 2.1 Show that the ring of integers in $\mathbb{Q}(\cos \frac{2\pi}{n})$ is $\mathbb{Z}[\cos \frac{2\pi}{n}]$.

? Show that the class group of $\mathbb{Q}(\zeta_{23})$ (is this the right one?) is nontrivial.

Chapter 7

Valuations and completions

Here is some motivation for considering \mathfrak{p} -adic fields.

1. One useful tool in arithmetic geometry is the *local to global* principle, which says that the existence of solutions modulo all primes tells us something about the existence of solutions in the original field or ring, such as \mathbb{Q} or \mathbb{Z} . For example, the Hasse-Minkowski Theorem. However, it is not enough to check for solutions modulo all powers of p — because a solution modulo p does not necessarily give a solution modulo powers of p . The solution is to look for solutions in a field which contains information *modulo all powers of p* , a p -adic field.
2. When we take a \mathfrak{p} -adic fields, the only prime ideal remaining is \mathfrak{p} ; all others primes become units. This vastly simplifies algebraic number theory; we don't have to worry about primes that split. Then we can recover facts about the global field.

1 Case study: p -adic integers

We first examine how p -adic rationals are defined, before generalizing to other number fields.

Often we look at the integers modulo higher and higher powers of a prime p ; for example, when we were looking at the existence of primitive roots (Theorem 1.6.2) or the structure of $\mathbb{Z}/p^n\mathbb{Z}$ (Theorem 1.7.1). Hensel's lemma told us that under certain conditions we can lift solutions modulo higher and higher powers of p .

Rather than work with powers of p piecemeal, we can devise a structure that holds information modulo all powers of p at once. To do this, we define the ring p -adic integers \mathbb{Z}_p and p -adic rationals \mathbb{Q}_p , which contain \mathbb{Z} and \mathbb{Q} , respectively. We will do this in two ways:

1. Define \mathbb{Z}_p as an *inverse limit* of the rings $\mathbb{Z}/p^n\mathbb{Z}$ and \mathbb{Q}_p as the fraction field.
2. Give \mathbb{Q} a topology (or even better, a metric) related to divisibility by p , and complete \mathbb{Q} with respect to this topology.

1.1 p -adics as an inverse limit

Definition 7.1.1: A p -adic integer is a compatible sequence

$$(x_n)_{n \geq 1}$$

where $x_n \in \mathbb{Z}/p^n\mathbb{Z}$ and such that $x_{n+1} \equiv x_n \pmod{p^n}$ for all n , i.e. x_{n+1} maps to x_n under the projection map $\mathbb{Z}/p^{n+1}\mathbb{Z} \rightarrow \mathbb{Z}/p^n\mathbb{Z}$.

The ring structure is defined by componentwise addition and multiplication. The ring of p -adic integers is denoted by \mathbb{Z}_p and its fraction field is denoted by

$$\mathbb{Q}_p = \text{Frac}(\mathbb{Z}_p).$$

In light of Theorem 5.7.5, we can phrase this definition in a more abstract way:

$$\mathbb{Z}_p = \varprojlim \mathbb{Z}/p^n\mathbb{Z}$$

where there are maps $\varphi_n^m : \mathbb{Z}/p^m\mathbb{Z} \rightarrow \mathbb{Z}/p^n\mathbb{Z}$ given by projection whenever $m \geq n$.

1.2 p -adics as completions

We can give define a topology on \mathbb{Z} by decreeing that it be invariant under translation and that a neighborhood base of 0 be $\{p^n\mathbb{Z}, n \geq 0\}$. This is the same as the topology induced by the norm

$$|a|_p = p^{-v} \text{ when } a = \frac{p^v b}{c}, p \nmid b, c.$$

Definition 7.1.2 (Alternate definition of p -adics): \mathbb{Q}_p is the completion of \mathbb{Q} with respect to the p -adic norm.

We show the equivalence more generally in ().

1.3 Units in \mathbb{Z}_p

Proposition 7.1.3: The group of units in \mathbb{Z}_p is

$$\mathbb{Z}_p^\times \cong \begin{cases} \mathbb{Z}_p \times \mathbb{Z}/(p-1)\mathbb{Z}, & p \neq 2 \\ \mathbb{Z}_2 \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}, & p = 2. \end{cases}$$

Proof. Note that

$$\mathbb{Z}_p^\times = \varprojlim_{n \geq 1} (\mathbb{Z}/p^n\mathbb{Z})^\times$$

because any inverse modulo p^n can be lifted to an inverse modulo p^{n+1} .

The proposition follows from taking inverse limits in Theorem 1.7.1. □

1.4 Monsky's Theorem*

We use the 2-adic valuation to prove the following theorem from combinatorial geometry. Surprisingly, no proof is known that does not use p -adics.

Theorem 7.1.4. *A unit square cannot be cut into an odd number of triangles of equal area.*

The idea of the proof is as follows.

1. Extend the 2-adic valuation to a nonarchimedean valuation on the real numbers.
2. Color each point in the plane one of three colors, based on the 2-adic valuation of the coordinates. We show that the sides of the square only have two colors, with the vertices alternating colors, and that a triangle of area $\frac{1}{m}$ where m is odd, cannot contain vertices of all three colors. The last fact depends crucially on the fact that the area formula for a triangle has a factor of $\frac{1}{2}$ in it.
3. By Sperner's Lemma (from graph theory), the coloring in such a subdivision is inconsistent.

Proof. We omit the proof of the first item.¹ Assuming it, color the points of the plane in three colors depending on which of the following conditions is satisfied.

(A) $|x|_2 < 1, |y|_2 < 1$

(B) $|x|_2 \geq 1, |x|_2 \geq |y|_2$

(C) $|y|_2 \geq 1, |y|_2 > |x|_2$

First, we show that if $(\Delta x, \Delta y)$ has color A, then translating by $(\Delta x, \Delta y)$ does not change the color of A. Indeed, consider 3 cases.

1. (x, y) is of color A. By the nonarchimedean property, we have

$$|x + \Delta x|_2 \leq \max(|x|_2, |\Delta x|_2) \leq 1, |y + \Delta y|_2 \leq \max(|y|_2, |\Delta y|_2) \leq 1,$$

so $(x + \Delta x, y + \Delta y)$ is again of color A.

2. (x, y) is of color B. Since $|x|_2 \geq 1 > |\Delta x|_2$, we have

$$|x + \Delta x|_2 = |x|_2 \geq 1.$$

Since $|x|_2 \geq |y|_2$ and $1 > |\Delta y|_2$ we have

$$|y + \Delta y|_2 \leq \max(|y|_2, |\Delta y|_2) \leq |x|_2 = |x + \Delta x|_2.$$

Hence $(x + \Delta x, y + \Delta y)$ is again of color B.

¹There is a way around it; see Proofs from the Book.

3. (x, y) is of color C . The proof is the same as above except x, y are interchanged and there is strict inequality in the dotted inequalities above.

Now suppose that A, B, C are three points of those respective colors. By translation we may assume that $A = O$. Let $B = (x, y)$ and $C = (x', y')$. We have

$$\begin{aligned} |x|_2 &\geq |y|_2 \\ |y'|_2 &> |x|_2 \\ \implies |xy'|_2 &> |x'y|_2. \end{aligned}$$

1. A, B, C cannot be collinear, as that would imply $xy' = x'y$.
2. We show A, B, C cannot form a triangle of area $\frac{1}{m}$ for m odd. The area is $\pm\frac{1}{2}(xy' - x'y)$, and we have

$$\left|\frac{1}{2}(xy' - x'y)\right| = \left|\frac{1}{2}\right|_2 |x|_2 |y'|_2 > 1,$$

while $\left|\frac{1}{m}\right| = 1$.

Next we establish the following combinatorial lemma.

Lemma 7.1.5 (Sperner's lemma). *Suppose \mathcal{P} is a polygon that has been subdivided into triangles. Define a vertex or segment to be a vertex or edge of one of these triangles, and say a segment is of type $\mathcal{C}_1\mathcal{C}_2$ if the endpoints are colored \mathcal{C}_1 and \mathcal{C}_2 . We say a triangle is rainbow if it has vertices of all 3 colors.*

Suppose every vertex of the subdivision is colored with either \mathcal{A} , \mathcal{B} , or \mathcal{C} , such that the following hold.

1. *No outer edge of \mathcal{P} contains vertices of all three colors.*
2. *There are an odd number of segments of type \mathcal{AB} on the outer edges.*

Then \mathcal{P} contains a triangle whose vertices are all different colors.

Proof. We count the number of segments of type \mathcal{AB} . In a monochromatic triangle the count is 0, in a two-colored triangle the count is 0 or 2, and in a three-colored triangle the count is 1. Let n be the sum of the counts over all triangle. Every interior segment of type \mathcal{AB} is counted twice, as it is part of two triangles, so

$$n = 2i + e,$$

where i and e denote the number of interior and exterior segments of type \mathcal{AB} . Since e is odd by assumption, n is also odd. But this can only happen if there is a three-colored triangle. \square

Now the points $O = (0, 0)$, $X = (1, 0)$, $Y = (1, 1)$, and $Z = (0, 1)$ are colored with \mathcal{A} , \mathcal{B} , \mathcal{B} , \mathcal{C} , respectively. We've shown that each side contains segments of at most 2 colors; segments of type \mathcal{AB} can only appear on side OX and XY ; in the former there must be an odd number (since O, X are different colors) and in the latter there must be an even number. Thus the conditions of Sperner's Lemma are satisfied, and any subdivision must contain a rainbow triangle, which cannot have area $\frac{1}{m}$ for m odd. \square

2 Valuations

Definition 7.2.1: A **valuation** on a field K is a function $|\cdot| : K \rightarrow \mathbb{R}$ such that

1. $|x| \geq 0$ with equality only when $x = 0$.
2. $|xy| = |x||y|$.
3. $|x + y| \leq |x| + |y|$.

If the stronger condition $|x + y| \leq \max(|x|, |y|)$ holds, then $|\cdot|$ is **nonarchimedean**.

Example 7.2.2: For a number field K , any embedding $\sigma : K \hookrightarrow \mathbb{C}$ gives a valuation on K :

$$|a| := |\sigma a|.$$

Example 7.2.3: The **\mathfrak{p} -adic valuation** is

$$|a|_{\mathfrak{p}} = \left(\frac{1}{\mathfrak{N}\mathfrak{p}} \right)^{v_{\mathfrak{p}}(a)}.$$

In the special case $K = \mathbb{Q}$, $\mathfrak{p} = (p)$, we have

$$|a|_p = \left(\frac{1}{p} \right)^{v_p(a)}.$$

Proposition 7.2.4: A valuation is nonarchimedean if and only if it is bounded on \mathbb{Z} (in fact, bounded by 1). Hence if $\text{char}(K) \neq 0$, then K only has nonarchimedean valuations.

Proof. If $|\cdot|$ is archimedean, then $|1 + \cdots + 1| \leq |1| = 1$, so $|n| \leq 1$ for any $n \in \mathbb{Z}$.

Conversely, suppose that $|\cdot|$ is bounded on \mathbb{Z} , say by C . We have

$$\begin{aligned} |(a + b)^n| &= \left| \sum_{k=0}^n \binom{n}{k} a^k b^{n-k} \right| \\ &\leq \sum_{k=0}^n C |a|^k |b|^{n-k} \\ &\leq C(n + 1) \max(|a|, |b|)^n. \end{aligned}$$

Hence for all $n \geq 1$, $|a + b| \leq (C(n + 1))^{\frac{1}{n}} \max(|a|, |b|)$. Taking $n \rightarrow \infty$ gives the result. \square

Proposition 7.2.5 (Relationship between additive and multiplicative valuations): Fix a base b . There is a correspondence between additive and multiplicative valuations, given by

$$\begin{aligned} |x| &= b^{-v(x)} \\ v(x) &= -\log_b(x). \end{aligned}$$

Different values of b give equivalent valuations. If $v(K^\times)$ is discrete in \mathbb{R} , then it is a multiple of a discrete valuation.

We say $|\cdot|$ is discrete when $|K^\times|$ is a discrete subgroup of $\mathbb{R}_{>0}$.

Proposition 7.2.6 (Valuation defines subrings): Suppose K is equipped with a nonarchimedean valuation. Then

1. $A := \{a \in K : |a| \leq 1\}$ is a subring of K , with
2. $U := \{a \in K : |a| = 1\}$ is a group, and
3. $\mathfrak{p} := \{a \in K : |a| < 1\}$ is a prime ideal.

Proof. Straightforward. □

Proposition 7.2.7 (Elementary properties of discrete valuations):

1. $|a + b| \leq \max(|a|, |b|)$ with equality if $|a| \neq |b|$.
2. (“All triangles are isosceles.”) If $d(c, b) < d(c, a)$ then $d(a, c) = d(a, b)$. (The longer side is the repeated one.)
3. If $a_1 + \cdots + a_n = 0$, then the maximum valuation of the summands must be attained for at least two of them.

Proof. Straightforward. □

2.1 Equivalent valuations

A valuation on K defines a metric (and hence a topology) on K by

$$d(a, b) = |a - b|.$$

For example, high powers of p have small p -adic valuation, so numbers differing by high powers of p are close together in the p -adic valuation.

Proposition 7.2.8: Let $|\cdot|_1, |\cdot|_2$ be valuations on K , with the first being nontrivial. Then the following are equivalent.

1. $|\cdot|_1, |\cdot|_2$ determine the same topology on K .
2. If $|\alpha|_1 < 1$, then $|\alpha|_2 < 1$.
3. $|\cdot|_1 = |\cdot|_2^a$ for some $a > 0$.

We say that $|\cdot|_1$ and $|\cdot|_2$ are **equivalent** if the above conditions hold.

Proof.

(1) \implies (2): Note $|\alpha|_j < 1$ if and only if $|\alpha^n|_j = |\alpha|_j^n \rightarrow 0$, i.e. α^n converges to 0 in the topology of $|\cdot|_j$. Since the topologies are the same,

$$|\alpha|_1 < 1 \iff \alpha^n \text{ converges to } 0 \iff |\alpha|_2 < 1.$$

(2) \implies (3): Take y so that $|y|_1 > 1$, and let $a = \frac{|y|_2}{|y|_1}$, so that $|y|_2 = |y|_1^a$. We show that $|x|_2 = |x|_1^a$ for all $x \in K$.

Suppose $|x|_1 = |y|_1^{b_1}$ and $|x|_2 = |y|_2^{b_2}$. We need to show $b_1 = b_2$, i.e. so the following commutes.

$$\begin{array}{ccc} |x|_1 & \xrightarrow{\wedge a} & |x|_2 \\ \uparrow \wedge b_1 & & \uparrow \wedge b_2 \\ |y|_1 & \xrightarrow{\wedge a} & |y|_2 \end{array}$$

We approximate b_1 with rational numbers $\frac{m}{n}$. First suppose $b_1 > \frac{m}{n}$. Then

$$\left| \frac{y^m}{x^n} \right|_1 = |y|_1^{m-b_1n} < 1$$

so by hypothesis

$$|y|_2^{m-b_2n} = \left| \frac{y^m}{x^n} \right|_2 < 1$$

giving $b_2 > \frac{m}{n}$. Similarly, if $b_1 < \frac{m}{n}$, then the above argument with $\frac{x^n}{y^m}$ shows $b_2 < \frac{m}{n}$. Since \mathbb{Q} is dense in \mathbb{R} , we have $b_1 = b_2$.

(3) \implies (1): The open ball of radius r with respect to $|\cdot|_1$ is the same as the open ball of radius r^a with respect to $|\cdot|_2$. \square

3 Places

Definition 7.3.1: A **place** is an equivalence class of nontrivial valuations on K .² We denote by V_K the set of places of K , by V_K^0 the set of nonarchimedean places and V_K^∞ the set of archimedean places.

We aim to classify all places in a number field K .

Proposition 7.3.2: Let K/\mathbb{Q} be an algebraic extension. Then the places on K are exactly the \mathfrak{p} -adic valuations $|\cdot|_{\mathfrak{p}}$ for \mathfrak{p} a prime ideal of \mathcal{O}_K .

²Some books use “prime” instead of “place.” We use the latter term to avoid confusion.

Proof. Since K is algebraic over \mathbb{Q} , an element $\alpha \in \mathcal{O}_K$ satisfies a monic polynomial equation with coefficients in \mathbb{Z} :

$$x^n + a_{n-1}x^{n-1} + \cdots + a_0 = 0.$$

By Proposition 7.2.4, $a_j \in \mathbb{Z}$ gives $|a_j| \leq 1$. By the nonarchimedean property,

$$|\alpha|^n = |a_{n-1}\alpha^{n-1} + \cdots + a_0| \leq \max_{0 \leq m \leq n-1} |a_m| |\alpha|^m \leq \max_{0 \leq m \leq n-1} |\alpha|^m.$$

Hence $|\alpha| < 1$. This means if A is as in Proposition 7.4.5—the ring of integers of $|\cdot|$, then $\mathcal{O}_K \subseteq A$. Let \mathfrak{P} be the prime ideal corresponding to $|\cdot|$. Since \mathfrak{P} is prime in A , $\mathfrak{p} := \mathfrak{P} \cap \mathcal{O}_K$ is prime in \mathcal{O}_K . Note $\mathfrak{p} \neq (0)$ because if so $|\cdot|$ is trivial.

Now suppose $v_{\mathfrak{p}}(y) = n$. Let $\pi \in \mathfrak{p} \setminus \mathfrak{p}^2$ be a uniformizer. Then $(y\pi^{-n})$ is a fractional ideal; suppose ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_m$ appear in its factorization with exponents at least $-k$. Take $b \in \bigcap_{j=1}^m \mathfrak{p}_j^k$. Then $(y\pi^{-n}b)$ is an integral ideal (c) not divisible by \mathfrak{p} . We have $c \in A \setminus \mathfrak{p}$. Writing $|\pi| = \left(\frac{1}{\mathfrak{N}\mathfrak{p}}\right)^a$, we have

$$|y| = \left|\frac{c}{b}\right| |\pi^n| = \left|\frac{1}{\mathfrak{N}\mathfrak{p}}\right|^n = |y|_{\mathfrak{p}}^a.$$

Moreover, two equivalent nonarchimedean valuations would have the same maximal ideals and hence correspond to the same prime \mathfrak{p} . \square

Theorem 7.3.3 (Ostrowski). *The following is a list of all places on \mathbb{Q} .*

1. Archimedean: $|\cdot|_{\infty}$.³
2. Nonarchimedean: $|\cdot|_p$, where p ranges over all primes.

Proof. Let $|\cdot|$ be a valuation on \mathbb{Q} and m, n be integers greater than 1. To compare $|m|$ and $|n|$, we write m in base n :

$$m = a_r n^r + \cdots + a_0, \quad 0 \leq a_k \leq n-1, \quad a_r > 0.$$

Let $N = \max\{1, |n|\}$. Then by the triangle inequality,

$$|m| \leq \sum_{k=0}^r a_k N^k.$$

Since $r \leq \frac{\ln m}{\ln n}$, we get

$$|m| \leq \left(1 + \frac{1}{N} + \frac{1}{N^2} + \cdots\right) n N^{\frac{\ln m}{\ln n}} \leq 2n N^{\frac{\ln m}{\ln n}}$$

³A stronger version of part 1 is as follows. Let K be complete with respect to an archimedean norm. Then $K = \mathbb{R}$ or \mathbb{C} , and the norm is the normal absolute value raised to a power in $(0, 1]$.

Replacing m by m^t and taking the t th root gives

$$|m| \leq (2n)^{\frac{1}{t}} N^{\frac{\ln m}{\ln n}}.$$

Taking $t \rightarrow \infty$ gives

$$|m| \leq N^{\frac{\ln m}{\ln n}}. \quad (7.1)$$

Consider two cases.

1. For all integers $n > 1$, $|n| > 1$. Then (7.1) gives $|m|^{\frac{1}{\ln m}} \leq |n|^{\frac{1}{\ln n}}$. By symmetry, we get $|m|^{\frac{1}{\ln m}} = |n|^{\frac{1}{\ln n}}$. Since this is true for all m and n , $|n|^{\frac{1}{\ln n}} = c$ is constant, i.e.

$$|n| = c^{\ln n} = n^{\frac{\ln n}{\ln c}}$$

for all $n \in \mathbb{Z}$. Since \mathbb{Z} generates \mathbb{Q} as a group, we get that $|\cdot|$ is equivalent to the standard archimedean valuation.

2. For some $n > 1$, $|n| \leq 1$. Then (7.1) shows that $|m| \leq 1$ for all $m > 1$. Thus by Proposition 7.2.4, $|\cdot|$ is nonarchimedean. The nonarchimedean valuations are given by Proposition 7.3.2. \square

Later on we will return to the question of finding all valuations on an extension of \mathbb{Q} (Theorem 7.4.2).

3.1 Approximation

Theorem 7.3.4 (Weak approximation theorem). *Let v_1, \dots, v_n be all the places of K , with valuations $|\cdot|_1, \dots, |\cdot|_n$. The map*

$$\phi : K \rightarrow \prod_{j=1}^n K_{v_j}$$

induced by the inclusions $K \hookrightarrow K_{v_j}$ has dense image.

In other words, given $a_1, \dots, a_n \in K$, for any $\varepsilon > 0$, there exists $a \in K$ such that

$$|a - a_j|_j < \varepsilon \text{ for all } j.$$

Proof. Step 1: We show that there exists a such that

$$\begin{aligned} |a|_1 &> 1, \\ |a|_j &< 1, \quad i = 2, \dots, n. \end{aligned} \quad (7.2)$$

We induct on n . For $n = 2$, note that by Proposition 7.2.8(2), we can find b, c so that

$$\begin{aligned} |b|_1 &< 1, & |b|_2 &\geq 1 \\ |c|_1 &\geq 1, & |c|_2 &< 1. \end{aligned}$$

Now take $a = \frac{c}{b}$.

For the induction step, suppose we've found b so that (7.2) holds for $n - 1$. Choose c so that

$$|c|_1 > 1, \quad |c|_n < 1;$$

we will use it to "correct" $|b|_n$ as necessary. Consider three cases.

1. $|b|_n < 1$: We can let $a = b$.
2. $|b|_n = 1$: Let $a = b^r c$, for large enough r . This works because

$$\lim_{r \rightarrow \infty} |b^r c|_j = \begin{cases} \infty, & j = 1 \\ 0, & 2 \leq j \leq n - 1 \\ |c|_n < 1, & j = n. \end{cases}$$

3. $|b|_n > 1$: First note that from $1 - |a^r| \leq |1 + a^r| \leq 1 + |a^r|$ we get

$$\lim_{r \rightarrow \infty} \left| \frac{x^r}{1 + x^r} \right| = \begin{cases} 0, & |x| < 1 \\ 1, & |x| > 1. \end{cases} \quad (7.3)$$

Let $a = \frac{cb^r}{1+b^r}$, for large enough r . This works because the above gives

$$\lim_{r \rightarrow \infty} \left| \frac{cb^r}{1 + b^r} \right|_j = \begin{cases} |c|_1 > 1, & j = 1 \\ 0, & 2 \leq j \leq n - 1 \\ |c|_n < 1, & j = n. \end{cases}$$

Step 2: Now we show that there are points in the image of ϕ arbitrarily close to $(1, 0, \dots, 0)$. Indeed, choosing a as in step 1, we have by (7.3) that

$$\lim_{r \rightarrow \infty} \varphi \left(\frac{a^r}{1 + a^r} \right) = (1, 0, \dots, 0).$$

Step 3: From step 2, choose b_j sufficiently close to $(0, \dots, 0, \underbrace{1}_j, 0, \dots, 0)$. Let

$$a = \sum_{j=1}^n a_n b_n$$

to find $\varphi(a)$ can be arbitrarily close to (a_1, \dots, a_n) . □

Note that if we include only the finite places, then this follows from the Chinese remainder theorem.

4 Completion

Definition 7.4.1: Let K be a field with valuation $|\cdot|$. The **completion** of K , denoted \hat{K} is the field containing K (i.e. there is an injection $K \hookrightarrow \hat{K}$ preserving valuation) satisfying the following properties.

1. \hat{K} is complete in its topology.
2. (UMP) For any homomorphism φ from K to a complete field L , there exists a unique homomorphism $\hat{K} \rightarrow L$ making the following commute.

$$\begin{array}{ccc} \hat{K} & \xrightarrow{\quad \quad} & L \\ \uparrow & \nearrow & \\ K & & \end{array} .$$

I.e., \hat{K} is the smallest complete field containing K .

Proof of existence. For existence, let \hat{K} be the set of equivalence classes of Cauchy sequences in K , and deem two sequences $\{a_n\}$ and $\{b_n\}$ equivalent if $\lim_{n \rightarrow \infty} |a_n - b_n| = 0$. Define $K \hookrightarrow \hat{K}$ by sending a to (a, a, \dots) . Extend the valuation by letting defining the norm of a $\{a_n\}$ to be $\lim_{n \rightarrow \infty} |a_n|$. See any book on real analysis for the details.

For the second part, given a sequence $\{a_n\} \in \hat{K}$, map it to $\lim_{n \rightarrow \infty} \varphi(a_n) \in L$. Uniqueness follows from the universal property. \square

4.1 Completions of archimedean fields; Classification of places

Theorem 7.4.2 (Ostrowski). *The only complete archimedean fields, up to isomorphism of valued fields and equivalence of valuation, are \mathbb{R} and \mathbb{C} .*

Proof. Suppose K is a complete archimedean field. Then on \mathbb{Q} , by the first Ostrowski Theorem 7.3.3, $\|\cdot\|$ is equivalent to the usual valuation $\|\cdot\|_\infty$. By Proposition 7.2.8, by raising to a power we may assume it is exactly the usual valuation. We must have $\hat{\mathbb{Q}} = \mathbb{R} \subseteq K$.

We show every element $\xi \in K$ satisfies a quadratic equation over \mathbb{R} . This is equivalent to saying that $\min_{z \in \mathbb{C}} f(z) = 0$ where

$$f(z) = |\xi^2 - (z + \bar{z})\xi + z\bar{z}|.$$

Let $m = \min |f(z)|$; by way of contradiction suppose $m \neq 0$. Note $S = \{z : f(z) = m\}$ is compact; let z_0 be an element of S of maximal absolute value.

The idea is the following: perturb (the n th power of) $\xi^2 - (z_0 + \bar{z}_0)\xi + z_0\bar{z}_0$ so that the valuation is very close to m^n ; taking $n \rightarrow \infty$ we force all roots to have absolute value m ; however, we can ensure one of the roots has larger absolute value, a contradiction.

Consider the polynomial

$$G(x) = (x^2 - (z_0 + \bar{z}_0)x + z_0\bar{z}_0)^n - (-\varepsilon)^n.$$

We have by the triangle inequality

$$G(\xi) \leq m^n + \varepsilon^n \implies |G(\xi)| \leq (m^n + \varepsilon^n)^2.$$

On the other hand, note $|G(x)|^2$ factors into quadratics so by definition of m , we have $|G(\xi)|^2 \geq m^{2n}$. We can do better: let z_1 be the root of $x^2 - (z_0 + \bar{z}_0)x + z_0\bar{z}_0 + \varepsilon = 0$; we have $|z_1| > m = |z_0|$. Now z_1 is a root of $G(x)$, so one of the quadratic factors of $|G(x)|^2$ is $x^2 - (z_1 + \bar{z}_1)x + z_1\bar{z}_1$ which by the assumption on z_0 , must be $> m$ at $x = \xi$. We get

$$m^{2n-1}(\xi^2 - (z_1 + \bar{z}_1)\xi + z_1\bar{z}_1) \leq |G(\xi)|^2 \leq (m^n + \varepsilon^n)^2$$

which is a contradiction for n large enough. \square

We can now finish our classification of places on K/\mathbb{Q} .

Theorem 7.4.3 (Classification of places of K). *Let K be a number field. There is exactly one place of K for each*

1. prime ideal \mathfrak{p} ,
2. real imbedding, and
3. conjugate pair of complex embeddings.

The valuations corresponding to prime ideals, i.e. \mathfrak{p} -adic valuations, are called **finite places**, while the those corresponding to real and complex embeddings are called **infinite (real or complex) places**.

Proof. The nonarchimedean valuations of K are given by Proposition 7.3.2, while each archimedean valuation v corresponds to an embedding (respecting valuations)

$$K \hookrightarrow K_v \cong \mathbb{R} \text{ or } \mathbb{C},$$

the isomorphism coming from Theorem 7.4.2. Note that complex conjugate embeddings give the same valuation. \square

Corollary 7.4.4. *Let L/K be extensions of number fields. If v is a place corresponding to a prime \mathfrak{p} of K , then the places $w \mid v$ in L correspond to primes $\mathfrak{P} \mid \mathfrak{p}$. If v is a place of K corresponding to an embedding $\sigma : K \rightarrow \mathbb{R}$ or \mathbb{C} , then the places $w \mid v$ correspond to σ to L .*

4.2 Completions of nonarchimedean fields

Suppose K is a field with a discrete nonarchimedean valuation $|\cdot|$. Since K is dense in \hat{K} and

$$|K \setminus \{0\}| = \{|\pi|^m : m \in \mathbb{Z}\}$$

is discrete in \hat{K} , we get $|K| = |\hat{K}|$.

Proposition 7.4.5 becomes much nicer when we look at the completion \hat{K} .

Proposition 7.4.5 (Valuation defines subrings): Let \hat{K} be complete with a nonarchimedean valuation. Then

1. $A = \mathcal{O}_{\hat{K}} := \{a \in \hat{K} : |a| \leq 1\}$ is a subring of \hat{K} , with
2. $U = \mathcal{O}_{\hat{K}}^\times := \{a \in \hat{K} : |a| = 1\}$ the unit group, and
3. $\mathfrak{m} := \{a \in K : |a| < 1\}$ its unique maximal ideal

Let π be a **uniformizer**, i.e. the largest element of K with $|\pi| < 1$. Then π generates \mathfrak{m} .

Proposition 7.4.6: Suppose K is a field with a discrete nonarchimedean valuation $|\cdot|$. Let S be a set of representatives for A/\mathfrak{m} . Then every element of \hat{K} has a unique expression in the form

$$\sum_{n \geq N} a_n \pi^n.$$

(More precisely, the sum represents $\lim_{m \rightarrow \infty} \sum_{n=N}^m a_n \pi^n$.) The norm is given by

$$\left| \sum_{n \geq N} a_n \pi^n \right| = |\pi|^N, \quad a_N \neq 0.$$

In other words, we can write elements of \hat{K} as “numbers with infinite π -expansions going off to the left,” as we saw in section 1.

Proof. Suppose $|x_n| = |\pi|^n$. Then $x\pi^{-n} \in A$, so there exists $s \in S$ with $x\pi^{-n} - a \in \mathfrak{m}$. Then

$$|x_n - a_n \pi^n| = |\pi|^{n+1};$$

set $x_{n+1} = x_n - a_n \pi^n$. Starting this process at $x = x_N$, we get

$$x = \sum_{n \geq N} a_n \pi^n.$$

We have $x - x_N \pi^N \in \pi^{-N+1}$ so $|x| = |\pi|^N$. □

We have a surprising result that is not true in the archimedean case.

Proposition 7.4.7: Let K be a field complete with respect to a discrete nonarchimedean valuation. Suppose $x_n \rightarrow 0$. Then $\sum_{n=1}^{\infty} x_n$ converges.

Proof. We show $s_n = \sum_{m=1}^n x_m$ be the sequence of partial sums. We show s_n is Cauchy. For each $\varepsilon > 0$ there is N such that for $n \geq N$, $|x_n| \leq \varepsilon$. Then by the nonarchimedean property, for all $n' > n \geq N$, $|s_{n'} - s_n| = |x_{n+1} \cdots + x_{n'}| \leq \varepsilon$. Thus s_n is Cauchy and hence converges. □

Thus we have two ways to think of a \mathfrak{p} -adic valuation.

We've thought of it analytically: give K a metric and take the completion of K as a metric space to obtain \widehat{K} . The algebraic notion of completion is the following: the completion of a ring A with respect to an ideal \mathfrak{p} is defined as $\hat{A} = \varprojlim_{n \geq 0} A/\mathfrak{p}^n$.

To connect up the analytic and algebraic definitions of the completion, note that $\hat{A} = \varprojlim_{n \geq 0} A/\mathfrak{p}^n$ is the completion of \hat{A} with the topology given by the neighborhood base $\{\mathfrak{p}^n\}_{n \geq 0}$; this gives A the structure of a topological ring.

Now if K is a field with a discrete nonarchimedean valuation, then we can define A and \mathfrak{p} as in Proposition 7.4.5. The topology given by the valuation is exactly the topology that has the neighborhood base of 0 being $\{\mathfrak{p}^n\}_{n \geq 0}$. This shows the following.

Proposition 7.4.8: Let K be a field with a discrete nonarchimedean valuation. Then defining A, \mathfrak{p} as in Proposition 7.4.5,

$$\widehat{K} = \text{Frac}(\varprojlim A/\mathfrak{p}^n).$$

Definition 7.4.9: Define the exponential function as a power series

$$e^x = \sum_{n=1}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \cdots .$$

We investigate the convergence of e^x .

Proposition 7.4.10: e^x converges for $v_p(x) > \frac{1}{p-1}$.

Proof. Writing $a = a_r p^r + \cdots + a_0$ in base p , we find by Example ?? that

$$v_p(n!) = \frac{n - \sum_{i=0}^r a_i}{p-1}.$$

Hence

$$v_p\left(\frac{x^n}{n!}\right) = n v_p(x) - \frac{n - \sum_{i=0}^r a_i}{p-1} = n \left(v_p(x) - \frac{1}{p-1} \right) + o(n).$$

By Proposition 7.4.7, e^x converges if and only if $v_p\left(\frac{x^n}{n!}\right) \rightarrow -\infty$; this gives the result. \square

5 Hensel's lemma

The following is the first version of Hensel's lemma for π -adics.

Lemma 7.5.1 (Hensel's lemma, I). *Let A be a complete local ring, $f(X) \in A[X]$, and a_0 be a simple root of $f(X)$ modulo π , i.e. $f(a_0) \equiv 0 \pmod{\pi}$ and $f'(a_0) \not\equiv 0 \pmod{\pi}$. Then there exists a unique root a of $f(X)$ with $a \equiv a_0 \pmod{\pi}$.*

Note this can be generalized as follows: Suppose $f(a_0) \equiv 0 \pmod{\pi^n}$ and $v_\pi(f'(a_0)) = k < n$. Then there is a unique root a of $f(X)$ with $a \equiv a_0 \pmod{\pi^{n-k}}$. The proof is the same, and is left to the reader!

Proof. We find zeros of $f(X)$ modulo higher and higher powers of π .

Using induction, we find a_n satisfying

$$f(a_n) \equiv 0 \pmod{\pi^{n+1}}.$$

The base case holds by hypothesis. For the induction step, note that by Taylor expansion of polynomials,

$$\begin{aligned} f(a_n + h\pi^{n+1}) &= f(a_n) + h\pi^{n+1}f'(a_n) + \cdots \\ &\equiv f(a_n) + h\pi^{n+1}f'(a_n) \pmod{\pi^{n+2}}. \end{aligned}$$

Since $f'(a_n) \not\equiv 0 \pmod{\pi}$ and $f(a_n) \equiv 0 \pmod{\pi^{n+1}}$, we can choose h so that this is 0 modulo π^{n+1} . (Explicitly, $h = -\frac{f(a_n)}{\pi^{n+1}} \cdot \frac{1}{f'(a_n)}$.) We let $a_{n+1} = a_n + h\pi^{n+1}$. By construction, the sequence a_n converges; let a be its limit. Since $a \equiv a_n \pmod{\pi^n}$, we get $f(a) \equiv f(a_n) \equiv 0 \pmod{\pi^{n+1}}$ for all n , and therefore $f(a) = 0$. \square

The first form of Hensel's lemma tells us about lifting a root a_0 of \bar{f} (f modulo π) to a root a of f in K . We can think of this as lifting a linear factor $x - a_0$ of \bar{f} to a linear factor $x - a$ of f . A stronger form of Hensel's lemma says that we can in fact lift any factor of \bar{f} to one of f .

Theorem 7.5.2 (Hensel's lemma, II). *Let k be the residue field of A and f be a monic polynomial. If $\bar{f} = g_0h_0$ where g_0 and h_0 are monic and relatively prime, then $f = gh$ for some g and h such that $\bar{g} = g_0$ and $\bar{h} = h_0$.*

If $\bar{f} = g_1 \cdots g_n$ is the complete factorization of \bar{f} in $k[X]$, then the complete factorization of f in $K[X]$ is $f = f_1 \cdots f_n$ where $\bar{f}_j = g_j$.

Proof. First we need the following lemma, which tells us that if the reductions of polynomials are relatively prime, then so are the original polynomials.

Lemma 7.5.3. *Let A be a local ring with residue field k . If $g, h \in A[X]$ are such that \bar{g} and \bar{h} are relatively prime, then g and h are relatively prime in $A[X]$ and there exist polynomials u, v with $\deg u < \deg h$ and $\deg v < \deg g$ such that*

$$ug + vh = 1.$$

Proof. Since \bar{g} and \bar{h} are relatively prime in $k[X] = (A/\mathfrak{m})[X]$, $(\bar{g}, \bar{h}) = A[X]/\mathfrak{m}A[X]$ and $(g, h) + \mathfrak{m}A[X] = A[X]$. Nakayama's lemma says that if A is a local ring with maximal ideal \mathfrak{m} and $M \supseteq N$ are A -modules with $M = N + \mathfrak{m}M$, then $M = N$. Apply this with $M = A[X]$, which is finitely generated over A to get $A[X] = (g, h)$. This means we can choose u, v such that $ug + vh = 1$. To satisfy the degree condition, drop all terms with higher degree. \square

We proceed as in the proof of Theorem 7.5.1. Suppose we have found g_n and h_n such that

$$f \equiv g_n h_n \pmod{\pi^{n+1}}.$$

We have

$$(g_n + v\pi^{n+1})(h_n + u\pi^{n+1}) \equiv g_n h_n + (ug_n + vh_n)\pi^{n+1} \pmod{\pi^{n+2}}.$$

By Lemma 7.5.3 we can choose u and v such that the above is congruent to f modulo π^{n+2} . Again let $g_{n+1} = g_n + v\pi^{n+1}$, $h_{n+1} = h_n + u\pi^{n+1}$, and take the limit as $n \rightarrow \infty$.

The second part follows from induction. Note $f = f_1 \cdots f_n$ is the complete factorization because any factorization of f gives a factorization for \bar{f} . \square

Definition 7.5.4: A **henselian field** is a field with nonarchimedean valuation v which satisfies Hensel's Lemma (with \mathfrak{p} the maximal ideal corresponding to v).

Hensel's lemma says that a field that is complete with respect to a discrete valuation is henselian.

6 Extending valuations

Theorem 7.6.1 (Extending discrete valuations). *Let K be henselian and let L/K be finite separable of degree n . Then $|\cdot|_K$ extends uniquely to a discrete valuation $|\cdot|_L$ on L , given by*

$$|\beta|_L = |\mathrm{Nm}_{L/K} \beta|_K^{\frac{1}{n}}.$$

Proof. Neukirch, pg. 131-132. \square

Definition 7.6.2: Let K be henselian. Let $v : K^\times \rightarrow \mathbb{Z}$ be the corresponding additive valuation, extended to $K^{\mathrm{al}\times} \rightarrow \mathbb{Q}$. Given a polynomial

$$f(X) = X^n + a_{n-1}X^{n-1} + \cdots + a_0 \in K[X]$$

define the **Newton polygon** of $f(X)$ to be the lower convex hull⁴ of

$$P_i := (i, v(a_i)).$$

Proposition 7.6.3: Suppose the bottom of the Newton polygon has segments of x -length n_i and slope $-s_i$. Then

1. $f(X)$ has exactly n_i roots $\alpha \in K^{\mathrm{al}}$ with $v(\alpha) = s_i$, and
2. $f_i(X) = \prod_{v(\alpha_i)=s_i} (X - \alpha_i)$ has coefficients in K .

Proof. We prove the following statement by induction: if $f(X) = \prod (X - \alpha_j) \in \bar{K}[X]$ and exactly n_i of the roots α_j have order equal to s_i , then the Newton's polygon of $f(X)$ has a segment of slope $-s_i$ and x -length n_i .

⁴draw the convex hull, and remove the segments joining $(0, v(a_0))$ and $(n, 0)$ from the top

The case $n = 1$ follows since the only line segment on the bottom joins $(0, v(\alpha_i))$ and $(1, 0)$. Now suppose the claim proved for n . Consider


$$g(X) = (X - \alpha)f(X) = \sum_{k=0}^{n+1} (a_{k-1} - \alpha a_k) X^k$$

(where nonexistent coefficients are set to 0). Let $t = v(\alpha)$. Let k_0 be the point such that the slopes of the line segments of Newton's polygon N for $k < k_0$ are $s \leq -t$, and such that the slopes of the line segments of N for $k > k_0$ are greater than $s > -t$. Let

$$\begin{aligned} d_k &= v(a_k) \\ \ell_k &= y\text{-value of intersection of } N \text{ with } x = k \\ d'_k &= v(a_{k-1} - \alpha a_k) \\ \ell'_k &= \begin{cases} \ell_k + t, & 0 \leq k \leq k_0 \\ \ell_{k-1} & k_0 < k \leq n. \end{cases} \end{aligned}$$

Let N' be the broken line formed by joining (k, ℓ'_k) . N' consists of segments of the same slopes as N , plus one more segment of slope $-t$ and x -length 1, in increasing order. It suffices to show that N' is the lower convex hull of the points (k, d'_k) .

Here is an example with $p = 5$, $f(X) = (X - 5)(X - 10)(X - 15)(X - 125)$ and $\alpha = 25$.⁵



Consider 2 cases. We will use

$$d'_k = v(a_{k-1} - \alpha a_k) \geq \min(v(a_{k-1}), v(\alpha a_k)) = \min(d_{k-1}, d_k + t),$$

with equality holding if $d_{k-1} \neq d_k + t$.

1. $k \leq k_0$: We have

$$\begin{aligned} d_{k-1} &\geq \ell_{k-1} \stackrel{(*)}{\geq} \ell_k + t = \ell'_k, \\ d_k + t &\geq \ell_k + t = \ell'_k \end{aligned}$$

⁵Of course, f does not have to split over $\mathbb{Q}[X]$ and the valuations don't have to be integers.

where in (*) we use the fact that the slope of the segment $(k-1, \ell_{k-1})(k, \ell_k)$ is at most $-t$. Hence (k, d'_k) lies above N' . Now suppose (k, d_k) lies on a corner of L (excluding $k = k_0$). Then $d_k = \ell_k$ and inequality holds in (*):

$$d_{k-1} > \ell_k + t = \ell'_k = d_k + t$$

so $d'_k = \ell'_k$ and (k, d'_k) lies on N' .

2. $k > k_0$: We have

$$\begin{aligned} d_{k-1} &\geq \ell_{k-1} = \ell'_k \\ d_k + t &\geq \ell_k + t \stackrel{(*)}{>} \ell_{k-1} = \ell'_k. \end{aligned}$$

where in (*) we use the fact that the slope of the segment $(k-1, \ell_{k-1})(k, \ell_k)$ is greater than $-t$. Hence (k, d'_k) lies above (k, ℓ'_k) . Now suppose $(k-1, d_{k-1})$ lies on a corner of L . Then $d_{k-1} = \ell_{k-1}$ so

$$d_k + t \geq \ell_k + t > d_{k-1} = \ell_{k-1} = \ell'_k,$$

showing $d'_k = \ell'_k$ and (k, d'_k) lies on N' . □

7 Places as Galois orbits

Here is an alternate definition of a place.

Definition 7.7.1: Let (K, v) be a field with valuation and L/K be an extension. A **place** on L over v is a $G(\overline{K}_v/K_v)$ -orbit on $\text{Hom}_K(L, \overline{K}_v)$.

Example 7.7.2: Let $K = \mathbb{R}$, and L a finite extension of K . Then the places of L over \mathbb{R} are just $\text{Hom}_K(L, \mathbb{R})$, the real embeddings of L , and the complex places are just $G(\mathbb{C}/\mathbb{R}) \backslash \text{Hom}_K(L, \mathbb{C})$, i.e. pairs of complex conjugate embeddings.

We show this is equivalent to our previous definition.

Theorem 7.7.3. *Let K be a number field. There is a bijective correspondence between equivalence classes of valuations $w \mid v$, v on K , and $G(\overline{K}_v/K_v)$ -orbits on $\text{Hom}_K(L, \overline{K}_v)$:*

$$\{w \mid v : w \in M_L\} \xrightarrow{\cong} G(\overline{K}_v/K_v) \backslash \text{Hom}_K(L, \overline{K}_v).$$

Letting \bar{v} be the unique extension of v to \overline{K}_v , the embedding $\tau : L \hookrightarrow \overline{K}_v$ is associated to the valuation $|\cdot|_{\bar{v}}$ restricted to L .

Proof. □

8 Krasner's lemma and consequences

The following shows a surprising difference between local and global fields.

Lemma 7.8.1 (Krasner's lemma). *Let K be complete with respect to a nonarchimedean valuation $|\cdot|$, and extend $|\cdot|$ to an algebraic closure K^{al} . Let $\alpha, \beta \in K^{\text{al}}$. If β is separable over $K(\alpha)$, and*

$$|\beta - \alpha| < |\beta' - \beta| \tag{7.4}$$

for any conjugate $\beta' \neq \beta$ of β over K , then $\beta \in K(\alpha)$.

We say that α belongs to β if inequality (7.4) holds.

Proof. By the fixed field theorem, it suffices to show that for all embeddings $\sigma : K(\alpha, \beta) \hookrightarrow K^{\text{al}}$ fixing $K(\alpha)$, that $\sigma(\beta) = \beta$. We have

$$|\sigma(\beta) - \alpha| = |\sigma(\beta) - \sigma(\alpha)| = |\beta - \alpha|$$

since $|\bullet| = |\sigma \bullet|$ and $\sigma(\alpha) = \alpha$. Hence

$$|\sigma(\beta) - \beta| = |(\sigma(\beta) - \alpha) + (\alpha - \beta)| \leq |\beta - \alpha|,$$

the last following since $|\cdot|$ is nonarchimedean. By the minimality assumption we must have $\sigma(\beta) = \beta$. \square

We define a norm on polynomials by setting

$$\left\| \sum_{k=0}^n c_k X^k \right\| = \max_{0 \leq k \leq n} |c_k|.$$

Using Krasner's Lemma, we show that polynomials that are close together have roots that are closely related. In fact, we have the following stronger result. Using Krasner's Lemma, we show that polynomials that are close together have roots generating the same extensions.

Theorem 7.8.2. *Given f , there exists $\varepsilon > 0$ such that if $\|f - g\| < \varepsilon$, then there is an ordering of roots $\alpha_1, \dots, \alpha_n$ and β_1, \dots, β_n of f and g , respectively, counting multiplicities, such that $K(\alpha_j) = K(\beta_j)$.*

Proof. Step 1: First we show that the roots of g approach the roots of f , as $\|f - g\| \rightarrow 0$.

Lemma 7.8.3. *Keep the hypothesis of the theorem. Suppose $\varepsilon > 0$. Then there exists $\delta > 0$ such that if $\|f - g\| < \delta$, then for every root β of g , there exists a root α of f such that $|\beta - \alpha| < \varepsilon$.*

Proof. First note that the roots of a monic polynomial h are bounded in terms of $\|h\|$. Indeed, letting $h(X) = \sum_{k=0}^n c_k X^k$, if γ is a root of h , then by Proposition 7.2.7(3), we must have $c_k \gamma^k \geq \gamma^n$ for some $0 \leq k < n$, and hence

$$\gamma \leq c_k^{\frac{1}{n-k}} \leq \max(1, \|h\|).$$

Suppose $\|f - g\| \leq \delta$ is small (say, less than 1). Then $\|g\| \leq \|f\| + \delta$, which is bounded. Hence the roots of $\|g\|$ are bounded, say by C . Let β be a root of g . On the one hand, we have

$$(f - g)(\beta) \leq \|f - g\| \max\{|\beta|^n, 1\} \leq \delta \max\{C^n, 1\} \tag{7.5}$$

and on the other,

$$(f - g)(\beta) = f(\beta) = \prod_{k=1}^n (\beta - \alpha_k).$$

Hence $|\beta - \alpha_k| \leq (\delta \max\{C^n, 1\})^{\frac{1}{n}}$ for some n . We can choose δ so this is less than ε . \square

Step 2: We strengthen the lemma to account for multiplicities.

Lemma 7.8.4. *Keep the hypotheses of the theorem. For every $\varepsilon > 0$ there exists $\delta > 0$ such that whenever $\|f - g\| < \delta$, there exist orderings $\alpha_1, \dots, \alpha_n$ and β_1, \dots, β_n such that $|\beta_k - \alpha_k| < \varepsilon$ for all k .*

Proof. By Lemma 7.8.3, as $\|f - g\| \rightarrow 0$, the distance from the roots of g to the closest roots of f approaches 0. Let $\beta_1(g), \dots, \beta_n(g)$ be the roots of g . For each k let $\alpha_k(g)$ be the root of f closest to $\beta_k(g)$. We have $\max_k |\beta_k(g) - \alpha_k(g)| \rightarrow 0$ as $g \rightarrow f$. Suppose the distinct roots $\alpha'_1, \dots, \alpha'_m$ of f have multiplicities r_1, \dots, r_m , and suppose that they occur with multiplicities s_1, \dots, s_m in the $\alpha_k(g)$. Suppose by way of contradiction that $(s_1(g), \dots, s_m(g))$ is not constantly (r_1, \dots, r_m) for g close enough to f . Then we can find a sequence $g_j \rightarrow f$ such that $(s_1(g_j), \dots, s_m(g_j))$ is constant and not equal to (r_1, \dots, r_m) . Then

$$g_j(X) = \prod_{k=1}^n (X - \beta_k(g_j)) \rightarrow \prod_{k=1}^m (X - \alpha'_k(g_j))^{s_k} \neq \prod_{k=1}^m (X - \alpha'_k)^{r_k} = f(X),$$

contradiction. \square

Step 3: Take $\varepsilon = \min_{i \neq j} |\alpha'_i - \alpha'_j|$ in Lemma 7.8.4. Then Krasner's Lemma 7.8.1 gives the conclusion. \square

From this we get that every field extension of \mathbb{Q}_p can be described by a field extension of \mathbb{Q} , by choosing a close enough approximation to a minimal polynomial.

Corollary 7.8.5. *Let L/\mathbb{Q}_p be a finite extension. Then there is a finite extension K/\mathbb{Q} such that $[K : \mathbb{Q}] = [L : \mathbb{Q}_p] = n$ and $K \cdot \mathbb{Q}_p = L$.*

Proof. Using the primitive element theorem, choose α so that $\mathbb{Q}_p(\alpha) = L$. Let $f \in \mathbb{Q}_p[X]$ be the minimal polynomial of α . By Theorem 7.8.2, for g close enough to f , there is a root β of g such that $\mathbb{Q}_p(\alpha) = \mathbb{Q}_p(\beta)$. Take $g \in \mathbb{Q}[X]$ sufficiently close, and $L = K(\beta)$. Then

$$K \cdot \mathbb{Q}_p = K(\alpha) = K(\beta) = L. \tag{7.6}$$

Chapter 8

Local and global fields

1 Topology of local fields

Definition 8.1.1: A local field is a field K with a nontrivial valuation $|\cdot|$ such that K is locally compact.

Note this requires that K is complete.

Proposition 8.1.2: Let K be complete with respect to a discrete nonarchimedean valuation. (Then we can define $A := \mathcal{O}_K = \{x \in K : |x| \geq 0\}$ and $\mathfrak{m} = \{x \in K : |x| > 0\}$.) Then A is compact if and only if $k := A/\mathfrak{m}$ is finite.

Proof. Suppose A is compact. Note $\mathfrak{m} = \{x : |x| < 1\}$ is open, and any translate of it is open. Note $A = \bigsqcup_{a \in A/\mathfrak{m}} a + \mathfrak{m}$ where the union is over representatives in A/\mathfrak{m} . A finite number of these cover A , so k is finite.

Conversely, suppose $k := A/\mathfrak{m}$ is finite. It suffices to show that A is closed and totally bounded¹.

1. A is closed since $A = \{x : |x| \leq |\pi|\}$.
2. A is totally bounded: Given $\varepsilon > 0$, choose r so that $|\pi|^{r+1} < \varepsilon$. Now every element is in a ball of radius 1 centered at one of the finite number of points in the form $a_0 + a_1\pi + \cdots + a_r\pi^r$. □

Proposition 8.1.3: If K has finite residue field then \mathcal{O}_K^\times , \mathfrak{p}^n , and $1 + \mathfrak{p}^n$ are all compact.

Proof. From Proposition 8.1.2, A is compact. The above are all closed subsets of A so compact. □

Theorem 8.1.4. *The following is a complete classification of local fields, up to isomorphism.*

¹A set is *totally bounded* if for every r , A can be covered by a finite number of sets with diameter at most r .

1. \mathbb{R} and \mathbb{C} with the usual metric.
2. Finite extensions of \mathbb{Q}_p .
3. Field of formal Laurent series $k((T))$ over finite field.

All local fields are either \mathbb{R} , \mathbb{C} , or are complete with respect to a discrete nonarchidean valuation.

Proof. [Neu99], p. 135. □

1.1 Open sets and continuity

Proposition 8.1.5: For any local field K and any n , the n th power map is open on K^\times , i.e. it takes open subsets of K^\times to open sets.

Proof. For $K = \mathbb{R}$ or \mathbb{C} , this is clear.

For K π -adic, this is an easy consequence of Hensel's Lemma. Let $y \in K^{\times n}$. We may suppose $v(y) = 0$. Suppose $x_0^n - y = 0$. Let $k = v(\pi)$ and let ε be such that $v(\varepsilon - y) \geq 2k + 1$. Consider the polynomial $f(x) = x^n - y$. Now $f(x_0) \equiv 0 \pmod{\pi^{2k+1}}$ so by Hensel's Lemma x_0 lifts to a solution of f in K . □

Proposition 8.1.6: For any extension of local fields L/K , any $\sigma \in G(L/K)$ acts as a homeomorphism, and the norm map $\text{Nm}_{L/K}$ is continuous on K^\times .

Proof. In the case \mathbb{C}/\mathbb{R} , we can verify this directly. Otherwise, L is complete with respect to a discrete nonarchimedean valuation. We show that σ preserves distance. Because σ is a field automorphism it suffices to show $|\sigma(x)| = |x|$. Let π be a uniformizer. For $\sigma \in G(L/K)$, $\sigma\mathfrak{m} = \mathfrak{m}$, so it must take the uniformizer to another uniformizer, $\sigma(\pi) = \pi' = a_1\pi + a_2\pi^2 + \dots$ with $a_1 \neq 0$. Then $\sigma(a_j\pi^j + a_{j+1}\pi^{j+1} + \dots) = a_j\pi'^j + a_{j+1}\pi'^{j+1}$ have the same valuation, as needed.

$\text{Nm}_{L/K}(x) = \prod_{\sigma \in G(L/K)} \sigma(x)$ by Proposition 1.2.3. σ and multiplication are continuous, so $\text{Nm}_{L/K}$ is continuous. □

2 Unramified extensions

Definition 8.2.1: Let K be a complete field with residue field k ; let L be a finite extension of K with residue field l . We say L/K is **unramified** if l/k is separable and the prime ideal \mathfrak{p} in \mathcal{O}_K does not ramify in L .

L/K is **totally ramified** if \mathfrak{p} ramifies completely; by the degree equation this is equivalent to $l = k$.

Note from the residue equation that

$$\mathfrak{p} \text{ does not ramify} \iff [L : K] = [l : k]. \tag{8.1}$$

Our main theorem of this section is Theorem 8.2.4. We will show that if L/K is unramified, then l/k is separable. If l/k is separable, though, we need an extra condition to make sure L/K is unramified; namely that a minimal polynomial for L/K stays a minimal polynomial for l/k , so that (8.1) holds.

Proposition 8.2.2: Let K be a complete field with residue field k ; let L be a finite extension of K with residue field l . Suppose $\alpha \in \mathcal{O}_K$, $L = K(\alpha)$, and let $g(x) \in K[x]$. The following are equivalent.

1. L/K is unramified, and g is the minimal polynomial of α .
2. l/k is separable, with $l = k(\bar{\alpha})$, g has α as a root, \bar{g} is the minimal polynomial of $\bar{\alpha}$, and \bar{g} has no repeated roots.

Proof. Suppose (1) holds. Then \bar{g} has $\bar{\alpha}$ as a root. Note $L = K(\alpha)$ gives $l = k(\bar{\alpha})$. By (8.1), $\bar{\alpha}$ has degree $[l : k] = [L : K]$ over k . Since \bar{g} has degree $[L : K]$, it must be the minimal polynomial of $\bar{\alpha}$, and have no repeated roots. This shows l/k is separable.

Suppose (2) holds. We have


$$[L : K] = \deg g = \deg \bar{g} = [l : k],$$

the first equality is because α is a root of g and the fact that \bar{g} is irreducible means g is irreducible (hence the minimal polynomial of α); and the last equality is because \bar{g} is the minimal polynomial of $\bar{\alpha}$. Hence \mathfrak{p} (the prime ideal of \mathcal{O}_K) is unramified by (8.1). Thus L/K is unramified. \square

For local fields, the property of being unramified behaves well under extensions and products.

Proposition 8.2.3:

1. Suppose that $K \subseteq L \subseteq M$ are finite extensions. If M/L and L/K are unramified, then M/K is unramified.
2. Suppose that $K \subseteq L, M$ are finite extensions. If L/K is unramified, then LM/M is unramified.
3. Suppose that $K \subseteq L, M$ are finite extensions. If L/K and M/K are unramified, then LM/K is unramified.



Proof. Let k, l, m, n be the residue fields of K, L, M, LM , and $\mathfrak{p}, \mathfrak{P}$, and \mathfrak{P}' be the prime ideals of $\mathcal{O}_K, \mathcal{O}_L, \mathcal{O}_M$, respectively.

1. We have $\mathfrak{p}\mathcal{O}_M = \mathfrak{P}\mathcal{O}_L = \mathfrak{P}'$. Separability is transitive, so M/K is unramified.
2. Write $L = K(\alpha)$. By Proposition 8.2.2, we can find g with α as root such that \bar{g} is the minimal polynomial of $l = k(\alpha)$ over k , and is separable. Then the minimal polynomial for $n = m(\alpha)$ over m divides \bar{g} , hence is separable. By Proposition 8.2.2 again, LM/M is unramified.
3. By part 2, LM/M is unramified. Since M/K is unramified, by part 1 LM/K is unramified. \square

Theorem 8.2.4. *Let K be a field; fix an algebraic closure. There is an equivalence of categories between*

- finite unramified extensions L/K , and
- finite separable extensions l/k ,

where the morphisms are field inclusions. (All maps between fields are inclusions.) The equivalence is given by $L \mapsto \mathcal{O}_L/\mathfrak{P}$ where \mathfrak{P} is the prime in \mathcal{O}_L .

Moreover,

1. $L \subseteq M$ if and only if $l \subseteq m$.²
2. The residue field of LM is lm .
3. L/K is Galois if and only if l/k is Galois, and

$$G(L/K) \xrightarrow{\cong} G(l/k)$$

by restricting $\sigma \in G(L/K)$ to $B = \mathcal{O}_L$ and modding out by $\mathfrak{P}B$.

Proof. By Proposition 8.2.2, L does get sent to a separable extension.

First we show the map is surjective. Given l/k separable, choose β so that $l = k(\beta)$ and choose f so that \bar{f} be the minimal polynomial of β . Since β is a simple root of \bar{f} , by Hensel's Lemma 7.5.1 we can lift it to a root α of f . Then $K(\alpha)$ is mapped to $k(\beta)$.

Part (2) is clear. For (1), if $L \subseteq M$ then clearly $l \subseteq m$. Conversely, suppose $l \subseteq m$. Now LM is also unramified (Proposition 8.2.3) and has residue field $l \cdot m = m$. Hence,

$$[M : K] = [m : k] = [lm : k] = [LM : K],$$

showing $L \subseteq M$.

If $l = m$, then the above shows that $L = M$. Hence the map is injective.

²This is part of the statement of equivalence of categories.

For (3), note an extension is Galois iff it is the (minimal) splitting field of a separable polynomial f . Take g to be the minimal polynomial of a primitive element α ; note $\bar{\alpha}$ generates l/k . Note by Proposition 8.2.2, \bar{g} is separable. If L/K is Galois, then g splits over L so \bar{g} splits over l . Combining the previous two statements, l/k is Galois. Conversely, suppose l/k is Galois. Since \bar{g} splits into nonrepeated linear factors, Hensel's Lemma 7.5.2 lifts it to a factorization of g . Hence g splits over K into distinct linear factors, showing L/K is Galois. \square

Suppose k is a finite field. In this case, the separable extensions l/k are exactly the finite extensions. Moreover, we understand what these extensions are; there is one of each degree, and we can find the corresponding L/K explicitly. Furthermore, by surjectivity in (3), $G(L/K)$ contains a unique element mapping to the Frobenius element in $G(l/k)$; see Definition 11.1.1.

Lemma 8.2.5. *Let α be a root of*

$$f(X) := X^n - a = 0$$

where a is a unit and $p \nmid n$. Then $K(\alpha)/K$ is unramified.

Proof. Let $g(X) \mid f(X)$ be the minimal polynomial of α . Let $L = K(\alpha)$ and l be its residue field.

Note that $f'(X) = nX^{n-1} \neq 0$ has no common factor with $f(X) = X^n - a$, even when reduced modulo \mathfrak{p} , as $p \nmid n$ and $a \notin \mathfrak{p}$. Hence $f(X)$, and *a fortiori* $g(X)$, has no repeated root in k . Any factorization of $g(X)$ in k gives a factorization of $g(X)$ in K by Hensel's Lemma. Hence g remains irreducible in $k[X]$. This shows $[l : k] = [L : K]$. By the degree equation, L/K must be unramified. \square

Theorem 8.2.6. *Let L/K be an extension of complete fields with finite residue fields. Then there exists a field $K \subseteq L_u \subseteq L$ such that L_u/K is unramified and every unramified extension of K contained in L is contained in L_u . Moreover,*

1. L_u is obtained by adjoining to K all roots of unity in L whose order is relatively prime to $q := \text{char}(K)$.
2. L/L_u is totally ramified.

$$\begin{array}{c} L \\ \left| \text{totally ramified} \right. \\ L_u \\ \left| \text{unramified} \right. \\ K \end{array}$$

We call L_u the **maximal unramified extension** of K contained in L .

Proof. Let L_u be the compositum of all unramified extensions of K contained in L . Then L_u is unramified by Proposition 8.2.3, and it contains all unramified extensions of K contained in L .

For each n not a multiple of p , $K(\zeta_n)/K$ is unramified by Lemma 8.2.5. Letting $q = |k|$, the corresponding extension of residue fields is $k(\zeta_n)/k = \mathbb{F}_{q^{\text{ord}_q(n)}}/\mathbb{F}_q$. We get all finite extensions l/k in this way, thus all unramified extensions L'/K in this way. Taking the roots of unity inside L gives the result. \square

3 Ramified extensions

Definition 8.3.1: Let L/K be a ramified extension of local fields, with $q := \text{char}(k) = p^n$. We say

1. L/K is **tamely ramified** if $p \nmid [l : k]$.
2. L/K is **wildly ramified** if $p \mid [l : k]$.

We seek analogues of Lemma 8.2.5 in for ramified extensions.

For a prime \mathfrak{p} of a Dedekind domain A (not necessarily corresponding to a local field) let $v_{\mathfrak{p}}$ denote the corresponding valuation. (That is, if $v_{\mathfrak{p}}(a)$ is defined such that $\mathfrak{p}^{v_{\mathfrak{p}}(a)}$ is the highest power of \mathfrak{p} dividing (a) .) Note the following two facts.

1. If $\mathfrak{p}B = \mathfrak{P}^e$, then

$$v_{\mathfrak{p}}(a) = v_{\mathfrak{P}}(a)^e.$$

2. If $a_1 + \cdots + a_n = 0$, then the minimum value of $v_{\mathfrak{p}}(a_i)$ is attained for two indices.

Definition 8.3.2: An **Eisenstein extension** relative to \mathfrak{p} is an extension $K(\alpha)/K$ where the minimal polynomial of α is of the form

$$f(x) = x^n + a_{n-1}x^{n-1} + \cdots + a_0$$

where $v_{\mathfrak{p}}a_i > 0$ and $v_{\mathfrak{p}}a_0 = 1$.

Theorem 8.3.3. *The prime ideal \mathfrak{p} totally ramifies in any Eisenstein extension relative to \mathfrak{p} :*

$$\mathfrak{p}B = \mathfrak{P}^e, \quad \mathfrak{P} = (f(\alpha), \mathfrak{P})^e.$$

Proof. Let $\mathfrak{P}^e \parallel \mathfrak{p}$. Note $e \leq n = [L : K]$. We calculate the valuation of $f(\alpha)$ with respect to \mathfrak{P} .

$$\begin{aligned} v_{\mathfrak{P}}(\alpha^n) &= nv_{\mathfrak{P}}(\alpha) \\ v_{\mathfrak{P}}(a_k \alpha^k) &= e + kv_{\mathfrak{P}}(\alpha) > e, & 1 \leq k \leq n-1 \\ v_{\mathfrak{P}}(a_0) &= e. \end{aligned}$$

Since $f(\alpha) = \alpha^n + \cdots + a_0 = 0$, the minimum valuation must be attained for two terms. The only way this is possible is if $nv_{\mathfrak{P}}(\alpha) = e$. Then $n = e$ and $v_{\mathfrak{P}}(\alpha) = 1$, as needed. \square

Theorem 8.3.4. *Let K be complete with respect to a nonarchimedean valuation. The totally ramified extensions of K are exactly those of the form $K(\alpha)$ where α is the root of an Eisenstein polynomial.*

Proof. The forward direction follows directly from Theorem 8.3.3.

Conversely, let L/K be a totally ramified extension. Take α to be a generator of the maximal ideal \mathfrak{P} of \mathcal{O}_L . Note $v(\alpha) = \frac{1}{n}$ since $(\alpha)^n = \mathfrak{p}$. Note that for any a_{n-1}, \dots, a_0 , we have

$$v(a_k \alpha^k) = v(a_k) + \frac{k}{n} \equiv \frac{k}{n} \pmod{1},$$

since $v(a_k)$ is an integer. Thus, the nonzero terms $a_k \alpha^k$, $0 \leq k < n$, have different orders. Thus by Proposition 7.2.7, $a_{n-1} \alpha^{n-1} + \dots + a_0 \neq 0$ unless all coefficients are 0. This shows that α must have degree n ; suppose $\alpha^n + a_{n-1} \alpha^{n-1} + \dots + a_0 = 0$. Again by Proposition 7.2.7, the minimum order is attained for two terms. We have

$$\begin{aligned} v_{\mathfrak{p}}(\alpha^n) &= n v_{\mathfrak{p}}(\alpha) = 1 \\ v_{\mathfrak{p}}(a_k \alpha^k) &= k v(a_k) + \frac{k}{n}, \quad 0 \leq k \leq n-1. \end{aligned}$$

The only way this can happen is if α^n and $v(a_0)$ are the nonzero terms with least order. This gives $v(a_0) = 1$, and $v(a_k) > 0$ for $1 \leq k \leq n$, i.e. the polynomial is Eisenstein. \square

Theorem 8.3.5. *Suppose L/K is a totally and tamely ramified extension of degree n . Then $L = K(\alpha)$ for some α a root of*

$$X^n - \pi = 0$$

for $\pi \in \mathfrak{p}$.

Proof. Take $\beta \in \mathfrak{P}$. Since L/K is totally ramified, $v_{\mathfrak{p}}(\beta^n) = 1$. Hence $\beta^n = u\pi$ for some $u \in B^\times$, and β is a zero of

$$g(X) := X^n - u\pi.$$

Unfortunately, u may not be in A . However, we show that this polynomial is close enough to

$$f(X) := X^n - u'\pi$$

for some $u' \in A$ and proceed as in Theorem 7.8.2 to show that the roots of these two polynomials generate the same extension.

Since L/K is totally ramified, $l = k$, i.e. $A/\mathfrak{p}A \xrightarrow{\cong} B/\mathfrak{P}B$. Thus there exists $u' \equiv u \pmod{\mathfrak{P}}$ with $u' \in A$. This means $|u' - u| < 1$. Letting $\alpha_1, \dots, \alpha_n$ be the roots of $f(X) = 0$,

$$|\beta - \alpha_1| \cdots |\beta - \alpha_n| = |f(\beta)| = |u\pi - u'\pi| < |\pi| = |\alpha_1| \cdots |\alpha_n|$$

so $|\beta - \alpha_j| < |\alpha_j|$ for some j ; without loss of generality $j = 1$.

Since L/K is tamely ramified, $p \nmid n$ and $f'(\alpha_1) = n\alpha_1^{n-1}$ has valuation $|\alpha_1|^{n-1}$. Hence

$$|\alpha_1|^{n-1} = |f'(\alpha_1)| = |(\alpha_1 - \alpha_2) \cdots (\alpha_1 - \alpha_n)|. \quad (8.2)$$

Note $|\alpha_j| = |u'\pi|^{\frac{1}{n}} = |\alpha_1|$; hence $|\alpha_1 - \alpha_j| \leq |\alpha_1|$. By (8.2), equality must hold. Hence $|\beta - \alpha_1| < |\alpha_1 - \alpha_j|$ for all $j \neq 1$, and by Krasner's Lemma 7.8.1, $K(\alpha_1) \subseteq K(\beta)$. Since both extensions are totally ramified of degree n , $L = K(\beta) = K(\alpha_1)$. \square

The analogues of Proposition 8.2.3 carry over exactly.

Proposition 8.3.6:

1. Suppose that $K \subseteq L \subseteq M$ are finite extensions. If M/L and L/K are tamely ramified, then M/K is tamely ramified.
2. Suppose that $K \subseteq L, M$ are finite extensions. If L/K is tamely ramified, then LM/M is tamely ramified.
3. Suppose that $K \subseteq L, M$ are finite extensions. If L/K and M/K are tamely ramified, then LM/K is tamely ramified.

Theorem 8.3.7. *Let K be a field with characteristic 0 and finite residue field, and let \mathfrak{p} be a prime in \mathcal{O}_K . Given n , there are only finitely many extensions of $K_{\mathfrak{p}}$ with degree at most n .*

Proof. First we show that there are finitely many totally ramified extensions of degree n . By Theorem 8.3.4, very such extension is realized by adjoining a root of an Eisenstein polynomial of degree n . By taking the coefficients, an Eisenstein polynomial can be identified with a point of

$$\underbrace{\mathfrak{p} \times \cdots \times \mathfrak{p}}_{n-1} \times A^\times \pi. \quad (8.3)$$

The topology given by $\|\cdot\|$ is exactly the product topology here; this is compact by Proposition 8.1.3. Now for each polynomial f , by Theorem 7.8.2 there exists an open set U_f such that any $g \in U_f$ has roots generating the same extensions as those of f . Since (8.3) is compact, a finite number of U_f cover f . The roots corresponding to those f generate all the totally ramified extensions of degree n .

By Theorem 8.2.6. Any finite extension L of degree n is an totally ramified extension of degree $\frac{n}{m}$ of an unramified extension L_u of degree m for some m . By the remark after Theorem 8.2.4, there is exactly one unramified extension of degree m ; for each L_u by the above there are a finite number of possibilities for L . \square

4 Witt vectors*

We know from Proposition 7.4.6 that every element of \hat{K} can be written as $\sum_{n \geq N} a_n \pi^n$ where the a_n come from a fixed set of representatives for A/\mathfrak{m} . Although this allows us to write down any element, unless the set of representatives is closed under addition and multiplication (i.e. form a copy of k in A), we cannot simply add and multiply the coefficients. Instead, we find

that addition and multiplication are governed by *Witt vectors*. We will actually develop this theory in a more general context.

Definition 8.4.1: Let p be a prime number. A ring R is a **strict p -ring** if R is complete and Hausdorff with respect to the p -adic topology, p is not a zero-divisor in R , and the residue ring $R/(p)$ is perfect. (A ring of characteristic p is **perfect** if the map $x \mapsto x^p$ is bijective.)

We will primarily be interested in the case where R is an unramified extension of \mathbb{Z}_p .

Theorem 8.4.2. *Let K be a perfect ring of characteristic p .*

1. *There is a strict p -ring R with residue ring K , unique up to canonical isomorphism.*
2. *There is a unique system of representatives $\tau : K \rightarrow R$, called the **Teichmüller representatives**, such that*

$$\tau(xy) = \tau(x)\tau(y)$$

for all $x, y \in K$.

The main example of interest to us is the following.

Example 8.4.3: Fix f ; then there is a unique unramified extension of \mathbb{Z}_p with residue field \mathbb{F}_q , $q = p^f$, namely $\mathbb{Z}_p[\zeta_{p^f-1}]$. The Teichmüller representatives are the $(q-1)$ th roots of unity μ_{q-1} . They are multiplicative, but not additive. The following construction will tell us how to add them.

Lemma 8.4.4. *Given $X = (X_0, X_1, \dots)$, define*

$$W_n(X) = X_0^{p^n} + pX_1^{p^{n-1}} + \dots + p^n X_n, \quad n \geq 0.$$

Then there exist polynomials

$$S_0, S_1, \dots; P_0, P_1, \dots \in \mathbb{Z}[X_0, X_1, \dots, Y_0, Y_1, \dots]$$

such that

$$\begin{aligned} W_n(S) &= W_n(X) + W_n(Y) \\ W_n(P) &= W_n(X) \cdot W_n(Y). \end{aligned}$$

where $X = (X_0, X_1, \dots)$, $Y = (Y_0, Y_1, \dots)$, $S = (S_0, S_1, \dots)$, and $P = (P_0, P_1, \dots)$.

The motivation for defining these polynomials is that they tell us how to add in strict p -rings using the base- p representation with Teichmüller representatives as coefficients.

Theorem 8.4.5. *Let R be a strict p -ring, k its residue ring, and $\tau : k \rightarrow R$ be the system of Teichmüller representatives. Then*

$$\sum_{n=0}^{\infty} \tau(x_n)p^n + \sum_{n=0}^{\infty} \tau(y_n)p^n = \sum_{n=0}^{\infty} \tau(S_n(x_0^{p^{-n}}, x_1^{p^{-(n-1)}}, \dots, x_n; y_0^{p^{-n}}, y_1^{p^{-(n-1)}}, \dots, y_n)p^n).$$

Proof of Lemma 8.4.4. We will abbreviate

$$\begin{aligned} W(X) &= (W_0(X), W_1(X), \dots) \\ R &= \mathbb{Z}[X_0, X_1, \dots; Y_0, Y_1, \dots]. \end{aligned}$$

All comparisons between X, Y will be done componentwise, and we define $X^n = (X_0^n, X_1^n, \dots)$.

We find the S_m, P_m inductively, with the additional condition that S_m, P_m are polynomials in $X_0, \dots, X_m, Y_0, \dots, Y_m$. To begin, note $W_0(X) = X_0$ so we set

$$\begin{aligned} S_0(X, Y) &= X_0 + Y_0 \\ P_0(X, Y) &= X_0 Y_0. \end{aligned}$$

Lemma 8.4.6. *If $F_m, G_m \in R$ and $F_m \equiv G_m \pmod{p}$ for every m , then*

$$W_n(F) \equiv W_n(G) \pmod{p^{n+1}}.$$

Proof. First note that for any $f, g \in R$ such that $f \equiv g \pmod{p}$,

$$f^{p^j} \equiv g^{p^j} \pmod{p^{j+1}}.$$

The proof is by induction, with the induction step following by the binomial theorem: if $f^{p^{j-1}} = g^{p^{j-1}} + p^j h$ then

$$f^{p^j} = (g^{p^{j-1}} + p^j h)^p = g^{p^j} + \underbrace{\binom{p}{1} p^{j-1} h g^{p^{j-1}(p-1)}}_{p^j} + p^{j+1} k$$

for some $k \in R$.

This claim gives $f_j^{p^{n-j}} \equiv g_j^{p^{n-j}} \pmod{p^{n-j+1}}$ and hence

$$p^j f_j^{p^{n-j}} \equiv p^j g_j^{p^{n-j}} \pmod{p^{n+1}}.$$

Summing these up give the result. □

Directly from the definitions, we have

$$W_n(X) = W_{n-1}(X^p) + p^n X_n.$$

Hence the equations

$$\begin{aligned} W_n(S) &= W_n(X) + W_n(Y) \\ W_n(P) &= W_n(X)W_n(Y) \end{aligned}$$

are equivalent to

$$\begin{aligned} W_{n-1}(S^p) + p^n S_n &= W_{n-1}(X^p) + p^n X_n + W_{n-1}(Y^p) + p^n Y_n \\ &= W_{n-1}(S(X^p, Y^p)) + p^n (X_n + Y_n) \end{aligned} \quad (8.4)$$

$$\begin{aligned} W_{n-1}(P^p) + p^n P_n &= (W_{n-1}(X^p) + p^n X_n)(W_{n-1}(Y^p) + p^n Y_n) \\ &= W_{n-1}(P(X^p, Y^p)) + p^n (X_n W_{n-1}(Y^p) + Y_n W_{n-1}(X^p) + p^n X_n Y_n) \end{aligned} \quad (8.5)$$

where (8.4) and (8.5) follow from the hypothesis for $n - 1$. Solving for S_n and P_n , these are equivalent to

$$S_n = X_n + Y_n + \frac{W_{n-1}(S(X^p, Y^p)) - W_{n-1}(S^p)}{p^n}$$

$$P_n = X_n W_{n-1}(Y^p) + Y_n W_{n-1}(X^p) + p^n X_n Y_n + \frac{W_{n-1}(P(X^p, Y^p)) - W_{n-1}(P^p)}{p^n}.$$

However, since taking p th powers is a homomorphism modulo p , for any $f \in R$ we have $f(X, Y)^p \equiv f(X^p, Y^p) \pmod{p}$. Applying this to $f = S_j, P_j$, we see the conditions of the lemma are satisfied, so the numerators are divisible by p^n , and we can successfully define S_n and P_n . \square

Theorem 8.4.7. *Let A be a commutative ring. For*

$$a = (a_0, a_1, \dots), \quad b = (b_0, b_1, \dots), \quad a_i, b_i \in A_i,$$

the operations

$$a \overset{W}{+} b = S(a, b), \quad a \overset{W}{\cdot} b = P(a, b).$$

turn the set $A^{\mathbb{N}_0}$ into a commutative ring $W(A)$.

This is called the **ring of Witt vectors** over A .

Proof. We first prove that associativity, commutativity, and distributivity hold as polynomial identities in the a_j, b_j . The result then follows by considering the substitution homomorphism $\mathbb{Z}[a_0, \dots; b_0, \dots] \rightarrow A$.

Lemma 8.4.8. *The function $W : R^{\mathbb{N}} \rightarrow R^{\mathbb{N}}$, where $R := \mathbb{Z}[a_0, \dots; b_0, \dots]$, is injective.*

Proof. Suppose $X = (X_0, X_1, \dots)$ and $W(X) = (Y_0, Y_1, \dots)$. We show the X_j are determined by induction. We have $X_0 = W_0(X) = Y_0$. For the induction step, note

$$Y_n = W_n(X) = X_0^{p^n} + pX_1^{p^{n-1}} + \dots + p^n X_n;$$

since X_0, \dots, X_{n-1}, Y_n are determined and multiplication by p^n is injective in R , X_n is determined. \square

Lemma 8.4.4 gives

$$W(X \overset{W}{+} Y) = W(S(X, Y)) = W(X) + W(Y)$$

$$W(X \overset{W}{\cdot} Y) = W(P(X, Y)) = W(X) \cdot W(Y).$$

Hence $W : W(A) \rightarrow R^{\mathbb{N}}$ is a map that preserves addition and multiplication; moreover, it is injective. Its image is a subalgebra of R , since it contains 0 and 1:

$$W(0, 0, \dots) = (0, 0, \dots)$$

$$W(1, 0, \dots) = (1, 1, \dots).$$

Hence $\overset{W}{+}$ and $\overset{W}{\cdot}$ turn $W(A)$ into a commutative algebra with unit (we are basically “pulling back” the algebra structure from $R^{\mathbb{N}}$ to $W(A)$ using W). \square

4.1 Frobenius and Transfer maps

5 Extending valuations on global fields

Theorem 8.5.1. *Let $|\cdot|$ be a valuation on K and let \hat{K} be the completion of K with respect to $|\cdot|$. Let $L = K(\alpha)$ be a finite separable extension of K , and let f be the minimal polynomial of α .*

The completions of L with respect to the extensions $|\cdot|'$ of $|\cdot|$ are exactly $\hat{K}[X]/(h)$ as h ranges over irreducible factors of f in \hat{K} .

Proof. Suppose we are given an extension $|\cdot|'$. Let \hat{L} be the completion of L with respect to $|\cdot|'$.

$$\begin{array}{ccc} L = K(\alpha) & \hookrightarrow & \hat{L} = \hat{K}(\alpha) \\ \downarrow & & \downarrow \\ K & \longrightarrow & \hat{K} \end{array}$$

To see $\hat{L} = \hat{K}(\alpha)$, note we have $\hat{L} = \widehat{K(\alpha)} \supseteq \widehat{K}(\alpha)$; for the reverse inclusion note $\widehat{K}(\alpha)$ contains α and is complete (as it is a finite-dimensional vector space over a complete field).

Now considering the extension \hat{L}/\hat{K} , α is the root of one of the irreducible factors of f in $\hat{K}[X]$.

Conversely, given an irreducible factor h of f in $\hat{K}[X]$, consider $\hat{K}(\alpha') = \hat{K}[X]/\langle h \rangle$. The valuation on \hat{K} extends uniquely to $\hat{K}(\alpha')$ by Theorem 7.6.1. Then let $K(\alpha) \hookrightarrow \hat{K}(\alpha')$ be the map sending α to α' . (This makes sense as the minimal polynomials of α, α' over K are both f .) By the same argument as before, $\hat{L} = \widehat{K(\alpha)} = \hat{K}(\alpha')$, as desired.

$$\begin{array}{ccc} L = K(\alpha) & \hookrightarrow & \hat{L} = \hat{K}(\alpha') \\ \downarrow & & \downarrow \\ K & \longrightarrow & \hat{K} \end{array}$$

□

Theorem 8.5.2. *Let \hat{K} be the completion of K with respect to a archimedean or discrete nonarchimedean valuation $|\cdot|$. Let L/K be a finite separable extension. There are finitely many extensions of $|\cdot|$ to L ; denoting them by $|\cdot|_i$ and the respective completions of L be L_i , we have the natural isomorphism*

$$\hat{K} \otimes_K L \cong \prod_i L_i.$$

Proof. By the primitive element theorem, we can write $L = K(\alpha)$. Let f be the minimal polynomial of α . Let f factor into irreducibles in $\hat{K}[X]$ as

$$f = f_1 \cdots f_n.$$

Then

$$\hat{K} \otimes_K L \cong \hat{K} \otimes_K K[x]/(f) \cong \hat{K}[x]/(f) \stackrel{\text{CRT}}{\cong} \prod_{i=1}^n \hat{K}[x]/(f_i) \stackrel{\text{Thm 8.5.1}}{\cong} \prod_{i=1}^n L_i. \quad \square$$

Note the map in the theorem sends

$$a \otimes b \mapsto (a_1 b, \dots, a_n b),$$

where a_i is the embedding of a into L_i . We now have a way to calculate norms and traces in terms of completed fields.

Corollary 8.5.3. *Keep the same notation as above. Then*

1. $\text{Nm}_{L/K}(\alpha) = \prod_{i=1}^n \text{Nm}_{L_i/\hat{K}}(\alpha)$.
2. $\text{Tr}_{L/K}(\alpha) = \prod_{i=1}^n \text{Tr}_{L_i/\hat{K}}(\alpha)$.

Proof. Using Proposition 1.2.3(1) and Theorem 8.5.1, we see

$$\begin{aligned} \text{Nm}_{L/K}(\alpha) &= \prod_{\alpha' \text{ root of } f} \alpha' = \left(\prod_{\alpha' \text{ root of } f_1} \alpha' \right) \cdots \left(\prod_{\alpha' \text{ root of } f_n} \alpha' \right) = \prod_{i=1}^n \text{Nm}_{L_i/\hat{K}}(\alpha) \\ \text{Tr}_{L/K}(\alpha) &= \sum_{\alpha' \text{ root of } f} \alpha' = \left(\sum_{\alpha' \text{ root of } f_1} \alpha' \right) + \cdots + \left(\sum_{\alpha' \text{ root of } f_n} \alpha' \right) = \sum_{i=1}^n \text{Tr}_{L_i/\hat{K}}(\alpha). \end{aligned}$$

□

6 Product formula

Lemma 8.6.1. *Let L/K be a finite extension of number fields, with normalized nonarchimedean valuations $w \mid v$, as in Example 7.2.3. Let $|\cdot|'_w$ be w normalized so it extends v . Then*

$$|\cdot|_w = |\cdot|'_w{}^{[L_w:K_v]}.$$

Proof. We check equality on the completions. Let p be the characteristic of the residue fields, and $\pi_v \in K_v, \pi_w \in L_w$ be prime. Since $\|\cdot\|_w$ is a map $L^\times \rightarrow p^{\mathbb{Z}}$, it suffices to check that $\|\cdot\|_w$ and $\|\cdot\|'_w{}^{[L_w:K_v]}$ agree for one element that does not get sent to 1. We check them on π_v . Note that if \mathfrak{P} is the prime corresponding to w , $\mathfrak{N}\mathfrak{P} = |\mathcal{O}_L/\mathfrak{P}\mathcal{O}_L| = |\mathcal{O}_{L_w}/\pi_w\mathcal{O}_{L_w}| = |l|$ where $l = \mathcal{O}_{L_w}/\pi_w\mathcal{O}_{L_w}$ and similarly for K . Then

$$|\pi_v|_w = \left(\frac{1}{|l|} \right)^{e(\pi_w/\pi_v)} = \left(\frac{1}{|k|} \right)^{e(L_w/K_v)f(L_w/K_v)} = |\pi_v|'_w{}^{[L_w:K_v]}$$

by the degree equation 2.5.2. □

Theorem 8.6.2 (Product formula). *For any nonzero $\alpha \in K$,*

$$\prod_{v \in V_K} |\alpha|_v = 1.$$

Proof.

Step 1: We first show the result for $K = \mathbb{Q}$. Given $n \in \mathbb{Q}$, factor it as $n = \pm \prod_{i=1}^{\infty} p_i^{a_i}$ where p_i are all the prime numbers; note only a finite number of the a_i are nonzero. Then

$$|\alpha| = \left(\prod_{i=1}^{\infty} |\alpha|_{p_i} \right) |\alpha|_{\infty} = \left(\prod_{i=1}^{\infty} p_i^{-a_i} \right) \left(\prod_{i=1}^{\infty} p_i^{a_i} \right) = 1.$$

Step 2: We pass to field extensions of \mathbb{Q} using the following lemma.

Lemma 8.6.3 (Extension formula). *Let $K \subseteq L$ be number fields and let v be a place of K . Then*

$$\prod_{w|v} |\alpha|_w = \left| \text{Nm}_{L/K} \alpha \right|_v.$$

Proof. For a place on L let $|\cdot|'_w$ be the valuation normalized so that it extends v . We have

$$\begin{aligned} \left| \text{Nm}_{L/K} \alpha \right|_v &= \prod_{w|v} \left| \text{Nm}_{L_w/K_v}(\alpha) \right|_v \\ &= \prod_{w|v} \left| \text{Nm}_{L_w/K_v}(\alpha) \right|'_w \\ &= \prod_{w|v} \left| \text{Nm}_{L_w/K_v}(\alpha) \right|_w^{\frac{1}{[L:K]}} && \text{by Lemma 8.6.1} \\ &= \prod_{w|v} |\alpha|_w && \text{by Theorem 7.6} \square \end{aligned}$$

Step 3: Since every place on K restricts to a unique place on \mathbb{Q} ,

$$\prod_{w \in V_K} |\alpha|_w = \prod_{v \in V_{\mathbb{Q}}} \prod_{w|v} |\alpha|_w = \prod_{v \in V_{\mathbb{Q}}} \left| \text{Nm}_{L/K}(\alpha) \right|_v \stackrel{\text{Step 1}}{=} 1,$$

where we apply step 1 to $\text{Nm}_{L/K}(\alpha)$. □

The product formula will be useful when defining a measure of size independent of scaling (see Chapter ??).

7 Problems

1. Let K be a complete nonarchimedean field whose residue field has characteristic p . Prove that the maximal tamely ramified (separable) extension of K is

$$K_{\text{tr}} = K_u \left(\left\{ \pi^{\frac{1}{m}} : p \nmid m \right\} \right).$$

Chapter 9

Ramification

We seek to generalize the definition of discriminant over Dedekind domains A which are not PID's. To do this we will first define the *different*, which measure how much we can enlarge B so that the image of the trace map is still in A , then define the discriminant as the discrepancy between B and the enlarged B , using χ_A . We will find that the different is the (ideal) norm of the discriminant.

We will see that our definition coincides with our previous definition when A is a PID. Fortunately, we don't have to prove everything from scratch again: by localization we can always reduce to the DVR/PID case.

The main use of the discriminant is to measure ramification: The primes dividing the discriminant are those that ramify. On a deeper level, the exponents measure the degree of ramification.

1 Lattices and χ

Definition 9.1.1: Let A be a Dedekind domain, $K = \text{Frac}(A)$, and V a finite dimensional K -vector space. An A -submodule $X \subseteq V$ is a **lattice** if it is finitely generated A -module and $\text{span}_K(X) = V$.

The most basic example of a lattice is a fractional ideal of K .

We would like to measure the discrepancy between two lattices—like the norm, but measured by an *ideal* instead. To do this, we first need some facts from commutative algebra.

1.1 Filtrations of modules

Definition 9.1.2: A module is **simple** if it is nonzero and has no nonzero proper submodule. A composition series of length m is a chain of submodules

$$M = M_0 \supset M_1 \supset \cdots \supset M_m = 0$$

where M_{i-1}/M_i is simple for each i . M has **finite length** if it has a finite composition series.

Proposition 9.1.3: The simple modules are exactly those in the form R/\mathfrak{m} where \mathfrak{m} is a maximal ideal of R . If M is simple, $M = R/\mathfrak{m}$ where $\mathfrak{m} = \text{Ann}(M)$.

The main theorem on filtrations is the following.

Theorem 9.1.4 (Jordan-Hölder). *Suppose M has a composition series.*

1. (Existence) *Any chain of submodules of M can be refined to a composition series.*
2. (Uniqueness) *Any composition series of M has the same length; moreover the number of times R/\mathfrak{m} appears as a quotient M_{i-1}/M_i in the filtration is invariant.*

We will be applying this when R is a Dedekind domain, so the maximal ideals are simply the nonzero prime ideals.

We also need the following.

Proposition 9.1.5: If M/M' and M' have finite length, then so does M .

1.2 The function χ_A

Definition 9.1.6: Let A be a Dedekind domain. Define

$$\chi_A : \{A\text{-module of finite length}\} \rightarrow \{\text{ideals of } A\}$$

as follows: Given M of finite length, with composition series

$$M = M_0 \supset M_1 \supset \cdots \supset M_m = 0$$

and $A/\mathfrak{p}_i \cong M_{i-1}/M_i$, define

$$\chi_A(M) = \prod_{i=1}^m \mathfrak{p}_i.$$

Example 9.1.7: The primes appearing in the filtration of an ideal $\mathfrak{a} \subset A$ are just the primes dividing \mathfrak{a} with multiplicity, so

$$\chi_A(\mathfrak{a}) = (\mathfrak{a}).$$

Proposition 9.1.8: If M' and M'' have finite length and $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ exact sequence of A -modules, then

$$\chi_A(M) = \chi_A(M')\chi_A(M'').$$

Definition 9.1.9: Let A be a Dedekind domain, $K = \text{Frac}(A)$, and $X_1, X_2 \subseteq V$ be A -lattices. Choose $X_3 \subseteq X_1 \cap X_2$ any A -lattice and define

$$\chi_A(X_1, X_2) := \chi_A(X_1/X_3)\chi_A(X_2/X_3)^{-1}$$

as fractional ideals of K .

Proof of well-definedness. We show this is independent of choice of X_3 .

Observe $\chi_A(X_1, X_2)\chi_A(X_2, X_1) = (1)$. Note this is independent of choice of X_3 . It suffices to show that

$$\chi_A(X_1/X_3)\chi_A(X_2/X_3)^{-1} = \chi_A(X_1/X_4)\chi_A(X_2/X_4)^{-1}$$

when $X_4 \subseteq X_3$. This follows by the exact sequence

$$0 \rightarrow X_3/X_4 \rightarrow X_1/X_4 \rightarrow X_1/X_3.$$

and Proposition 9.1.8. □

1.3 χ and localization

It is easier to study χ_A when A is local; in this case $\chi_A(X)$ is simply a power of the maximal ideal. To understand χ_A (and hence the discriminant) for general A , we thus consider the localization of A at all primes. The following says that χ_A is well-behaved under localization.

Proposition 9.1.10: Let A be a Dedekind domain and $\mathfrak{p} \subset A$ be a nonzero prime. Then

$$v_{\mathfrak{p}}(\chi_A(\chi_1, \chi_2)) = v_{\mathfrak{p}A_{\mathfrak{p}}}(\chi_{A_{\mathfrak{p}}}((X_1)_{\mathfrak{p}}, (X_2)_{\mathfrak{p}})).$$

Proof. Note $X_{\mathfrak{p}} = A_{\mathfrak{p}} \cdot X = A_{\mathfrak{p}} \otimes_A X$ is an $A_{\mathfrak{p}}$ -lattice of V .

Localization is exact, so preserves quotients. Suppose $M \supseteq N$ are adjacent terms in the filtration of A . If $M/N = A/\mathfrak{p}$ then

$$M_{\mathfrak{p}}/N_{\mathfrak{p}} = (M/N)_{\mathfrak{p}} = (A/\mathfrak{p})_{\mathfrak{p}} = A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$$

while if $M/N = A/\mathfrak{q}$, $\mathfrak{q} \neq \mathfrak{p}$, then $M_{\mathfrak{p}}/N_{\mathfrak{p}} = 0$. Only the quotients with A/\mathfrak{p} remain; the result follows. □

Proposition 9.1.11: Let A be a Dedekind domain with fraction field K , X an A -lattice in V , and $\sigma \in \text{Aut}_K(V)$. Then

$$\chi_A(X, \sigma X) = (\det \sigma).$$

Proof. It suffices to check both sides have the same \mathfrak{p} -valuation for every prime \mathfrak{p} of A ; by Proposition 9.1.10 this is equivalent to

$$\chi_{A_{\mathfrak{p}}}(X_{\mathfrak{p}}, \sigma_{\mathfrak{p}}X_{\mathfrak{p}}) = (\det \sigma_{\mathfrak{p}}).$$

Thus we only need to check the proposition for the case where A is a DVR, hence a PID. For all nonzero $\alpha \in A$,

$$\chi(X, \alpha\sigma X) = \alpha^n \chi(X, \sigma X) = \det[\alpha] \cdot \chi(X, \sigma X);$$

note we used $\chi(uX, \alpha uX) = \alpha^n$ since X is free over A , and that the matrix of the transformation $[\alpha]$ is simply αI . Thus by choosing α such that $\alpha\sigma X \subseteq X$ we may assume $\sigma(X) \subseteq X$.

By the structure theorem for modules, $X/\sigma X \cong A/\alpha_1 \times \cdots \times A/\alpha_n$ for some α_j , giving

$$\chi_A(X, \sigma X) = (\alpha_1 \cdots \alpha_n) = (\det \sigma).$$

□

1.4 Discriminant of bilinear forms

In this section we will define the discriminant of a bilinear form T on a lattice X over K , the fraction field of a Dedekind domain A . When we specialize to the case that X is an extension of A and $T = \text{tr}$, then we get a generalization of our original definition 1.3.1, in the case where X is not necessarily free over A .

Definition 9.1.12: Keep the above assumptions. Let V be a finite-dimensional K -vector space and $T : (V, V) \rightarrow K$ be a nondegenerate K -bilinear form. Thinking of T as a map $V \otimes_K V \rightarrow K$, we get a map

$$\wedge^n T : \wedge^n V \otimes_K \wedge^n V \rightarrow K$$

defined by

$$\wedge^n T(v_1 \wedge \cdots \wedge v_n, w_1 \wedge \cdots \wedge w_n) = \sum_{\pi \in S_n} (-1)^{\text{sign}(\pi)} T(v_1 \wedge w_{\pi(1)}) \cdots T(v_n \wedge w_{\pi(n)}). \quad (9.1)$$

Note $\wedge^n T \otimes \wedge^n T$ is a 1-dimensional vector space over K , with lattice $\wedge^n X \otimes_K \wedge^n X$. Define the **discriminant** of T on X to be

$$\mathfrak{d}_{X,T} := \chi_A(\wedge^n T, \wedge^n X \otimes \wedge^n X).$$

The main reason for defining the discriminant as above is because the “ \wedge ” construction is natural and makes it easy to prove a few basic properties.

Proposition 9.1.13: If X is free over A with basis (e_1, \dots, e_n) , then

$$\mathfrak{d}_{X,T} = (\det(T(e_i, e_j))).$$

Proof. Note that $X \otimes_K X$ is generated by $\wedge^n T(e_1 \wedge \cdots \wedge e_n, e_1 \wedge \cdots \wedge e_n)$. By (9.1), this is exactly $(\det(T(e_i, e_j)))$. \square

We now give an alternative characterization of the discriminant, in terms of the *dual lattice*.

Definition 9.1.14: Define the **dual** of X with respect to T by

$$X_T^* := \{y \in V : T(x, y) \in A \text{ for all } x \in X\}.$$

This is an A -lattice of V .

We first need the following.

Proposition 9.1.15: If e_1, \dots, e_n is a basis for X over A , and e_1^*, \dots, e_n^* is a *dual basis*, i.e. $T(e_i, e_j^*) = \delta_{ij}$ for each j , then e_1^*, \dots, e_n^* is a basis for X^* over A .

Proof. Note $y \in X^*$ iff $T(e_j, y) \in A$ for each j . Writing $y = \sum_{j=1}^n a_j e_j^*$, we find $T(e_j, y) = a_j$, so $y \in X^*$ iff $a_j \in A$ for each j , i.e. $y \in \text{span}_A(e_1^*, \dots, e_n^*)$. \square

Proposition 9.1.16: We have

$$\chi_A(X_T^*, X) = \mathfrak{d}_{X,T}.$$

Proof. We use the fact that a fractional ideal is determined by its localizations at all primes (this follows since the exponent of \mathfrak{p} in \mathfrak{a} is the same as that of $\mathfrak{p}A_{\mathfrak{p}}$ in $\mathfrak{a}A_{\mathfrak{p}}$, Proposition 2.2.5).

By using Proposition 9.1.10, we may localize at nonzero $\mathfrak{p} \subset A$. Hence it suffices to prove may assume A is DVR, i.e. free over A .

Write

$$\begin{pmatrix} e_1 \\ \vdots \\ e_n \end{pmatrix} = B \begin{pmatrix} e_1^* \\ \vdots \\ e_n^* \end{pmatrix}$$

where $B = (b_{i,j})$ is a $n \times n$ matrix. Then by Proposition 9.1.13,

$$\mathfrak{d}_{X,T} = (\det(T(e_i, e_j))) = (\det(b_{i,j})) = \chi(X_T^*, BX_T^*) = \chi(X_T^*, X),$$

as needed. \square

2 Discriminant and different

For the AKLB setup with L/K finite separable, consider the nondegenerate K -bilinear map

$$\begin{aligned} \text{tr} : L \times L &\rightarrow K \\ (x, y) &\mapsto \text{tr}_{L/K}(xy). \end{aligned}$$

Definition 9.2.1: Define the **codifferent**

$$B^* := B_{\text{tr}}^* = \{y \in L : \text{tr}(xy) \in A \text{ for all } x \in B\}$$

and the **different** and **discriminant** by

$$\begin{aligned} \mathfrak{D}_{B/A} &= \mathfrak{D}_{L/K} := (B^*)^{-1} \\ \mathfrak{d}_{B/A} &= \mathfrak{d}_{L/K} := \mathfrak{d}_{B,\text{tr}}. \end{aligned}$$

(These are fractional ideals of K .)

Observe that $B \subseteq B^*$ (in light of $\text{tr}_{L/K}(B) \subseteq A$) so $B \supseteq \mathfrak{d}_{B/A}$.

The following gives the precise relationship between the discriminant and different.

Proposition 9.2.2: $\text{Nm}_{L/K}(\mathfrak{D}_{B/A}) = \mathfrak{d}_{B/A}$.

Proof. We have

$$\mathfrak{D}_{B/A} = \chi_A(B^*, B) = \chi_A(B^*/B)$$

and

$$\mathfrak{D}_{B/A} = (B^*)^{-1} = \chi_B(B^*/B).$$

The result thus follows from commutativity of the following diagram.

$$\begin{array}{ccc} \{\text{finite length } B\text{-module}\} & \xrightarrow{\chi_B} & I_B \\ & \searrow \chi_A & \downarrow \text{Nm}_{L/K} \\ & & I_K. \end{array}$$

We have commutativity since if B/\mathfrak{P} is a quotient of adjacent terms in the B -filtration of M , then when we refine it to a A -filtration, since $B/\mathfrak{P} = (A/\mathfrak{p})^{f(\mathfrak{P}/\mathfrak{p})}$ as vector spaces, we get $f(\mathfrak{P}/\mathfrak{p})$ copies of A/\mathfrak{p} . \square

2.1 Basic properties

First, a slightly cleaner characterization of the codifferent.

Lemma 9.2.3. $\mathfrak{a} \in I_K$ and $\mathfrak{b} \in I_L$. Then

$$\text{tr}_{L/K}(\mathfrak{b}) \subseteq \mathfrak{a} \iff \mathfrak{b} \subseteq \mathfrak{a}\mathfrak{D}_{B/A}^{-1}.$$

Proof. We check $\text{tr}(\mathfrak{a}^{-1}\mathfrak{b}) \subseteq A$ iff $\mathfrak{a}^{-1}\mathfrak{b} \subseteq \mathfrak{D}_{L/K}^{-1}$.

The reverse direction is clear. For the forward direction, note that if $x \in \mathfrak{a}^{-1}\mathfrak{b}$ and $y \in B$, then $xy \in \mathfrak{a}^{-1}\mathfrak{b}$ and hence $\text{tr}(xy) \in A$. This shows $x \in \mathfrak{D}_{B/A}^{-1}$. \square

Proposition 9.2.4:

- (Transitivity) Let M/L be a finite separable extension, with C the integral closure of A in M . Then

$$\mathfrak{D}_{C/A} = \mathfrak{D}_{C/B}\mathfrak{D}_{B/A}.$$

- (Localization) For $S \subseteq A$ a multiplicative subset,

$$S^{-1}\mathfrak{D}_{B/A} = \mathfrak{D}_{S^{-1}B/S^{-1}A}.$$

- (Completion)

$$\mathfrak{D}_{B/A} \cdot \hat{B}_{\mathfrak{P}} = \mathfrak{D}_{\hat{B}_{\mathfrak{P}}/\hat{A}_{\mathfrak{p}}}.$$

Proof. 1. We have

$$\begin{aligned}
 e &\in \mathfrak{D}_{C/B}^{-1} \mathfrak{D}_{B/A}^{-1} \\
 \iff \mathfrak{D}_{B/A} e &\subseteq \mathfrak{D}_{C/B}^{-1} \\
 \iff \mathrm{tr}_{M/L}(\mathfrak{D}_{B/A} e) &\subseteq B && \text{Lemma 9.2.3 with } M/L \\
 \iff \mathfrak{D}_{B/A} \mathrm{tr}_{M/L}(e) &\subseteq B \\
 \iff \mathrm{tr}_{M/L}(e) &\in \mathfrak{D}_{B/A}^{-1} \\
 \iff \mathrm{tr}_{L/K}(\mathrm{tr}_{M/L}(e)) &\subseteq A && \text{Lemma 9.2.3 with } L/K \\
 \iff e &\in \mathfrak{D}_{C/A}^{-1}.
 \end{aligned}$$

2. Omit.

3. Localize at \mathfrak{p} . May assume A is a DVR. (B may not be a DVR.) Consider

$$\begin{array}{ccc}
 \prod_{\mathfrak{P}|\mathfrak{p}} \hat{B}_{\mathfrak{P}} & \hookrightarrow & \prod_{\mathfrak{P}|\mathfrak{p}} \hat{L}_{\mathfrak{P}} \\
 \downarrow \cong & & \downarrow \cong \\
 B \otimes_A \hat{A}_{\mathfrak{p}} & \hookrightarrow & L \otimes_K \hat{K}_{\mathfrak{p}} \\
 \downarrow & & \downarrow \mathrm{tr}_{L/K} \otimes_K \hat{K}_{\mathfrak{p}} \\
 \hat{A}_{\mathfrak{p}} & \hookrightarrow & \hat{K}_{\mathfrak{p}}
 \end{array}$$

The top-to-bottom map on the right is $\sum \mathrm{tr}_{\hat{L}_{\mathfrak{P}}/\hat{K}_{\mathfrak{p}}}$. Then

$$\begin{aligned}
 \mathfrak{D}_{B/A}^{-1} \otimes_A \hat{A}_{\mathfrak{p}} &\cong \mathfrak{D}_{B \otimes_A \hat{A}_{\mathfrak{p}}/\hat{A}_{\mathfrak{p}}}^{-1} \\
 &\cong \mathfrak{D}_{\prod_{\mathfrak{P}|\mathfrak{p}} \hat{B}_{\mathfrak{P}}/\hat{A}_{\mathfrak{p}}}^{-1} \\
 &\cong \prod_{\mathfrak{P}|\mathfrak{p}} \mathfrak{D}_{\hat{B}_{\mathfrak{P}}/\hat{A}_{\mathfrak{p}}}^{-1} \\
 &\cong \prod_{\mathfrak{P}|\mathfrak{p}} \mathfrak{D}_{B/A}^{-1} \otimes_B \hat{B}_{\mathfrak{P}}
 \end{aligned}$$

□

3 Discriminant and ramification

Recall $\mathrm{ord}_{\hat{\mathfrak{P}}}(\mathfrak{D}_{\hat{B}_{\mathfrak{P}}/\hat{A}_{\mathfrak{p}}}) = \mathrm{ord}_{\mathfrak{P}}(\mathfrak{D}_{B/A})$. Our goal is to show that $e_{\mathfrak{P}/\mathfrak{p}} = 1$ and $\kappa(\mathfrak{P})/\kappa(\mathfrak{p})$ separable (i.e. \mathfrak{P} is unramified over K , iff $\mathfrak{P} \nmid \mathfrak{D}_{B/A}$).

In the CDVR case,

$$\begin{array}{ccccc}
 \mathfrak{P} & \hookrightarrow & B & \text{---} & L \\
 \downarrow & & \downarrow & & \downarrow \\
 \mathfrak{p} & \hookrightarrow & A & \text{---} & K
 \end{array}$$

we have $B/\mathfrak{p}B = B/\mathfrak{P}^e$.

Lemma 9.3.1.

$$\mathrm{tr}_{L/K}(b) \bmod \mathfrak{p} = e \mathrm{tr}_{l/k}(\bar{b}).$$

Proof. For all $b \in B$,

$$0 = \mathfrak{P}^e/\mathfrak{P}^e \subseteq \mathfrak{P}^{e-1}/\mathfrak{P}^e \subseteq \cdots \subseteq \mathfrak{P}/\mathfrak{P}^e \subseteq B/\mathfrak{P}^e.$$

Each adjacent quotient is 1 dimensional over l and hence f -dimensional over k . Choose a basis $\{\bar{w}_i\}_{i=1}^n$ ($n = ef$) for B/\mathfrak{P}^e as k -vector space, such that

$$\mathrm{span}_k(\{w_i\}_{i=(e-j)f+1}^{ef}) = \mathfrak{P}^{e-j}/\mathfrak{P}^e.$$

(The last jf vectors span $\mathfrak{P}^{e-j}/\mathfrak{P}^e$.) Lift $\{\bar{w}_i\}$ to $w_i \in B$ such that $w_i \bmod \mathfrak{P}^e = \bar{w}_i$. The w_i are a basis of B over A . Now

$$\begin{aligned} \mathrm{tr}_{L/K}(b) &= \mathrm{tr}_K(m_b) \\ bw_i &= (b_{i,j})(w_j)_{j=1}^n \\ \bar{b}\bar{w}_i &= (\bar{b}_{i,j})(\bar{w}_j). \end{aligned}$$

We have

$$\mathrm{tr}_{L/K}(b) = \sum_{i=1}^n b_{ii} \bmod \mathfrak{p}.$$

Now $(b_{i,j} \bmod \mathfrak{p})_{fk+1 \leq i, j \leq f(k+1)}$ represents the linear map (multiplication by \bar{b})

$$\frac{\mathfrak{P}^k/\mathfrak{P}^e}{\mathfrak{P}^{k+1}/\mathfrak{P}^e} \rightarrow \frac{\mathfrak{P}^k/\mathfrak{P}^e}{\mathfrak{P}^{k+1}/\mathfrak{P}^e}.$$

The trace as a k -linear map is $\mathrm{tr}_{l/k}(\bar{b})$. There are e such $f \times f$ blocks. □

Corollary 9.3.2.

$$\mathrm{ord}_{\mathfrak{p}}(\mathfrak{d}_{B/A}) \geq e - 1.$$

Proof. It suffices to show

$$\mathrm{ord}_{\mathfrak{p}}(\mathfrak{d}_{B/A}) \geq (e - 1)f.$$

This is since $\mathfrak{D}_{B/A} = \mathfrak{P}^c$ implies $\mathfrak{d}_{B/A} = \mathrm{Nm}_{L/K}(\mathfrak{P}^c) = \mathfrak{P}^{ef}$.

Now

$$\mathfrak{d}_{B/A} = (\det \mathrm{tr}_{L/K}(w_i w_j))$$

same as in the previous proof. Now $w_i \in \mathfrak{P}$ if $f + 1 \leq i \leq n = ef$, therefore $\bar{w}_i \in \mathfrak{P}/\mathfrak{P}^e$. For all j , $\mathrm{tr}_{L/K}(w_i w_j) \in \mathfrak{P} \cap K = \mathfrak{p}$, giving the result. □

Now consider the general case.

Theorem 9.3.3. *Suppose A, B are Dedekind. Then*

1. $\text{ord}_{\mathfrak{p}}(\mathfrak{D}_{B/A}) \geq e_{\mathfrak{p}/\mathfrak{p}} - 1$.
2. \mathfrak{P} is unramified over K iff $\mathfrak{P} \nmid \mathfrak{D}_{L/K}$.

Serre does this by Eisenstein polys.

Proof. 1. $\text{ord}_{\mathfrak{p}}(\mathfrak{D}_{B/A}) = \text{ord}_{\hat{\mathfrak{p}}}(\mathfrak{D}_{\hat{B}_{\mathfrak{p}}/\hat{A}_{\mathfrak{p}}})$. and $e_{\mathfrak{p}/\mathfrak{p}} = e_{\hat{\mathfrak{p}}/\hat{\mathfrak{p}}}$. Use the CDVR case.

2. For “ \Leftarrow ”, note $\text{ord}_{\mathfrak{p}}(\mathfrak{D}_{L/K})$ implies $0 \geq e_{\mathfrak{p}/\mathfrak{p}} - 1$ i.e. $e_{\mathfrak{p}/\mathfrak{p}} = 1$.

For “ \Rightarrow ”, it suffices to prove $\mathfrak{p} \nmid \mathfrak{D}_{B/A}$. Reduce to the CDVR case. Now

$$\det(\text{tr}_{L/K}(w_i w_j)) \bmod \mathfrak{p} = e_{\mathfrak{p}/\mathfrak{p}} \text{tr}_{l/k}(\overline{w_i w_j}) \neq 0$$

if l/k is separable (Neukirch I.2). □

3.1 Types of ramification

Definition 9.3.4: \mathfrak{P} is unramified if $e_{\mathfrak{p}/\mathfrak{p}}$ and l/k separable. For \mathfrak{P} ramified,

1. \mathfrak{P} is **tamely ramified** if either $\text{char } k = 0$ or $\text{char } k \nmid e_{\mathfrak{p}/\mathfrak{p}}$.
2. \mathfrak{P} is **wildly ramified** otherwise.

Theorem 9.3.5. \mathfrak{P} is tamely ramified over K iff

$$\text{ord}_{\mathfrak{p}}(\mathfrak{D}_{L/K}) = e_{\mathfrak{p}/\mathfrak{p}} - 1.$$

Proof. Reduce to the CDVR case.

Step 1: We show that \mathfrak{P} is tamely ramified iff $\text{tr}_{L/K}(B) = A$. Observe that $\text{tr}_{L/K}(B)$ is an ideal of A , so the latter is equivalent to $\text{tr}_{L/K}(B) \pmod{\mathfrak{p}} \neq 0$. But we know

$$\text{tr}_{L/K}(b) \bmod \mathfrak{p} = e_{\mathfrak{p}/\mathfrak{p}} \text{tr}_{l/k}(\overline{b}),$$

and $\text{tr}_{l/k}(\overline{b}) \neq 0$ (not identically 0). Hence $e_{\mathfrak{p}/\mathfrak{p}} \not\equiv 0 \pmod{\mathfrak{p}}$ iff $\text{tr}_{L/K}(b) \not\equiv 0 \pmod{\mathfrak{p}}$.

Step 2: $\text{tr}_{L/K}(B) = A \iff \text{ord}_{\mathfrak{p}}(\mathfrak{D}_{L/K}) = e_{\mathfrak{p}/\mathfrak{p}} - 1$.

We've seen

$$\text{tr}_{L/K}(\mathfrak{b}) \subseteq \mathfrak{a} \iff \mathfrak{b} \subseteq \mathfrak{a} \mathfrak{D}_{B/A}^{-1}.$$

Plug in $\mathfrak{b} = B$ to get, as ideals of B ,

$$\begin{aligned} A' := \text{tr}(B) \subseteq \mathfrak{a} &\iff B \subseteq \mathfrak{a} \mathfrak{D}_{L/K}^{-1} \\ &\iff \mathfrak{D}_{L/K} \subseteq \mathfrak{a} B \end{aligned}$$

Write $A' = \mathfrak{p}^a$. We have $\mathfrak{p}^a \mid \mathfrak{D}$ iff $\mathfrak{p}^a \mid A'$ for $a \in \mathbb{Z}$. (Power can be rational.)

For $a \in \mathbb{Z}$, $\text{ord}_{\mathfrak{p}}(A') \geq a$ iff $\text{ord}_{\mathfrak{p}}(\mathfrak{D}) \geq a$.

Thus we get

$$\text{ord}_{\mathfrak{p}}(A') \leq \underbrace{\text{ord}_{\mathfrak{p}}(\mathfrak{D}_{L/K})}_{\frac{\text{ord}_{\mathfrak{p}}(\mathfrak{D})}{e_{\mathfrak{p}/p}}} < \text{ord}_{\mathfrak{p}}(A') + 1.$$

Thus $\text{tr}_{L/K}(B) = A$ iff $a = 0$ iff $\text{ord}_{\mathfrak{p}}(\mathfrak{D}) = e - 1$.

Thus \mathfrak{P} is tamely ramified iff $v(\mathfrak{D}) = e - 1$. \square

3.2 Computation of different

Proposition 9.3.6: When A and B are CDVR's, B is generated by one element over A as an A -algebra:

$$B = A[\beta].$$

(We say that B is *monogenous* over A .)

Let $L := \text{Frac}(A)$, $K := \text{Frac}(B)$. When L/K is totally ramified, then we can choose β to be any uniformizer π_L .

Proof. Any element of B can be written as $\sum_{k \geq 0} a_k \pi_k$ where a_k are fixed representatives of $l = B/(\pi_L)$. But we can choose the a_k to be representatives of $k = A/(\pi_K)$, since $k = l$. \square

Theorem 9.3.7. (*Residue field extension separable.*) $\mathfrak{d}_{B/A} = (f'_{\beta}(\beta))$ where $f_{\beta}(x) \in A[x]$ is the minimal polynomial of β over K .

Proof.

Lemma 9.3.8.

$$\text{tr}_{L/K} \left(\frac{\beta^k}{f'(\beta)} \right) = \begin{cases} 0, & 0 \leq i \leq n-2 \\ 1, & i = n-1. \end{cases}$$

Proof. The eigenvalues of multiplication by β are just the roots β_1, \dots, β_n of the characteristic polynomial. Note that if A is a linear operator with eigenvalues λ_i and P is a polynomial then $P(A)$ has eigenvalues $P(\lambda_i)$. Hence

$$\text{tr} \left(\frac{\beta^k}{f'(\beta)} \right) = \sum_{i=1}^n \frac{\beta_i^k}{f'(\beta_i)}$$

Let $D(x_1, \dots, x_n) = \sum_{i < j} (x_i - x_j)$. Noting $f'(\beta_i) = \prod_{j \neq i} (\beta_i - \beta_j)$, the above equals

$$\frac{1}{D(x_1, \dots, x_n)} \underbrace{\sum_{i=1}^n \frac{x_i^k D(x_1, \dots, x_n)}{\prod_{j \neq i} (x_i - x_j)}}_{P(x_1, \dots, x_n)}$$

evaluated at $(x_1, \dots, x_n) = (\beta_1, \dots, \beta_n)$. Consider P . Note P is zero whenever $x_i = x_j$ for some $i \neq j$ (All except two terms are 0; those two cancel.). So $x_i - x_j \mid P$, and $D \mid P$.

However, P has degree less than $\frac{(n-1)n}{2}$ when $k < n - 1$, so must be 0. If $k = n - 1$ then we know P is a constant multiple of D , look at the coefficient of any term to see that in fact $P = D$. \square

It suffices to prove

$$(f'_\beta(\beta)^{-1}) = B^* := \{b \in L : \text{tr}_{L/K}(bb') \in A \text{ for all } b' \in B\}.$$

The condition inside is equivalent to

$$\text{tr}(b\beta^j) \in A, \quad 0 \leq j \leq n - 1.$$

(because $B = \bigoplus A\beta^i$.) But by the lemma,

$$\text{tr} \left(\sum_{i=0}^{n-1} a_i \frac{\beta^{i+j}}{f'_\beta(\beta)} \right) = a_{n-1-j} + \cdots (>).$$

“Triangular.” Therefore

$$B^* = \bigoplus A \cdot \frac{\beta^i}{f'_\beta(\beta)} = \left(\frac{1}{f'_\beta(\beta)} \right).$$

\square

Good exercise: Compute $\mathfrak{D}_{\mathbb{Q}_p(\zeta_{p^n})/\mathbb{Q}_p}$. This is tamely ramified only at $n = 1$. Totally ramified tower. The first step is $(\mathbb{Z}/p)^\times$, tame, everything else is p , wild.

(Note $G(\mathbb{Q}(\zeta_{p^n})/\mathbb{Q}) \cong (\mathbb{Z}/p^n\mathbb{Z})^\times$ because $D_p \cong G(\mathbb{Q}_p(\zeta_{p^n})/\mathbb{Q}_p)$.)

4 Ramification groups

Local, CDVR setup.

Definition 9.4.1: Let $i \geq -1$. The i th **ramification group** is

$$\begin{aligned} G_i &= \{\sigma \in G : b \in \mathcal{O}_L, v_L(\sigma(b) - b) \geq i + 1\} \\ &= \{\sigma \in G : v_L(\sigma(\beta) - \beta) \geq i + 1\}. \end{aligned}$$

Observe $G_{-1} = G$, that $G_i \supseteq G_j$ for $i \leq j$ and $\bigcap_{i \geq -1} G_i = \{1\}$. Also note for all i , G_i is a normal subgroup of G because

$$G_i = \ker(G \rightarrow \text{Aut}(\mathcal{O}_L/(\pi_L^{i+1}))).$$

In particular, G_i/G_{i+1} is a group. Furthermore G_i is defined even when i is not an integer; we have $G_i = G_{[i]}$.

We will study $\{G_i\}_{i \geq -1}$. We want

1. A formula for $v_L(\mathfrak{D}_{L/K})$. If at most tame, equals $e - 1$, else greater.
2. Look at quotients G_i/G_{i+1} . Abelian, cyclic, p -group, prime-to- p ?

4.1 $\mathfrak{D}_{L/K}$ and i_G

Definition 9.4.2: Let $\sigma \in G(L/K)$. Define $i_G : G \rightarrow \mathbb{N}_0 \cup \{\infty\}$ by

$$i_G(\sigma) = \min \{v_L(\sigma(\beta) - \beta) : \beta \in B\}.$$

Note that if $B = A[\beta]$, then

$$i_G(\sigma) = v_L(\sigma(\beta) - \beta).$$

Observe

- $i_G(\sigma) = \infty$ iff $\sigma = 1$.
- $G_i = \{\sigma \in G : i_G(\sigma) \geq i + 1\}$, so $\sigma \in G_i$ iff $i_G(\sigma) \geq i + 1$, so doesn't depend on choice of generator.

Note

$$i_G(\tau\sigma\tau^{-1}) = i_G(\sigma), \quad \sigma, \tau \in G.$$

Because $G_i \trianglelefteq G$. Note

$$i_G(\sigma\tau) \geq \min(i_G(\sigma), i_G(\tau)).$$

Because

$$\begin{aligned} i_G(\sigma\tau) &= v_L(\sigma\tau\beta - \beta) \\ &\geq \min(v_L(\sigma\tau(\beta) - \tau(\beta)), v_L(\tau(\beta) - \beta)) \\ &= \min(i_G(\sigma), i_G(\tau)). \end{aligned}$$

since $\mathcal{O}_L = \mathcal{O}_K[\beta] = \mathcal{O}_K[\tau\beta]$.

Proposition 9.4.3:

$$v(\mathfrak{D}_{L/K}) = \sum_{\sigma \neq 1} i_G(\sigma) = \sum_{i \geq 0} (|G_i| - 1).$$

($\mathfrak{a} = (\pi_L^i) \implies v_L(a) = i$.)

Proof. Let $f(x)$ be the minimal polynomial for β . Letting $n = [L : K]$,

$$f(x) = \prod_{i=1}^n (X - \beta_i).$$

Now

$$\begin{aligned} \mathfrak{D}_{L/K} &= (f'(\beta)) \\ &= \prod_{i>1} (\beta - \beta_i) \\ &= \prod_{\sigma \neq 1} (\beta - \sigma(\beta)). \end{aligned}$$

Take v_L for (1).

$$v_L(\mathfrak{D}_{L/K}) = \sum_{\sigma \neq 1} \underbrace{v_L(\beta - \sigma(\beta))}_{i_G(\sigma)}.$$

For (2), consider multiset

$$\bigsqcup_{i \geq 0} (G_i \setminus \{1\}).$$

finite. Note $\sigma \in G$ appears in $G_0, G_1, \dots, G_{i_G(\sigma)-1}$, there's $i_G(\sigma)$. Compute the size of the multiset in two different ways

$$\sum_{i \geq 0} (|G_i| - 1) = \sum_{\sigma \neq 1} i_G(\sigma).$$

□

Remark: $v_L(\mathfrak{D}_{L/K}) = e - 1$ iff $G_1 = \{1\}$ (because $|G_0| = e$, iff L/K is at most tame. Let's understand $\mathfrak{D}_{L/K}, i_G$ under sub and quotient group. Consider $L/L^H/K$. First, sub.

Proposition 9.4.4:

$$\begin{aligned} i_H(\sigma) &= i_G(\sigma) \text{ for all } \sigma \in H \\ H_i &= H \cap G_i. \end{aligned}$$

Proof. Same generator works for larger ring. $\mathcal{O}_L = \mathcal{O}_K[\beta] \implies \mathcal{O}_L = \mathcal{O}_{K'}[\beta]$. Then true by def. □

Corollary 9.4.5.

$$v_L(\mathfrak{D}_{L/K'}) = \sum_{\sigma \neq 1, \sigma \in H} \underbrace{i_G(\sigma)}_{i_H(\sigma)}.$$

For quotient.

Proposition 9.4.6: For $H \trianglelefteq G, \bar{G} \neq 1 \in G/H$,

$$i_{G/H}(\bar{\sigma}) = \frac{1}{e_{L/K'}} \sum_{\sigma \in G, \sigma \bmod H = \bar{\sigma}} i_G(\sigma).$$

Corollary 9.4.7.

$$v_{K'/K}(\mathfrak{D}_{K'/K}) = \frac{1}{e'_{L/K}} \sum_{\sigma \notin H, \sigma \in G} i_G(\sigma).$$

Because by prev. $\sum_{\bar{\sigma} \neq 1} i_{G/H}(\bar{\sigma})$ equals RHS by prop.

Proof. Choose $\alpha \in \mathcal{O}_{K'}$ and $\beta \in \mathcal{O}_L$ such that $\mathcal{O}_{K'} = \mathcal{O}_K[\alpha']$ and $\mathcal{O}_L = \mathcal{O}_K[\beta]$. Then

$$\begin{aligned} e_{L/K'} i_{G/H}(\bar{\sigma}) &= e_{L/K'} v_{K'}(\bar{\sigma}\alpha' - \alpha') \\ &= v_L(\bar{\sigma}\alpha' - \alpha'). \\ \sum_{\sigma \in G} &= i_G(\sigma) \\ &= \sum_{\tau \in H} \underbrace{i_G(\sigma\tau)}_{v_L(\sigma\tau\beta - \beta)} \\ &= v_L \left(\prod_{\tau \in H} (\sigma\tau(\beta) - \beta) \right) = \text{before.} \end{aligned}$$

fixing σ .

It suffices to prove $(\sigma\bar{\alpha}' - \alpha') = \prod_{\tau \in H} (\sigma\tau(\beta) - \beta)$. Call LHS, RHS **a**, **b**.

1. **a** | **b**: Consider

$$g(X) = \prod_{\tau \in H} (X - \tau(\beta)) \in \mathcal{O}_{K'}[x].$$

minimal polynomial of β/K' .

$$\sigma g(X) = \prod_{\tau \in H} (X - \sigma\tau(\beta)).$$

Observe $\sigma\alpha' - \alpha'$ divides coefficients of $\sigma g(X) - g(X)$. Because for all $a \in \mathcal{O}_{K'}$, $a = a_0 + a_1\alpha'$, $\sigma a = a_0 + a_1\sigma\alpha' + \dots$. Note $\sigma\alpha' - \alpha' \mid \sigma\alpha'^i - \alpha'^i$. Note $g(\beta) = 0$. Take $x = \beta$ to get

$$\sigma\alpha' - \alpha' \mid \sigma g(\beta) - \underbrace{g(\beta)}_0.$$

2. **b** | **a**. SWITCH f and g below. Cook up a minimal polynomial to show divisibility. $\alpha' = \mathcal{O}_K[\beta] = \mathcal{O}_L$. Write

$$\alpha' = \sum_{i=0}^{n-1} a_i \beta^i =: g(\beta).$$

$a_i \in \mathcal{O}_K$. $g(X) \in \mathcal{O}_K[X]$. Consider $g(X) - \alpha' \in \mathcal{O}_{K'}[X]$. By construction has β as a root.

Hence plugging in $x = \beta$,

$$\begin{aligned} f(X) &\mid g(X) - \alpha \\ \sigma f(X) &\mid \sigma(g(X)) - \sigma(\alpha') \\ \sigma f(\beta) &\mid f(\beta) - \sigma(\alpha') \end{aligned}$$

giving $\pm b \mid \pm a$.

□

End of unedited stuff.

4.2 Filtration of ramification groups

We know from (??) that

$$G_{-1}/G_0 \cong G/I_{L/K} \cong G(l/k).$$

In particular, if k is finite then G_{-1}/G_0 is finite cyclic and if $k = \bar{k}$ then G_{-1}/G_0 is trivial. From now on assume $i \geq 0$.

We aim to study the filtration

$$G \supseteq G_0 \supseteq G_1 \supseteq \cdots. \quad (9.2)$$

To do this, we first study the filtration

$$L^\times \supseteq U_L^0 \supseteq U_L^1 \supseteq \cdots \quad (9.3)$$

where

$$U_L^i = \begin{cases} \mathcal{O}_L^\times, & i = 0 \\ 1 + \pi_L^i \mathcal{O}_L, & i \geq 1. \end{cases}$$

The quotient groups in (9.3) can be understood explicitly (Proposition 9.4.8). We will relate the two filtrations by Proposition 9.4.10.¹ From this we get several important corollaries about the structure of the groups G_s . Understanding conjugates and commutators of elements in the G_s gives us several more important properties.

Proposition 9.4.8: Let K be a complete field with discrete valuation (for instance, a local field), k its residue field, and \mathfrak{m} the associated maximal ideal. Then we have isomorphisms

$$\begin{aligned} U_K/U_K^{(1)} &\xrightarrow{\cong} k^\times & U_K^{(m)}/U_K^{(m+1)} &\xrightarrow{\cong} k^+ \\ u &\mapsto u \pmod{\mathfrak{m}} & 1 + a\pi^m &\mapsto a \pmod{\mathfrak{m}}. \end{aligned}$$

Proof. For the first just note that $1 + \mathfrak{m}$ is the multiplicative unit of A/\mathfrak{m} . For the second, note $(1 + a\pi^m)(1 + b\pi^m) = 1 + (a + b)\pi^m + \cdots$. \square

To construct a map $G_i/G_{i+1} \rightarrow U_L^i/U_L^{i+1}$, we first need the following characterization of G_i .

Lemma 9.4.9. Suppose L/K is a finite Galois extension of local fields, π is a uniformizer of L , and $G = G(L/K)$. For $i \in \mathbb{N}_0$ and $\sigma \in G_0$,

$$\sigma \in G_i \iff \frac{\sigma(\pi)}{\pi} \equiv 1 \pmod{\pi_L^i}. \quad (9.4)$$

¹This will be important in local class field theory, which says there is a canonical isomorphism $K^\times/\text{Nm}_{L/K}(L^\times) \cong G(L/K)$ if L/K is finite abelian.

Proof. The RHS is equivalent to

$$\sigma(\pi) - \pi \equiv 0 \pmod{\pi_L^{i+1}}. \quad (9.5)$$

We need to show this is equivalent to

$$\sigma(\beta) - \beta \equiv 0 \pmod{\pi_L^{i+1}} \text{ for all } \beta \in L. \quad (9.6)$$

It is clear that (9.6) implies (9.5).

First suppose L/K is totally ramified. Then $\mathcal{O}_L = \mathcal{O}_K[\pi]$ by Proposition 9.3.6, giving that (9.5) implies (9.6).

Now consider the general case. We know $L/L^{L/K}$ is totally ramified (Theorem 2.7.2), so the theorem holds for $L/L^{L/K}$. Now, by Proposition 9.4.4, $G_i(L/L^{L/K}) = G_i \cap I_{L/K} = G_i$. Furthermore, since π_L is the same for L/K and $L/L^{L/K}$, the right hand-side of (9.4) does not change whether we are talking about L/K or $L/L^{L/K}$. Hence the theorem for $L/L^{L/K}$ implies the theorem for L/K . \square

Proposition 9.4.10: There is a well-defined injective group homomorphism

$$\begin{aligned} \theta_i : G_i/G_{i+1} &\hookrightarrow U_L^i/U_L^{i+1} \\ \sigma &\mapsto \frac{\sigma(\pi)}{\pi} \end{aligned}$$

that is independent of the choice of uniformizer π .

Proof. Note that

$$u \in \mathcal{O}_L, \sigma \in G_i \implies \sigma(u) \equiv u \pmod{\pi^{i+1}} \implies \frac{\sigma(u)}{u} \in U_L^{i+1}. \quad (9.7)$$

First we show θ_i is a group homomorphism $G_i \rightarrow U_L^i/U_L^{i+1}$. We have

$$\frac{\sigma\tau(\pi)}{\pi} = \frac{\sigma(\pi)}{\pi} \cdot \frac{\tau(\pi)}{\pi} \cdot \frac{\sigma\left(\frac{\tau(\pi)}{\pi}\right)}{\frac{\tau(\pi)}{\pi}}.$$

Since $\frac{\tau(\pi)}{\pi} \in \mathcal{O}_L$ and $\tau \in G_i$, (9.7) gives $\frac{\sigma\left(\frac{\tau(\pi)}{\pi}\right)}{\frac{\tau(\pi)}{\pi}} \in U_L^{i+1}$.

Lemma 9.4.9 gives that the kernel is exactly G_{i+1} , so θ_i induces an injective map $G_i/G_{i+1} \rightarrow U_L^i/U_L^{i+1}$.

Now suppose π' is another uniformizer. Write $\pi' = u\pi$ with $u \in \mathcal{O}_L^\times$. Then $\sigma \in G_i$ and (9.4.9) give

$$\frac{\sigma(\pi')}{\pi'} = \frac{\sigma(\pi)}{\pi} \cdot \underbrace{\frac{\sigma(u)}{u}}_{\in U_L^{i+1}}. \quad \square$$

Corollary 9.4.11. 1. G_0/G_1 is finite cyclic.

2. If $\text{char}(l) = 0$ then $G_1 = \{1\}$; if $\text{char}(l) = p \neq 0$, then for each $i \geq 1$,

$$G_i/G_{i+1} = (\mathbb{Z}/p\mathbb{Z})^{n_i}$$

for some n_i .

Proof. 1. Proposition 9.4.10 and 9.4.8 give $G_0/G_1 \hookrightarrow U_L/U_L^1 \cong l^\times$. But any finite subgroup of a finite field must be cyclic.

2. For $\text{char}(l) = 0$, l^+ has no finite nontrivial subgroup. For $\text{char}(l) = p$, we have $G_i/G_{i+1} \hookrightarrow U_L^i/U_L^{i+1} \cong l^+$. Just note l^+ is an abelian p -group. \square

Corollary 9.4.12. $G_0 = I_{L/K}$ is solvable. If $G(l/k) = G_{-1}/G_0$ is solvable (in particular, if k is finite) then G is solvable.

Proof. The series

$$G_0 \supseteq G_1 \supseteq \dots$$

is a solvable series for G . \square


4.3 First ramification group

Recall that we defined $G_0 = I_{L/K}$ so that we can split L/K into two parts: $L/L^{I_{L/K}}$ is totally ramified while $L^{I_{L/K}}/L$ is unramified. We can further split the extension $L/L^{I_{L/K}}$ into a wildly ramified and tamely ramified part.

Definition 9.4.13: Define the **wild inertia group** and **tame inertia group** to be

$$\begin{aligned} G_1 &= I_{L/K}^{\text{wild}} \\ G_0/G_1 &= I_{L/K}^{\text{tame}}. \end{aligned}$$

Theorem 9.4.14. The extension $L/L^{I_{L/K}^{\text{wild}}}$ is wildly ramified with Galois group $G_1 = I_{L/K}^{\text{wild}}$ and the extension $L^{I_{L/K}^{\text{wild}}}/L^{I_{L/K}^{\text{tame}}}$ is tamely ramified with Galois group G_1/G_0 .



Moreover, G_1 is the unique p -Sylow subgroup of G_0 , and

$$G_0 = G_1 \rtimes G_0/G_1.$$

Proof. Note $G_0/G_1 \hookrightarrow k^\times$ while $G_j/G_{j+1} \hookrightarrow k^+$ for $j \geq 1$; we have $p \nmid |k^\times|$ while $|k|$ is a power of p ; and $|G_1| = \prod_{1 \leq j < \infty} |G_j/G_{j+1}|$. Hence G_1 is a p -SSG of G_0 ; it is unique since it is normal and all p -SSGs are conjugate. Since the indices of the field extensions are the orders of the Galois groups, the result on tame and wild ramification follow.

Now we prove the semidirect product. This follows directly from the Schur-Zassenhaus Lemma: If H is a normal Hall subgroup of a finite group G , then H has a complement, and hence $G = H \rtimes G/H$. (A Hall subgroup $H \subseteq G$ is a group such that $\gcd(|H|, [G : H]) = 1$.)

The following is an alternate proof. We show the exact sequence

$$\begin{array}{ccccccc} 1 & \longrightarrow & G_1 & \longrightarrow & G_0 & \longrightarrow & G_0/G_1 \longrightarrow 1 \\ & & \parallel & & \parallel & & \parallel \\ & & I_{L/K}^{\text{wild}} & & I_{L/K} & & I_{L/K}^{\text{tame}} \end{array}$$

splits by showing there exists a right inverse $G_0/G_1 \rightarrow G_0$ of the projection $G_0 \rightarrow G_1$.² Since G_0/G_1 is cyclic of order $r := |l^\times|$, it suffices to find a lift $\sigma \in G_0$ of the generator $\bar{\sigma} \in G_0/G_1$ with order r . Write $|G_0| = p^s r$. Let

$$\sigma = \sigma^{p^{\varphi(r)t}}$$

where t is such that $\varphi(r)t \geq s$. Note $r \nmid p$ implies $p^{\varphi(r)t} \equiv 1 \pmod{r}$. Since $\sigma^r \in G_1$, this implies σ is still a lift of $\bar{\sigma}$. Moreover $\varphi(r)t \geq s$ gives that its order is r , so it is the desired lift. \square

Proposition 9.4.15: For $i \geq 1$, $\sigma \in G_0$, $\tau \in G_i/G_{i+1}$,

$$\theta_i(\sigma\tau\sigma^{-1}) = \theta_0(\sigma)^i \theta_i(\tau).$$

(Here $\theta_0(\sigma)^i$ is thought of as in $U_L/U_L^1 \cong l^\times$, and $\theta_i(\tau) \in U_L^i/U_L^{i+1} \cong l^+$.)

Proof. It is slightly more convenient to work additively rather than multiplicatively, so we consider

$$\begin{aligned} \theta'_i : G_i/G_{i+1} &\hookrightarrow U_L^i/U_L^{i+1} \cong (\pi^i)/(\pi^{i+1}) \\ \sigma &\mapsto \frac{\sigma(\pi)}{\pi} \mapsto \begin{cases} \frac{\sigma(\pi)}{\pi}, & i = 0 \\ \frac{\sigma(\pi)}{\pi} - 1, & i \geq 1, \end{cases} \end{aligned}$$

where π is any uniformizer.

Define

$$\pi' = \sigma^{-1}(\pi)$$

and let $a \in \mathcal{O}_L^\times$ be such that

$$\tau(\pi') = \pi' + a\pi'\pi^i.$$

²The image of G_0/G_1 is a *complement* Q of G_1 in G_0 ; the elements of Q act on G_1 by conjugation—this is what the semidirect product means.

Note that

$$\theta'_i(\tau) = \frac{\tau(\pi')}{\pi'} = a\pi^i.$$

Now we calculate, modulo $(\pi)^{i+1}$, that

$$\begin{aligned} \theta'_i(\sigma\tau\sigma^{-1}) &= \frac{\sigma\tau\sigma^{-1}(\pi)}{\pi} - 1 \\ &= \frac{\sigma\tau(\pi')}{\pi} - 1 \\ &= \frac{\sigma(\pi' + a\pi'\pi^i)}{\pi} - 1 \\ &= \frac{\pi + \sigma(a\pi'\pi^i)}{\pi} - 1 \\ &= \frac{a\sigma(\pi'\pi^i)}{\sigma(\pi')} && \text{since } \sigma(a) \equiv a \pmod{\pi^{i+1}} \\ &= \left(\frac{\sigma(\pi)}{\pi}\right)^i a\pi^i \\ &= \theta'_0(\sigma)^i \theta'_i(\tau). \end{aligned} \quad \square$$

Proposition 9.4.16: If $\sigma \in G_i$ and $\tau \in G_j$, $i, j \geq 1$, then

$$\sigma\tau\sigma^{-1}\tau^{-1} \in G_{i+j+1}.$$

Proof. □

Corollary 9.4.17. For $i \geq 1$,

$$\sigma\tau\sigma^{-1}\tau^{-1} \in G_{i+1} \iff \sigma^i \in G_1 \text{ or } \tau \in G_{i+1}.$$

Proof. We have

$$\begin{aligned} \sigma\tau\sigma^{-1}\tau^{-1} \in G_{i+1} &\iff \sigma\tau\sigma^{-1} = \tau && \text{in } G_i/G_{i+1} \\ &\iff \theta'_i(\sigma\tau\sigma^{-1}) = \theta'_i(\tau) && \text{in } (\pi^i)/(\pi^{i+1}) \\ &\iff \theta'_i(\tau)(\theta'_0(\sigma)^i - 1) = 0 && \text{by Proposition 9.4.15} \\ &\iff \theta'_i(\tau) = 0 \text{ or } \theta'_0(\sigma^i) = 1 \\ &\iff \tau \in G_{i+1} \text{ or } \sigma^i \in G_1. \end{aligned}$$

□

Corollary 9.4.18. Suppose G is abelian and $|G_0/G_1| \nmid i$. Then $G_i = G_{i+1}$.

Proof. Write $G_0/G_1 = \langle \bar{\sigma} \rangle$ where $r = |G_0/G_1|$. Since $r \nmid i$, $\bar{\sigma}^i \neq 1$; for any lift $\sigma \in G_0$ of $\bar{\sigma}$, $\sigma^i \notin G_1$. Since G is abelian, we get for all $\tau \in G_i$, $\sigma\tau\sigma^{-1}\tau^{-1} = 1$. By the previous corollary, noting $\sigma^i \notin G_1$, we must have $\tau \in G_{i+1}$. □

Definition 9.4.19: A **jump** for L/K is an integer i such that

$$G_i \neq G_{i+1}.$$

Corollary 9.4.18 tells us that jumps are divisible by $|G_0/G_1|$.

5 Herbrand's Theorem

5.1 Functions φ and ψ

Note that ramification groups behave nicely under taking subgroups (i.e. passing from M/K to M/L), by Proposition 9.4.4. However, the indices are screwed up when passing to quotient groups (i.e. passing from M/K to M/L). We calculate exactly how the index changes (Herbrand's Theorem 12.1), and use it to define a different numbering scheme that is invariant under passing to quotient groups.

It is important to know how ramification groups behave under quotients because this gives a compatible system that allows us to look at larger and larger field extensions, i.e. pass to the inverse limit.

Definition 9.5.1: Define $\varphi_{L/K} : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ by

$$\varphi_{L/K}(u) = \int_0^u \frac{1}{[G_0 : G_t]} dt$$

(recall $G_u = G_{\lceil u \rceil}$) and extend $\varphi_{L/K}$ to $\mathbb{R}_{\geq -1} \rightarrow \mathbb{R}_{\geq -1}$ by

$$\varphi(u) = u, \quad -1 \leq u \leq 0.$$

This is a piecewise linear increasing function with $\varphi_{L/K}(-1) = -1$ and with derivative at least $\frac{1}{|G_0|}$, so it is a bijection.

Definition 9.5.2: Define $\psi_{L/K} : \mathbb{R}_{-1} \rightarrow \mathbb{R}_{-1}$ by $\psi_{L/K} = \varphi_{L/K}^{-1}$. Define the **upper numbering** filtration by

$$G^v := G_{\psi_{L/K}(v)}, \quad v \geq -1.$$

5.2 Transitivity of φ and ψ

The function $\varphi_{L/K}$ gives the reindexing when we pass to the quotient Galois group.

Theorem 9.5.3 (Herbrand's Theorem). *Let $L/K'/K$ be finite Galois extension with separable residue field extension. For all $u \geq -1$,*

$$G_u H/H = (G/H)_{\varphi_{L/K'}(u)}.$$

Here, G_u is the ramification group of L/K and $(G/H)_{\varphi_{L/K'}(u)}$ is the ramification group of K'/K .

We will need several lemmas. First we relate the function $i_{G/H}(\bar{\sigma})$ and i_G evaluated at the lifts of $\bar{\sigma}$ in G .

Lemma 9.5.4. *For $\bar{\sigma} \in G/H$, $j(\bar{\sigma}) = \max_{\sigma \in \bar{\sigma}H} i_G(\sigma)$,*

$$i_{G/H}(\bar{\sigma}) - 1 = \varphi_{L/K'}(j(\bar{\sigma}) - 1).$$

Thus applying $\varphi_{L/K}$ has the effect of “turning” i_G into $i_{G/H}$. By writing out the criterion for $\sigma \in G_u$ or $(G/H)_u$ in terms of i_G and $i_{G/H}$, respectively, we will get Herbrand’s Theorem.

Proof. Pick $\sigma_0 \in G$ mapping to $\bar{\sigma}$ such that $i_G(\sigma_0) = j(\bar{\sigma})$. Then by Proposition 9.4.6,

$$i_{G/H}(\bar{\sigma}) = \frac{1}{e_{L/K'}} \sum_{\sigma \in \bar{\sigma}H} i_G(\sigma) = \frac{1}{e_{L/K'}} \sum_{\tau \in H} i_G(\sigma_0\tau). \quad (9.8)$$

We claim that

$$i_G(\sigma_0\tau) = \min(i_G(\sigma_0), i_G(\tau)) = \min(j(\bar{\sigma}), i_G(\tau))$$

for all $\tau \in H$. Indeed, by the nonarchimedean inequality,

$$i_G(\sigma_0\tau) = v_L(\sigma_0\tau(\beta) - \beta) \geq \min(v_L(\sigma_0\tau(\beta) - \tau(\beta)), v_L(\tau(\beta) - \beta)) = \min(i_G(\sigma_0), i_G(\tau)).$$

Consider two cases.

1. $i_G(\tau) = i_H(\tau) \geq i_G(\sigma_0)$. The above gives

$$i_G(\sigma_0\tau) \geq \min(i_G(\sigma_0), i_G(\tau)) \geq i_G(\sigma_0).$$

Equality holds by the maximality assumption on σ_0 .

2. $i_G(\tau) < i_G(\sigma_0)$. Then

$$i_G(\sigma_0\tau) = \min(i_G(\sigma_0), i_G(\tau)) = i_G(\tau).$$

The RHS of (9.8) then equals $\frac{1}{e_{L/K'}} \sum_{\tau \in H} \min(i_G(\sigma_0), i_G(\tau))$; the result then follows from the next lemma. \square

Lemma 9.5.5.

$$\varphi_{L/K}(u) = \frac{1}{e_{L/K}} \sum_{\sigma \in G} \min(j(\bar{\sigma}), u + 1) - 1.$$

Proof. Since both sides are piecewise linear functions, and both sides equal u for $-1 \leq u \leq 0$, it suffices to show their derivatives (slopes) are equal for $u > 0$.

If $i - 1 < u < i$ where $i \in \mathbb{N}$, then the slope of the LHS is $\frac{1}{[G:G_i]}$. For the RHS, since $i_G(\sigma)$ is an integer, each term is either $i_G(\sigma)$ or $u + 1$; each term where $u + 1$ is the minimum contributes to the slope. Hence the slope on the RHS is

$$\frac{1}{e_{L/K}} |\{\sigma \in G : u + 1 < i_G(\sigma)\}| = \frac{1}{e_{L/K}} |\{\sigma \in G : i_G(\sigma) \geq i + 1\}| = \frac{|G_i|}{e_{L/K}} = \frac{1}{[G_0 : G_i]},$$

as needed. \square

Proof of Theorem 9.5.3. We have the following string of equivalences.

1. $\bar{\sigma} \in G_u H/H = G_u/G_u \cap H$
2. There is $\sigma \in G$ lifting $\bar{\sigma}$ so that $\sigma \in G_u$.
3. $j_G(\bar{\sigma}) - 1 \geq u$.
4. $\varphi_{L/K'}(j_G(\bar{\sigma}) - 1) \geq \varphi_{L/K'}(u)$.
5. $i_{G/H}(\bar{\sigma}) - 1 \geq \varphi_{L/K'}(u)$.
6. $\bar{\sigma} \in (G/H)_{\varphi_{L/K'}(u)}$.

We have (3) \iff (4) because $\varphi_{L/K'}$ is monotonically increasing and (4) \iff (5) by Lemma 9.5.4. \square

Now we prove transitivity for φ and ψ .

Proposition 9.5.6:

$$\begin{aligned}\varphi_{L/K} &= \varphi_{K'/K} \circ \varphi_{L/K'} \\ \psi_{L/K} &= \psi_{L/K'} \circ \psi_{K'/K}.\end{aligned}$$

Proof. It suffices to prove the first equation; the first implies the second since φ and ψ are inverse. For $-1 \leq u \leq 0$ both sides equal u . Thus it suffices to show the derivatives of both sides are equal for $u \geq 0$. For $u \notin \mathbb{Z}$, the derivative on the LHS is

$$\varphi'_{L/K}(u) = \frac{1}{[G_0 : G_u]}.$$

By the chain rule, the slope on the RHS is

$$\begin{aligned}\varphi'_{K'/K}(\varphi_{L/K'}(u))\varphi'_{L/K'}(u) &= \frac{|(G/H)_{\varphi_{L/K'}(u)}| |H_u|}{|(G/H)_0| |H_0|} \\ &= \frac{|G_u H/H| |H_u|}{e_{K'/K} e_{L/K'}} && \text{by Herbrand's Theorem 12.1} \\ &= \frac{|G_u/H \cap G_u| |H_u|}{e_{L/K}} \\ &= \frac{|G_u|}{|G_0|}\end{aligned}$$

using $H \cap G_u = H_u$ (Proposition 9.4.4) and multiplicativity of ramification index. The derivatives are equal, as needed. \square

Finally, we prove the most important consequence of Herbrand's Theorem: namely, by using the upper numbering (i.e. numbering using the inverse of $\varphi_{L/K}$), quotients of ramification groups are preserved.

Proposition 9.5.7: For all $v \geq -1$,

$$G^v H/H = (G/H)^v.$$

Proof. By Herbrand's Theorem 12.1 and transitivity of ψ (Proposition 9.5.6) ($\psi_{L/K} = \psi_{L/K'} \circ \psi_{K'/K}$), we get

$$\begin{aligned} G^v H/H &= G_{\psi_{L/K}(v)} H/H \\ &= (G/H)_{\varphi_{L/K'}(\psi_{L/K}(v))} &= (G/H)_{\psi_{K'/K}(v)} = (G/H)^{\square} \end{aligned}$$

We can now define upper numbering for infinite algebraic extensions L/K .

Definition 9.5.8: Define

$$G(L/K)^v := \varprojlim_{K'/K \text{ finite Galois}} G(K'/K)^v.$$

6 Hasse-Arf Theorem

We have two different filtrations, the lower numbering filtration $\{G_u\}_{u \geq -1}$ and $\{G^v\}_{v \geq -1}$.

Definition 9.6.1: A **jump** is u such that $G_u \neq G_{u+\varepsilon}$ or v such that $G^v \neq G^{v+\varepsilon}$.

They are the x and y -coordinates of jump points, i.e. where the slope of φ changes.

Note a jump $u \in \mathbb{Z}$ since $G_u = G_{\lceil u \rceil}$. Moreover, u is a jump for the lower numbering iff $v = \varphi_{L/K}(u)$ is a upper numbering, because φ, ψ are monotonically increasing.

Theorem 9.6.2 (Hasse-Arf Theorem). *If G is finite abelian, then the jumps v are integers.*

In the cyclotomic case, G was abelian.

Remark 9.6.3: There is a nonabelian example where $v \notin \mathbb{Z}$. (See HW.)

We postpone the proof. Applications.

1. Used in local class field theory.
2. "Conductor of Galois representations" are in \mathbb{Z} , not just in \mathbb{Q} . Finite L/K , $G(L/K) \rightarrow \text{GL}_n(\mathbb{C})$.

Chapter 10

Geometric algebraic number theory

In this chapter we answer the following two questions.

1. Suppose, for every place v , we are given positive reals a_v , all but finitely many of them equal to 1. How many elements of $x \in K$ satisfy

$$|x|_v \leq a_v$$

for every v ?

2. Given a generalized ideal class \mathfrak{K} (to be defined) and a number L , how many ideals $\mathfrak{a} \in \mathfrak{K}$ satisfy $\mathfrak{N}\mathfrak{a} \leq L$? (What are the asymptotics as $L \rightarrow \infty$?) In particular, how many integral ideals satisfy $\mathfrak{N}\mathfrak{a} \leq L$?

The first question is known as the *Riemann-Roch problem* for number fields, because it is analogous to the Riemann-Roch problem in algebraic geometry¹: Given a curve C , and an integer a_P for every point (all but finitely many equal to 0), what is the dimension of the space of functions f with

$$\text{ord}_v(f) \geq -a_P$$

for every P ? ($\text{ord}_v(P)$ is the “order” of the zero of f at P .)

The second question is important because the answer will appear again when we define L -functions (because L -functions involve a sum over all ideals). This will allow us to get “explicit” formulas for quantities of interest (class number, regulator). And because it’s not much of a detour, we might as well answer the first question as well.

Our technique will be similar to that used in Chapters 3 and 5.

1 Generalized ideal classes

Also talk about adeles and stuff.

¹We will not attempt to draw a parallel in our discussion. The reader interested in seeing the correspondence should consult Neukirch [Neu99]. We follow Lang [Lan94], Chapter 6.

Proposition 10.1.1: We have the following diagram

$$\begin{array}{ccccccc}
 & & & & I(\mathfrak{c}) & \longrightarrow & I \\
 & & & & | & & | \\
 & & & & K(\mathfrak{c}) & \longrightarrow & P(\mathfrak{c}) \longrightarrow P \\
 & & & & | & & | \\
 U & \longrightarrow & UK_{\mathfrak{c}} & \longrightarrow & P_{\mathfrak{c}} & & \\
 | & & | & & & & \\
 U_{\mathfrak{c}} & \longrightarrow & K_{\mathfrak{c}} & & & &
 \end{array}$$

where each square

$$\begin{array}{ccc}
 A & \xrightarrow{\varphi} & B \\
 | & & | \\
 C & \longrightarrow & D
 \end{array}$$

means $C = \varphi^{-1}(D)$ and $A/C \cong B/D$.

Proposition 10.1.2: The group of \mathfrak{c} -ideal classes has order

$$h_{\mathfrak{c}} = \frac{h 2^{r(\mathfrak{c})} \prod_{\mathfrak{p}|\mathfrak{c}_0} \mathfrak{N}\mathfrak{p}^{m(\mathfrak{p})} \left(1 - \frac{1}{\mathfrak{N}\mathfrak{p}}\right)}{[U : U_{\mathfrak{c}}]}$$

where $r(\mathfrak{c})$ is the number of real places dividing \mathfrak{c} .

We will define the totient function by $\varphi(\mathfrak{c}) = 2^{r(\mathfrak{c})} \prod_{\mathfrak{p}|\mathfrak{c}_0} \mathfrak{N}\mathfrak{p}^{m(\mathfrak{p})} \left(1 - \frac{1}{\mathfrak{N}\mathfrak{p}}\right)$.

2 Counting lattice points

Definition 10.2.1: A subset $T \subseteq \mathbb{R}^N$ is k -Lipschitz parametrizable if there exist a finite number of Lipschitz maps $\varphi_j : [0, 1]^k \rightarrow T$ whose images cover T .

Theorem 10.2.2. Let $L \subset \mathbb{R}^N$ a lattice with fundamental domain F and $D \subset \mathbb{R}^N$ a subset whose boundary is $(N - 1)$ -Lipschitz parametrizable. Then

$$|\{x \in L : x \in tD\}| = \text{Vol}(D)\text{Vol}(F)t^N + O(t^{N-1}).$$

3 Riemann-Roch problem

4 Asymptotics of generalized ideal classes

Definition 10.4.1: For a generalized ideal class $\mathfrak{K} \in I(\mathfrak{c})/P_{\mathfrak{c}}$, let

$$j(\mathfrak{K}, t) = \{\mathfrak{a} \in \mathfrak{K} : \mathfrak{N}\mathfrak{a} \leq t\}.$$

Theorem 10.4.2.

$$j(\mathfrak{K}, t) = \frac{2^r (2\pi)^s R_{\mathfrak{c}}}{w_{\mathfrak{c}} \sqrt{d_k} \mathfrak{N}\mathfrak{c}}.$$

Part II
Class Field Theory

Chapter 11

Class Field Theory: Introduction

We give the main theorems of class field theory, deferring the proofs to the next five chapters. In this chapter we'll focus on the motivation and intuition behind the theorems. The reader may find it helpful to read this chapter along with Chapter 16, Applications.

In Section 1 we'll introduce the Frobenius map, which we need before we can state the theorems of class field theory. In Section 2 we state the theorems of local class field theory. We state two formulations of global class field theory: using ideals in Section 4 and using ideles in Section 6, after giving the relevant background on ray class groups and ideles. The formulation using ideals is less sophisticated to understand, but the formulation using ideles is more useful theoretically. We'll compare the two formulations in Section 6.1. Finally, we'll present a proof of the Kronecker-Weber Theorem using class field theory in Section 7. Throughout, we'll refer back to the cyclotomic case, because class field theory is easy to understand in this case, and it already shows much of what's at play.

1 Frobenius elements

In order to define the Artin map and state the main theorems of class field theory, we first need to understand the Frobenius map. This map takes prime ideals inside a field K to automorphisms in a Galois group $G(L/K)$. One reason for studying the Frobenius map is that $\text{Frob}_{L/K}(\mathfrak{p})$ gives information on how the prime ideal \mathfrak{p} splits in a Galois extension. First, we'll define the Frobenius element and explain what it tells us about the splitting of primes. Next, we'll look at the example of a cyclotomic extension, which suggests that something deeper is going on with the Frobenius map, which we'll attempt to explain with class field theory.

The reader may wish to review Section 7, on the decomposition and inertia groups.

The results in this section will apply to both local and global fields.

Definition 11.1.1: Let L/K be a Galois extension with Galois group G , and assume that the residue field k is finite.

1. Let \mathfrak{P} be an unramified prime of L . Define the **Frobenius element**

$$\text{Frob}_{L/K}(\mathfrak{P}) = (\mathfrak{P}, L/K)$$

to be the element $\sigma \in D_{L/K}(\mathfrak{P}) \subseteq G(L/K)$ that acts as the Frobenius automorphism on the residue field $l = \mathcal{O}_L/\mathfrak{P}$ fixing $k = \mathcal{O}_K/\mathfrak{p}$. In other words, letting $k = \mathbb{F}_q$,

$$\sigma\alpha = \alpha^q \text{ for all } \alpha \in l.$$

2. Let \mathfrak{p} be an unramified prime of K . Let \mathfrak{P} be any prime dividing \mathfrak{p} , and define $\text{Frob}_{L/K}(\mathfrak{p}) = (\mathfrak{p}, L/K)$ to be the conjugacy class of $(\mathfrak{P}, L/K)$. Equivalently (see lemma 11.1.2),

$$\text{Frob}_{L/K}(\mathfrak{p}) = (\mathfrak{p}, L/K) := \{(\mathfrak{P}, L/K) \mid \mathfrak{P}|\mathfrak{p}\}.$$

In the local case, when there is only one prime, we will simply write $\text{Frob}_{L/K}$.

Proof of existence of $(\mathfrak{P}, L/K)$. When \mathfrak{p} is unramified in L , $I(\mathfrak{P}) = 1$ so from Corollary 2.7.6, the map $D_{L/K}(\mathfrak{P}) \rightarrow G(l/k)$ is an isomorphism. Thus there is a unique element of $D_{L/K}(\mathfrak{P})$ whose image is the Frobenius element. \square

To show the above definition is valid, we need to show that changing the prime above \mathfrak{p} corresponds to conjugating the Frobenius element.

Lemma 11.1.2. *Let $\tau \in G(L/K)$. Then*

$$\begin{aligned} D(\tau\mathfrak{P}) &= \tau D(\mathfrak{P}) \tau^{-1} \\ (\tau\mathfrak{P}, L/K) &= \tau(\mathfrak{P}, L/K) \tau^{-1}. \end{aligned}$$

Therefore (since $G(L/K)$ operates transitively on the primes dividing \mathfrak{p}), the conjugacy class of $(\mathfrak{P}, L/K)$ is equal to $\{(\mathfrak{P}, L/K) \mid \mathfrak{P}|\mathfrak{p}\}$.

Proof. The first statement follows from the fact that if G acts on S and G is the stabilizer of $s \in S$, then tGt^{-1} is the stabilizer of ts . Recall that the decomposition group $D(\mathfrak{P})$ is defined as the stabilizer of \mathfrak{P} .

For the second statement, let $q = |k|$ and note that τ , as an automorphism, preserves q th powers. Hence for all $b \in \mathcal{O}_L$,

$$(\tau(\mathfrak{P}, L/K)\tau^{-1})(b) \equiv \tau(\tau^{-1}(a)^q) \equiv a^q \pmod{\tau(\mathfrak{P})}. \quad \square$$

Note that if G is abelian, then the conjugacy classes are just elements, so we can think of $(\mathfrak{P}, L/K)$ as an element of $G(L/K)$.

One of the most basic applications of the Frobenius map is to the splitting of primes in an extension.

Proposition 11.1.3: Let L/K be an extension of degree n , unramified at $\mathfrak{P} \mid \mathfrak{p}$. Then \mathfrak{p} splits into $\frac{n}{|(\mathfrak{P}, L/K)|}$ factors, where $\langle(\mathfrak{P}, L/K)\rangle$ is the subgroup of G generated by $(\mathfrak{P}, L/K)$.

In particular, \mathfrak{p} splits completely iff $(\mathfrak{p}, L/K) = 1$.

Proof. Let l and k be the residue fields.

The Frobenius element generates the decomposition group $D(\mathfrak{P})$, since it acts as the Frobenius automorphism on l/k and $D(\mathfrak{P}) \cong G(l/k)$. Hence $|D(\mathfrak{P})| = |\langle(\mathfrak{p}, L/K)\rangle|$. Since \mathfrak{p} is unramified in L , $e(\mathfrak{P}/\mathfrak{p}) = 1$ and $f(\mathfrak{P}/\mathfrak{p}) = |D(\mathfrak{P})| = |\langle(\mathfrak{p}, L/K)\rangle|$. Hence, letting g be the number of primes above \mathfrak{p} , we have

$$n = [L : K] = \underbrace{e(\mathfrak{P}/\mathfrak{p})}_1 \underbrace{f(\mathfrak{P}/\mathfrak{p})}_{|\langle(\mathfrak{p}, L/K)\rangle|} g.$$

Then

$$g = \frac{n}{|\langle(\mathfrak{p}, L/K)\rangle|},$$

as needed.

In particular, \mathfrak{p} splits completely iff $g = n$, iff $|\langle(\mathfrak{p}, L/K)\rangle| = 1$, iff $|\langle(\mathfrak{p}, L/K)\rangle| = 1$, i.e. the Frobenius element $(\mathfrak{p}, L/K)$ is trivial. \square

Next, we'll need a result of how the Frobenius element changes as we change the base field.

Proposition 11.1.4: Suppose that L/K is an unramified Galois extension, $K \subseteq K' \subseteq L$, and \mathfrak{p} is a prime of K' . Let k, k' be the residue fields of K and K' . Then

$$\text{Frob}_{L/K'}(\mathfrak{p}) = \text{Frob}_{L/K}(\mathfrak{p})^{[k':k]}$$

Note by taking the $[k' : k]$ th power we mean that if $\text{Frob}_{L/K}(\mathfrak{p})$ is the conjugacy class of σ , then $\text{Frob}_{L/K}(\mathfrak{p})^{[k':k]}$ is the conjugacy class of $\sigma^{[k':k]}$.

Proof. By definition, the left hand side induces the $|k'|$ th power map on l , while the right hand side induces the $|k| \cdot [k' : k]$ th power map on l . Hence they are equal. \square

1.1 Examples

We calculate the Frobenius map explicitly in two examples. First, a warm-up.

Example 11.1.5: For the field extension $\mathbb{Q}(i)/\mathbb{Q}$,

$$(p, \mathbb{Q}(i)/\mathbb{Q}) = \begin{cases} \text{complex conjugation,} & p \equiv 3 \pmod{4}, \\ 1, & p \equiv 1 \pmod{4}. \end{cases}$$

Proof. If $p \equiv 3 \pmod{4}$, then p remains prime in $\mathbb{Q}(i)$. The residue fields are

$$\begin{array}{c} l = \mathbb{Z}[i]/p\mathbb{Z}[i] = \mathbb{F}_{p^2} \\ | \\ k = \mathbb{Z}/p\mathbb{Z} = \mathbb{F}_p. \end{array}$$

Now $(p, \mathbb{Q}(i)/\mathbb{Q})$ must induce the p th power map on $\ell = \mathbb{F}_{p^2}$. Since this is not the identity, it must be the only element of $G(\mathbb{Q}(i)/\mathbb{Q})$ that is not the identity, i.e. complex conjugation. (This does act as the p th power, since recalling $p \equiv 3 \pmod{4}$, $(a + bi)^p \equiv a^p + b^p i^p \equiv a - bi \pmod{p}$.)

If $p \equiv 1 \pmod{4}$, then p splits in $\mathbb{Q}[i]$, say into \mathfrak{P}_1 and \mathfrak{P}_2 where $\mathfrak{P}_1, \mathfrak{P}_2$ are complex conjugate. Then $\mathbb{Z}[i]/\mathfrak{P}_1 = \mathbb{Z}[i]/\mathfrak{P}_2 = \mathbb{Z}/p\mathbb{Z} = \mathbb{F}_p$ so the extension of residue fields is trivial and the Frobenius automorphism is trivial. It is induced by the identity map, so $(p, \mathbb{Q}(i)/\mathbb{Q}) = 1$. (Note that in this case the decomposition group is trivial and does not contain complex conjugation.) \square

We generalize the above example to cyclotomic extensions.

Example 11.1.6: Let $K = \mathbb{Q}(\zeta_n)$ where ζ_n is a primitive n th root of unity. Then $G(K/\mathbb{Q}) \cong (\mathbb{Z}/n\mathbb{Z})^\times$ by identifying $k \in (\mathbb{Z}/n\mathbb{Z})^\times$ with the automorphism sending ζ_n to ζ_n^k (Proposition 6.3.1).

Suppose $\sigma := (p, L/K)$ is the map $\zeta_n \mapsto \zeta_n^k$. By definition σ reduces to the p th power map on the residue fields, so $\sigma(\zeta_n) \equiv \zeta_n^p \pmod{p\mathcal{O}_K}$. Hence

$$\zeta_n^p \equiv \zeta_n^k \pmod{p\mathcal{O}_K}.$$

But since $p \nmid n$, the n th roots of unity are distinct modulo p . (More precisely, they are distinct elements of \mathbb{F}_{p^m} where m is such that $p^m \equiv 1 \pmod{n}$.) Hence we must have $p \equiv k \pmod{n}$, i.e. σ is the p th power map.

This shows that for a prime $p \nmid n$, under the identification $G(K/\mathbb{Q}) \cong (\mathbb{Z}/n\mathbb{Z})^\times$, we have

$$(p, \mathbb{Q}(\zeta_n)/\mathbb{Q}) = p \pmod{n}.$$

This calculation of the Frobenius elements gives a complete characterization of how primes split in cyclotomic extensions. We obtain a simple proof of Theorem 6.2.4, which we restate here.

Theorem 11.1.7. *Suppose that $n = p^r m$, where $p \nmid m$. Let*

$$f = \text{ord}_m(p).$$

Then the prime factorization of (p) in $\mathbb{Q}(\zeta_n)$ is

$$(p) = (\mathfrak{P}_1 \cdots \mathfrak{P}_g)^{\varphi(p^r)}$$

where \mathfrak{P}_j are distinct primes, each with residue degree f over \mathbb{Q} , and $g = \frac{\varphi(m)}{f}$.

In particular,

$$(p) \text{ splits completely in } \mathbb{Q}(\zeta_n) \text{ iff } p \equiv 1 \pmod{n}.$$

Proof. For $r = 0$, i.e. $n = m$, the automorphism $\zeta_n \mapsto \zeta_n^p$ has order $\text{ord}_m(p)$, so the result follows from Example 11.1.6 and Proposition 11.1.3. For $r > 0$, note that (p) totally ramifies in $\mathbb{Q}(\zeta_{p^r})$ by Proposition 6.2.2, and $\mathbb{Q}(\zeta_n)$ is the compositum $\mathbb{Q}(\zeta_{p^r})\mathbb{Q}(\zeta_m)$. \square

1.2 The Frobenius map is a nice homomorphism

Because we've defined the Frobenius map on prime ideals \mathfrak{p} unramified in L , and the prime ideals are a free basis for the ideal group, we can extend the Frobenius map to the subgroup of ideals generated by unramified primes. Denoting this subgroup by I_K^S , we have a map

$$\text{Frob}_{L/K} : I_K^S \rightarrow G(L/K). \quad (11.1)$$

What is nice about this map? Look back to the cyclotomic case, Example 11.1.6. The Frobenius map didn't map the primes arbitrarily; it sent p to $p \pmod{n}$. What's to note here is that $(p, \mathbb{Q}(\zeta_n)/\mathbb{Q})$ *only depends on* $p \pmod{n}$, *information about* p *intrinsic to* \mathbb{Q} , even though $(p, \mathbb{Q}(\zeta_n)/\mathbb{Q})$ tells us about the field extension $\mathbb{Q}(\zeta_n)/\mathbb{Q}$. Thus (11.1) factors:

$$\begin{array}{ccc} I_{\mathbb{Q}}^S & \xrightarrow{\text{Frob}_{L/\mathbb{Q}}} & G(L/\mathbb{Q}) \\ \downarrow & \cong \nearrow & \\ I_{\mathbb{Q}}^S/I_{\mathbb{Q}}(1, \infty n) & & \end{array} \quad (11.2)$$

Here, $I_{\mathbb{Q}}^S$ denotes the prime ideals relatively prime to n and $I_{\mathbb{Q}}(1, \infty n)$ denotes the subgroup of ideals generated by (p) with $p \equiv 1 \pmod{n}$ and *positive*.

Something like this in fact happens in general: global class field theory tells us that for all abelian extensions, the Frobenius map “factors through a modulus,” that $(\mathfrak{p}, L/K)$ depends only on what \mathfrak{p} is modulo a nice subgroup of ideals in K . Our example essentially proves class field theory for cyclotomic extensions of \mathbb{Q} , by using the roots of unity to “keep book” on the action of Frobenius. Don't be deceived, though, the general case is much harder.

Before we look at global class field theory, we first study local class field theory. Since there's only one prime in a local field, rather than consider a map from the (rather boring) ideal group, we consider a map from the field itself.

2 Local reciprocity

When K is a nonarchimedean local field, there is a single prime ideal $\mathfrak{p} = (\pi)$. For every abelian unramified extension, the previous section gives an element of $G(L/K)$ corresponding to \mathfrak{p} , which we can think of as corresponding to π .

The main theorem of local class field theory is that we can extend this map to all elements of K^\times , and get elements in $\varprojlim_{\text{finite abelian } L/K} G(L/K) = G(K^{\text{ab}}/K)$. We will also show that this map behaves well under restricting to subextensions L/K .

Theorem 11.2.1 (Local reciprocity law). *For any nonarchimedean local field K , there exists a unique homomorphism*

$$\phi_K : K^\times \rightarrow G(K^{\text{ab}}/K),$$

*called the **local Artin (reciprocity) map** with the following properties.*

1. (*Relationship with Frobenius map*) For any prime element π of K and any finite unramified extension L of K , $\phi_K(\pi)$ acts on L as $\text{Frob}_{L/K}(\pi)$.
2. (*Isomorphism*) Let p_L be the projection $G(K^{\text{ab}}/K) \rightarrow G(L/K)$. For any finite abelian extension L/K , ϕ_K induces an isomorphism $\phi_{L/K} : K^\times / \text{Nm}_{L/K}(L^\times) \rightarrow G(L/K)$ making the following commute:

$$\begin{array}{ccc} K^\times & \xrightarrow{\phi_K} & G(K^{\text{ab}}/K) \\ \downarrow & & \downarrow p_L \\ K^\times / \text{Nm}_{L/K}(L^\times) & \xrightarrow[\cong]{\phi_{L/K}} & G(L/K). \end{array}$$

3. (*Compatibility with norm map*) For any $K \subseteq K'$, the following diagram commutes.

$$\begin{array}{ccc} K'^\times & \xrightarrow{\phi_{K'}} & G(K'^{\text{ab}}/K') \\ \downarrow \text{Nm}_{K'/K} & & \downarrow \bullet|_{K^{\text{ab}}} \\ K^\times & \xrightarrow{\phi_K} & G(K^{\text{ab}}/K) \end{array}$$

We can also say something about this map topologically.

Definition 11.2.2: A **norm group** is a subgroup of K^\times of the form $\text{Nm}_{L/K}(L^\times)$ for some finite extension L/K .

Let Frob denote the Frobenius element of l/k . The **Weil group** $W(L/K)$ of an extension L/K is equal to the inverse image of $\text{Frob}^{\mathbb{Z}}$ under the map $G(L/K) \rightarrow G(l/k)$. The topology on $W(L/K)$ is the topology from considering it as a disjoint union of cosets $I(L/K)\sigma_n$, where σ_n is any lift of Frob^n .

Note that the topology on $W(K^{\text{ab}}/K)$ as defined above is strictly finer than the topology it inherits from $G(K^{\text{ab}}/K)$ (see exercise 2.1).

Theorem 11.2.3 (Local existence theorem). *Let K be a nonarchimedean local field. The norm groups of K are exactly the open subgroups of finite index.*

Theorem 11.2.4 (Topological isomorphism for LCFT). *The image of the Artin map is the Weil group $W(L/K)$, and ϕ_K gives an isomorphism of topological groups $K^\times \rightarrow W(L/K)$. It restricts to an isomorphism $U_K \rightarrow I(L/K)$.*

Combining Theorems 11.2.1 and 11.2.3 gives the following bijective correspondence.

Theorem 11.2.5. *Let K be a nonarchimedean local field. Then there is a bijective correspondence between finite abelian extensions of K and the set of open subgroups of finite index of K^\times , given by*

$$L \mapsto \text{Nm}_{L/K}(L^\times).$$

Furthermore, this is an inclusion-reserving bijection that takes intersections to products and products to intersections:

$$\begin{aligned} L \subseteq M &\iff \text{Nm}_{L/K}(L^\times) \supseteq \text{Nm}_{M/K}(M^\times) \\ \text{Nm}_{L \cdot L'/K}((L \cdot L')^\times) &= \text{Nm}_{L/K}(L^\times) \cap \text{Nm}_{L'/K}(L'^\times) \\ \text{Nm}_{L \cap L'/K}((L \cap L')^\times) &= \text{Nm}_{L/K}(L^\times) \cdot \text{Nm}_{L'/K}(L'^\times). \end{aligned}$$

Finally, every subgroup of K^\times containing a norm group is a norm group.

The following gives a sort-of converse statement: nonabelian extensions cannot be described by norm groups.

Theorem 11.2.6 (Norm limitation theorem). *Let L be a finite extension of a local field K , and K' be the largest abelian extension of K contained in L . Then*

$$\text{Nm}_{L/K}(L^\times) = \text{Nm}_{K'/K}(K'^\times).$$

3 Ray class groups

In order to define the Frobenius element of a prime we need the extension to be unramified. However, when K is a global field, we cannot as easily say an extension L/K is “unramified,” because \mathcal{O}_K has many prime ideals. Requiring that L/K to be unramified at all primes of K is too restrictive, because most fields L do not satisfy this condition.

Thus, we instead focus on a set of primes S and consider extensions L/K that are unramified outside of S . When we define Frobenius elements, we have to exclude S , and when we define a reciprocity map we have to exclude the subgroup that these primes generate. (Note that unlike in local reciprocity, we will not define ϕ_K with domain K^\times , but rather with domain a subgroup of the ideal group I_K .)

Letting S range over all finite subsets, we will account for all finite abelian extensions L/K , because each extension is ramified at only finitely many primes (Theorem 2.6.1).

This motivates the following definition.

Definition 11.3.1: Let I_K be the group of fractional ideals of K . Define I_K^S to be the subgroup of I_K generated by prime ideals not in S .

Let L/K be an extension of K . Define $I_L^S := I_L^{S'}$, where S' is the set of prime ideals lying above a prime ideal in S .

Note that if $S \subseteq T$ then $I_K^S \supseteq I_K^T$.

Similar to Theorem 11.2.1, global class field theory will tell us there is a map

$$I_K^S / \text{Nm}_{L/K}(I_L^S) \rightarrow G(L/K)$$

when S contains the primes that ramify in L . However, this is not an isomorphism until we take a further quotient, namely, the quotient with a subgroup of principal ideals $P_K(1, \mathfrak{m})$, which we will define. First we need the following.

Definition 11.3.2: A **modulus** \mathfrak{m} is a formal product of places of K , where

1. Finite primes have exponents in \mathbb{N}_0 , and only finitely many exponents are nonzero.
2. Infinite real places have exponents 0 or 1.
3. Infinite complex places do not appear.

We say a place divides \mathfrak{m} if it appears with positive exponent. We write

$$\mathfrak{m} = \underbrace{\prod_{\mathfrak{p} \text{ finite}} \mathfrak{p}^{m(\mathfrak{p})}}_{\mathfrak{m}_0} \underbrace{\prod_{v \text{ real}} v^{m(v)}}_{\mathfrak{m}_\infty}.$$

In other words, a modulus is the product of a proper ideal with some number of real places.

Definition 11.3.3: Let $S(\mathfrak{m})$ denote the set of finite primes dividing \mathfrak{m} .

Define $K(1, \mathfrak{m})$ (“elements of K that are 1 modulo \mathfrak{m} ”) to be the subgroup of elements of K^\times satisfying the following.

$$\begin{cases} \text{ord}_{\mathfrak{p}}(a - 1) \geq m(\mathfrak{p}), & \text{finite } \mathfrak{p} \mid \mathfrak{m} \\ a_v > 0, & \text{real } v \mid \mathfrak{m}. \end{cases}$$

Let $i : K^\times \rightarrow I_K$ be the map sending a to (a) , and let

$$P_K(1, \mathfrak{m}) := i(K(1, \mathfrak{m})).$$

Define the **ray class group** of \mathfrak{m} to be

$$C_K(\mathfrak{m}) = I_K^{S(\mathfrak{m})} / P_K(1, \mathfrak{m}).$$

Note that $P_K(1, \mathfrak{m}) \in I_K^{S(\mathfrak{m})}$ because if $a \in K(1, \mathfrak{m})$ and $\mathfrak{p} \in S(\mathfrak{m})$, then $\text{ord}_{\mathfrak{p}}(a - 1) \geq 1$ and $\text{ord}_{\mathfrak{p}}(a) = 0$, i.e. $\mathfrak{p} \nmid (a)$. We will often abbreviate $I^{S(\mathfrak{m})}$ as $I^{\mathfrak{m}}$.

Example 11.3.4: If $\mathfrak{m} = 1$ then $P_K(1, \mathfrak{m})$ is the subgroup of principal ideals and $C_K(\mathfrak{m})$ is just the ideal class group.

If $\mathfrak{m} = \prod_{v \text{ real}} v$, then

$$C_K(\mathfrak{m}) = I_K / \{(a) \in I_K : a_v > 0 \text{ for all real } v\}$$

is called the **narrow class group**. We are only modding out by the “totally positive” principal ideals, so it is larger than the class group.

Definition 11.3.5: A **congruence subgroup** for K modulo \mathfrak{m} is a subgroup H such that

$$P_K(1, \mathfrak{m}) \subseteq H \subseteq I_K^{S(\mathfrak{m})}.$$

The corresponding **generalized ideal class group** is $I_K^{S(\mathfrak{m})} / H$.

We will show that generalized ideal class groups are exactly the Galois groups of abelian extensions of K .

Finally, in preparation for the global reciprocity theorem, we say what it means exactly for a map to only depend on modulo conditions, like the Frobenius map we considered in Section 1.2.

Definition 11.3.6: A homomorphism $\psi : I^S \rightarrow G$ **admits a modulus** if there exists a modulus \mathfrak{m} with $S(\mathfrak{m}) = \mathfrak{m}$ such that ψ factors through $I^S/P_K(1, \mathfrak{m})$. In other words, there exists a modulus \mathfrak{m} with $S(\mathfrak{m}) = S$ such that

$$\psi(P_K(1, \mathfrak{m})) = 0.$$

4 Global reciprocity

In this section K is a global field.

Theorem 11.4.1 (Global reciprocity theorem). *Let L/K be a finite abelian extension. Let S be the set of primes ramifying in L . There is a unique map $\psi_{L/K}$ such that for a prime ideal $\mathfrak{p} \notin S$, $\psi_{L/K}(\mathfrak{p})$ acts on L as $\text{Frob}_{L/K}(\mathfrak{p})$. Moreover, $\psi_{L/K}$ satisfies the following properties.*

1. (Isomorphism) $\psi_{L/K}$ admits a modulus \mathfrak{m} with $S(\mathfrak{m}) = S$ and $\psi_{L/K}$ induces an isomorphism

$$\psi_{L/K} : I_K^S / (P_K(1, \mathfrak{m}) \cdot \text{Nm}_{L/K}(I_L^S)) \xrightarrow{\cong} G(L/K).$$

2. (Compatibility over all extensions) Suppose $S \subseteq T$, and $L/K, M/K$ are finite abelian extensions such that $L \subseteq M$ and such that the set of primes ramifying in L, M are contained in S, T , respectively. Then the following commutes, where p_L is the projection map.

$$\begin{array}{ccc} I_K^T & \xrightarrow{\psi_{M/K}} & G(M/K) \\ \downarrow & & \downarrow p_L \\ I_K^S & \xrightarrow{\psi_{L/K}} & G(L/K). \end{array}$$

3. (Compatibility with norm map) For $K \subseteq K' \subseteq L$, the following diagram commutes.

$$\begin{array}{ccc} I_{K'}^S & \xrightarrow{\psi_{L/K'}} & G(L/K') \\ \downarrow \text{Nm}_{K'/K} & & \downarrow \\ I_K^S & \xrightarrow{\psi_{L/K}} & G(L/K) \end{array}$$

Remark 11.4.2: The uniqueness of $\psi_{L/K}$ is clear from the fact that I_K^S is freely generated by prime ideals. Part 2 follows immediately from the definition of $\psi_{L/K}$ and $\psi_{M/K}$, and part

3 follows immediately from the existence of $\psi_{L/K}$ and $\psi_{L/K'}$, as we show below. The crux of the theorem is part 1.

For part 2, since primes generate I_K^S , it suffices to show that for any prime $\mathfrak{p} \in I_K^S$,

$$\psi_{L/K}(\mathfrak{p}) = p_L(\psi_{M/K}(\mathfrak{p})).$$

But by definition, the left-hand side is $\text{Frob}_{L/K}(\mathfrak{p})$ and the right-hand side is $p_L(\text{Frob}_{M/K}(\mathfrak{p}))$. Now p_L induces the map $G(m/k) \rightarrow G(l/k)$, so both sides act on k as the $|k|$ th power Frobenius, and are equal.

For part 3, we need to show for any prime $\mathfrak{p} \in I_{K'}^S$,

$$\psi_{L/K'}(\mathfrak{p}) = \psi_{L/K}(\text{Nm}_{K'/K}(\mathfrak{p})).$$

But by definition, the left-hand side is $\text{Frob}_{L/K'}(\mathfrak{p})$ and the right-hand side is $\psi_{L/K}(\mathfrak{p}^{[k':k]}) = \text{Frob}_{L/K}(\mathfrak{p})^{[k':k]}$. The result now follows from Proposition 11.1.4.

Example 11.4.3 (Cyclotomic extensions): In Section 1.2, we showed that the global reciprocity theorem (part 1 above) holds for a cyclotomic extension $\mathbb{Q}(\zeta_n)/\mathbb{Q}$. Indeed, letting \mathfrak{m} be $n\infty$, we have that $I_K^{\mathfrak{m}}/P_K(1, \mathfrak{m}) \xrightarrow{\cong} G(\mathbb{Q}(\zeta_n)/\mathbb{Q})$ as in (11.2). (Note that $\text{Nm}_{L/K}(I_L^S) \subseteq P_K(1, \mathfrak{m})$ will follow from the first inequality 15.2.1.)

Note the modulus in Theorem 11.4.1 has to be divisible by all primes ramifying in L , and the primes have to have large enough exponents for $\ker(\psi_{L/K})$ to be a congruence subgroup modulo \mathfrak{m} . There is a canonical choice for \mathfrak{m} , namely the modulus with least exponents. It is called the **conductor** of the extension L/K , and denoted by $\mathfrak{f}(L/K)$.

We have the following analogue of Theorem 11.2.3.

Theorem 11.4.4 (Existence theorem). *Let H be a congruence subgroup modulo \mathfrak{m} . Then there exists an abelian extension L/K such that*

$$H = P_K(1, \mathfrak{m}) \cdot \text{Nm}_{L/K}(I_L^{\mathfrak{m}}) = \ker(\psi_{L/K}).$$

In particular, this applies when $H = P_K(1, \mathfrak{m})$.

Definition 11.4.5: For each modulus \mathfrak{m} there is a field $K_{\mathfrak{m}}$, called the **ray class field** of K modulo \mathfrak{m} such that $\psi_{K_{\mathfrak{m}}/K}$ defines an isomorphism

$$C_K(\mathfrak{m}) \xrightarrow{\cong} G(K_{\mathfrak{m}}/K).$$

Example 11.4.6: Since $\infty(n)$ is the smallest modulus such that $\psi_{\mathbb{Q}(\zeta_n)/\mathbb{Q}}$ factors through $I_K^{\mathfrak{m}}/P_K(1, \mathfrak{m})$, $\infty(n)$ is the conductor of $\mathbb{Q}(\zeta_n)$. Since we actually have an isomorphism

$$C_K(\infty n) = I_K^{\infty n}/P_K(1, \infty n) \xrightarrow{\cong} G(\mathbb{Q}(\zeta_n)/\mathbb{Q}),$$

$\mathbb{Q}(\zeta_n)$ is in fact the ray class field of $\infty(n)$.

We have that $\mathbb{Q}(\zeta_n + \zeta_n^{-1})$ is the ray class field of (n) (see exercise 1.2).

Putting this all together, if we fix a modulus \mathfrak{m} we have the following bijection between extensions and subgroups.

Theorem 11.4.7. *Fix a modulus \mathfrak{m} and a global field K . The map $L \mapsto \text{Nm}_{L/K}(C_L(\mathfrak{m}))$ is a bijection between*

1. *the set of abelian extensions of K in the ray class field $K_{\mathfrak{m}}$ and*
2. *the set of subgroups of $C_K(\mathfrak{m})$.*

Moreover, it reverses inclusions and switches products and intersections:

$$\begin{aligned} L \subseteq M &\iff \text{Nm}_{L/K}(C_L(\mathfrak{m})) \supseteq \text{Nm}_{M/K}(C_M(\mathfrak{m})) \\ \text{Nm}_{L_1 \cdot L_2/K}(C_{L_1 \cdot L_2}(\mathfrak{m})) &= \text{Nm}_{L_1/K}(C_{L_1}(\mathfrak{m})) \cap \text{Nm}_{L_2/K}(C_{L_2}(\mathfrak{m})) \\ \text{Nm}_{L_1 \cap L_2/K}(C_{L_1 \cap L_2}(\mathfrak{m})) &= \text{Nm}_{L_1/K}(C_{L_1}(\mathfrak{m})) \cdot \text{Nm}_{L_2/K}(C_{L_2}(\mathfrak{m})). \end{aligned}$$

Note Theorem 11.4.1 is like Theorem 11.2.1 except that we're only working with finite extensions L/K instead of putting them together into K^{ab}/K . We cannot combine the maps $\psi_{L/K}$ because they are defined on different groups. Hence we now take a different approach, using ideles.

5 Ideles

In this section we give an alternate statement of the main theorems of global class field theory.

In local class field theory, we had isomorphisms $K^\times / \text{Nm}_{L/K}(L^\times) \cong G(L/K)$. For this to be true, $\text{Nm}_{L/K}(L^\times)$ must have finite index in K^\times . However, this is no longer true when K is a global field. (If K is local, it is complete with respect to a valuation, and $\text{Nm}_{L/K}(x) = y$ has solutions in y for many x , in the same way that Hensel's lemma often gives solutions over complete fields.)

We want to work with complete fields but K comes with a bunch of different places. The solution is to complete K at *every place at once* and combine the information into the adèle ring and idele group. Then we will get statements for global class field theory that resemble local class field theory, with K^\times replaced by \mathbf{C}_K , a group related to the idele group (to be defined).

Definition 11.5.1: Abbreviate $\mathcal{O}_v = \mathcal{O}_{K_v}$. The **adèle ring** of K is

$$\mathbb{A}_K = \left\{ (a_v) \in \prod_{v \in V_K} K_v : a_v \in \mathcal{O}_v \text{ for all but finitely many } v \right\}.$$

We write this as $\prod'_{v \in V_K} (K_v, \mathcal{O}_v)$. Equip it with a topology by letting a basis for open sets be $\prod_v U_v$, where U_v is open in K_v for all v and $U_v = \mathcal{O}_v$ for almost all v . In other words, it is the unique topology from which $\prod_v \mathcal{O}_v$ inherits the product topology and is open.

The **idele group** of K is the group of units of the above:

$$\mathbb{I}_K = \mathbb{A}_K^\times = \prod_{v \in V_K}' (K_v^\times, \mathcal{O}_v^\times) = \left\{ (a_v) \in \prod_{v \in V_K} K_v^\times : a_v \in \mathcal{O}_v^\times \text{ for all but finitely many } v \right\}.$$

Equip it with a topology by letting a basis for open sets be $\prod_v U_v$, where U_v is open in K_v^\times for all v and $U_v = \mathcal{O}_v^\times$ for almost all v . In other words, it is the unique topology from which $\prod_v \mathcal{O}_v^\times$ inherits the product topology and is open.

Be careful: the topology of the idele group is not the subspace topology induced from the adèle ring.

Definition 11.5.2: For a finite set S containing all infinite places, let $\mathbb{I}_K^S = \prod_{v \in S} K_v^\times \times \prod_{v \notin S} \mathcal{O}_v^\times$. In other words, \mathbb{I}_K^S contains those ideles that are units away from S . Give \mathbb{I}_K^S the subspace topology inherited from \mathbb{I}_K .

Note the topology on \mathbb{I}_K^S is just the product topology, and that $\mathbb{I} = \bigcup_S \mathbb{I}_K^S$.

Proposition 11.5.3: \mathbb{I}_K^S is locally compact.

Proof. $\prod_{v \in S} K_v^\times$ is a finite product of locally compact spaces; $\prod_{v \notin S} \mathcal{O}_v^\times$ is a product of compact spaces (Proposition 8.1.3) so compact by Tychonoff's Theorem. Since a finite product of locally compact spaces is compact, the result follows. \square

Think of the ideles as a thickening of ideals: it includes factors for infinite places, and includes units at finite primes. We can embed K^\times via the diagonal map, and K_v^\times via the inclusion map.

Definition 11.5.4: Define $i : K \hookrightarrow \mathbb{A}_K$ by the diagonal map $i(a) = (a, a, \dots)$ and $i_v : K_v \hookrightarrow \mathbb{A}_K$ by the inclusion map $i_v(a) = (1, \dots, 1, \underbrace{a}_v, 1, \dots, 1)$. Also denote by i, i_v the maps restricted to $K^\times \hookrightarrow \mathbb{I}_K$ and $i_v : K_v^\times \hookrightarrow \mathbb{I}_K$.

Proposition 11.5.5: $i(K^\times)$ is discrete in \mathbb{I}_K .

Proof. Given $a \in K^\times$, let S be set of places containing the infinite places and the finite places where $v(a) \neq 0$. Consider the open set

$$U = \{ \mathbf{x} \in \mathbb{I}_K : |x_v - a|_v < \varepsilon \text{ for } v \in S, x_v \in U_v \text{ for } v \notin S \}$$

containing $i(a)$. If $i(b) \in U$ with $a \neq b$, then

$$\prod_v |b - a|_v < \varepsilon^{|S|} < 1,$$

contradicting the product formula 19.1. Hence $i(K^\times) \cap U = \{i(a)\}$. \square

Definition 11.5.6: The **idele class group** is defined to be

$$\mathbf{C}_K = \mathbb{I}_K / K^\times,$$

where K^\times is thought of as a subgroup of \mathbb{I}_K by the diagonal map i .

We define a norm on adeles by defining it componentwise.

Definition 11.5.7: The **norm**, from L to K is the function $\text{Nm}_{L/K} : \mathbb{A}_L \rightarrow \mathbb{A}_K$ defined by

$$\text{Nm}_{L/K}((x_w)_{w \in V_L}) = \left(\prod_{\substack{w|v \\ v \in V_K}} \text{Nm}_{L_w/K_v}(x_w) \right).$$

This descends to a function $\text{Nm}_{L/K} : \mathbb{I}_L \rightarrow \mathbb{I}_K$.

5.1 Ray class groups vs. ideles

We will need the following to go between the interpretations of global class field theory via ray class groups and via ideles. The statement in terms of ray class groups is easier for concrete applications, but the statement in terms of ideles is better abstractly, and more convenient to prove. (But to complicate things more, certain parts of the proof will be easier to think of in terms of ray class groups.)

We can go from $\mathbb{I}_K \rightarrow I_K$ easily, via the map

$$p(\mathbf{a}) = \prod_{v=v_p \text{ finite}} \mathfrak{p}^{v(a_v)} \tag{11.3}$$

(also denoted simply (\mathbf{a})). However, if we want the image to be in $I_K^{S(\mathfrak{m})}$, we need to focus our attention on a subset of ideles $\mathbb{I}_K(1, \mathfrak{m})$ (defined below). Taking the map $\mathbb{I}_K(1, \mathfrak{m}) \rightarrow I_K^{S(\mathfrak{m})}$ and modding out by appropriate groups then makes it a bijection. We also need to check that we don't lose anything when we consider only ideles of the form $\mathbb{I}_K(1, \mathfrak{m})$; that is, that the inclusion $\mathbb{I}_K(1, \mathfrak{m}) \hookrightarrow \mathbb{I}_K$ is a bijection, again after modding out by appropriate groups. This is Proposition 11.5.9 below.

Definition 11.5.8: For a place $v \mid \mathfrak{m}$, define

$$I(\mathfrak{m})_v = \begin{cases} \mathbb{R}_{>0}, & v \text{ real} \\ 1 + \mathfrak{p}^{m(p)}, & v = v_p \text{ finite.} \end{cases}$$

Let \mathcal{O}_v^\times be the group of units of K_v . (For v infinite, $\mathcal{O}_v^\times := K_v^\times$). Define

$$\mathbb{I}_K(1, \mathfrak{m}) = \prod_{v \mid \mathfrak{m}} I(\mathfrak{m})_v \times \prod'_{v \nmid \mathfrak{m}} (K_v^\times, \mathcal{O}_v^\times) \tag{11.4}$$

$$\mathbb{U}_K(1, \mathfrak{m}) = \prod_{v \mid \mathfrak{m}} I(\mathfrak{m})_v \times \prod_{v \nmid \mathfrak{m}} \mathcal{O}_v^\times$$

$$K(1, \mathfrak{m}) = i(K^\times) \cap \mathbb{I}_K(1, \mathfrak{m}).$$

Let $\mathbb{U}_K := \mathbb{U}_K(1, 1)$.

Compare (11.4) to the definition of $P_K(1, \mathfrak{m})$.

Proposition 11.5.9: We have the following maps.

$$\begin{array}{ccc} \mathbb{I}_K(1, \mathfrak{m})/K(1, \mathfrak{m}) & \xrightarrow{\cong} & \mathbb{I}_K/K^\times = \mathbf{C}_K \\ \downarrow & & \\ \mathbb{I}_K(1, \mathfrak{m})/K(1, \mathfrak{m})\mathbb{U}_K(1, \mathfrak{m}) & \xrightarrow{\cong} & C_K(\mathfrak{m}). \end{array}$$

The bottom map is induced by the map $p : \mathbb{I}_K \rightarrow I_K^{\mathfrak{m}}$ and the top map is induced by inclusion. Moreover, for any finite Galois L/K such that

$$\mathbb{U}_K(1, \mathfrak{m}) \subseteq \text{Nm}_{L/K}(\mathbb{I}_L),$$

this diagram induces isomorphisms

$$\begin{array}{ccc} \mathbb{I}_K(1, \mathfrak{m})/[K^\times \text{Nm}_{L/K} \mathbb{I}_L \cap \mathbb{I}_K(1, \mathfrak{m})] & \xrightarrow{\cong} & \mathbb{I}_K/K^\times \text{Nm}_{L/K} \mathbb{I}_L \\ & \searrow \cong & \\ & & I_K^{\mathfrak{m}}/(P_K(1, \mathfrak{m}) \cdot \text{Nm}_{L/K}(I_L^{\mathfrak{m}})). \end{array}$$

Proof. For the bottom map, consider the exact sequence

$$0 \rightarrow K^\times \cap \mathbb{I}_K(1, \mathfrak{m}) = K(1, \mathfrak{m}) \xrightarrow{i} \mathbb{I}_K(1, \mathfrak{m}) \xrightarrow{p} I^{S(\mathfrak{m})} \rightarrow 0.$$

We have that $\mathbb{I}_K(1, \mathfrak{m})/K(1, \mathfrak{m}) = \text{coker } i$, so we use the kernel-cokernel sequence.¹ We have $\ker p = \mathbb{U}_K(1, \mathfrak{m})$, and $\text{coker } p \circ i = I^{S(\mathfrak{m})}/p(K(1, \mathfrak{m})) = I^{S(\mathfrak{m})}/P_K(1, \mathfrak{m}) = C_K(\mathfrak{m})$, so this gives the exact sequence

$$\mathbb{U}_K(1, \mathfrak{m}) \rightarrow \mathbb{I}_K(1, \mathfrak{m})/K(1, \mathfrak{m}) \rightarrow C_K(\mathfrak{m}) \rightarrow 1,$$

which gives the bottom isomorphism.

The top map is clearly injective. For surjectivity, take $a \in \mathbb{I}_K$. By the weak approximation theorem 7.3.4, there exists b so that $\frac{av}{b_v} \in \mathfrak{p}^{m(\mathfrak{p})} + 1$ for every $v = v_{\mathfrak{p}}$ dividing \mathfrak{m} . Then $\frac{a}{b} \in \mathbb{I}_K(1, \mathfrak{m})$, and its image varies in $\mathbb{I}_K/K^\times \text{Nm}_{L/K} \mathbb{I}_L$ from a by the constant factor $b \in K^\times$.

Now we show the second diagram. (Warning: this proof is not very enlightening.) Let p and p' denote the maps $\mathbb{I}_K \rightarrow I_K$ and $\mathbb{I}_K(1, \mathfrak{m}) \rightarrow I_K^S$, respectively. Note that the first diagram gives isomorphisms

$$\begin{array}{ccc} \mathbb{I}_K(1, \mathfrak{m})/((K^\times \text{Nm}_{L/K} \mathbb{I}_L) \cap \mathbb{I}_K(1, \mathfrak{m})) & \xrightarrow{\cong} & \mathbb{I}_K/K^\times \text{Nm}_{L/K} \mathbb{I}_L \\ \downarrow & & \\ \mathbb{I}_K(1, \mathfrak{m})/K(1, \mathfrak{m})\mathbb{U}_K(1, \mathfrak{m})p'^{-1}(\text{Nm}_{L/K}(I_L^{\mathfrak{m}})) & \xrightarrow{\cong} & I_K^{\mathfrak{m}}/(P_K(1, \mathfrak{m}) \cdot \text{Nm}_{L/K}(I_L^{\mathfrak{m}})). \end{array}$$

¹Given $A \xrightarrow{f} B \xrightarrow{g} C$, there is an exact sequence

$$0 \rightarrow \ker f \rightarrow \ker g \circ f \rightarrow \ker g \rightarrow \text{coker } f \rightarrow \text{coker } g \circ f \rightarrow \text{coker } g \rightarrow 0.$$

This is proven using the snake lemma.

We have that

$$K(1, \mathfrak{m})\mathbb{U}_K(1, \mathfrak{m})p'^{-1}(\mathrm{Nm}_{L/K}(I_L^S)) \quad (11.5)$$

$$= K(1, \mathfrak{m})\mathbb{U}_K(1, \mathfrak{m})p'^{-1}(\langle \mathfrak{p}^{f(w/v)} \mid w \mid v \notin S \rangle), \quad f_v = \text{residue degree} \quad (11.6)$$

$$= K(1, \mathfrak{m})\mathbb{U}_K(1, \mathfrak{m})(\mathbb{I}_K(1, \mathfrak{m}) \cap \mathbb{U}_K \mathrm{Nm}_{L/K}(\mathbb{I}_L)) \quad (11.7)$$

$$= \mathbb{U}_K(1, \mathfrak{m})(\mathbb{I}_K(1, \mathfrak{m}) \cap (K^\times \mathrm{Nm}_{L/K} \mathbb{I}_L)) \quad (11.8)$$

$$= (K^\times \mathrm{Nm}_{L/K} \mathbb{I}_L) \cap \mathbb{I}_K(1, \mathfrak{m}). \quad (11.9)$$

(11.6) follows from the fact that if $\mathfrak{P} \mid \mathfrak{p}$, then $\mathrm{Nm}_{L/K}(\mathfrak{P}) = \mathfrak{p}^{f(\mathfrak{P}/\mathfrak{p})}$. To go between (11.6) and (11.7), note that $\mathfrak{p}^{f(w/v)} = p(\mathrm{Nm}_{L/K}(1, \dots, 1, \underbrace{\pi_w}_w, 1, \dots, 1))$, and that $\ker(p) = \mathbb{U}_K$.

Now we go between (11.7) and (11.8). For “ \subseteq ,” suppose $a \in \mathbb{U}_K$ and $b \in \mathrm{Nm}_{L/K}(\mathbb{I}_L)$ such that $a \mathrm{Nm}_{L/K} b \in \mathbb{I}_K(1, \mathfrak{m})$. Suppose c agrees with a for every $v \mid \mathfrak{m}$, and is 1 everywhere else. Then $ac^{-1} \in \mathbb{U}_K(1, \mathfrak{m}) \subseteq \mathbb{I}_K(1, \mathfrak{m})$. Since $a \mathrm{Nm}_{L/K} b \in \mathbb{I}_K(1, \mathfrak{m})$ as well and $\mathbb{I}_K(1, \mathfrak{m})$ is a group, we must have $c \mathrm{Nm}_{L/K} b \in \mathbb{I}_K(1, \mathfrak{m})$. Hence

$$a \mathrm{Nm}_{L/K} b = \underbrace{ac^{-1}}_{\in \mathbb{U}_K(1, \mathfrak{m})} \underbrace{c \mathrm{Nm}_{L/K} b}_{\in \mathbb{I}_K(1, \mathfrak{m}) \cap (K^\times \mathrm{Nm}_{L/K} \mathbb{I}_L)},$$

as needed. Furthermore note $K(1, \mathfrak{m}) \subseteq \mathbb{I}_K(1, \mathfrak{m}) \cap K^\times \mathrm{Nm}_{L/K} \mathbb{I}_L$. For “ \supseteq ,” suppose $a \in K^\times$ and $b \in \mathrm{Nm}_{L/K}(\mathbb{I}_L)$ such that $a \mathrm{Nm}_{L/K} b \in \mathbb{I}_K(1, \mathfrak{m})$. By weak approximation, take $c \in K^\times$ sufficiently close to $\frac{1}{b_v}$ with respect to v , for every $v \in \mathfrak{m}$, so that $\mathrm{Nm}_{L/K}(cb) \in \mathbb{I}_K(1, \mathfrak{m})$. Then $a \mathrm{Nm}_{L/K}(c^{-1}) \in \mathbb{I}_K(1, \mathfrak{m})$ as well, and in fact in $K(1, \mathfrak{m})$. Then

$$a \mathrm{Nm}_{L/K} b = \underbrace{a \mathrm{Nm}_{L/K}(c^{-1})}_{\in K(1, \mathfrak{m})} \underbrace{\mathrm{Nm}_{L/K} cb}_{\in \mathbb{I}_K(1, \mathfrak{m}) \cap \mathrm{Nm}_{L/K} \mathbb{I}_L},$$

as needed.

The last step (11.9) follows from the assumption on \mathfrak{m} . \square

Example 11.5.10: Recall how we realized the class group and narrow class group as ray class groups in Example 11.3.4. We now realize them as quotients of the idele class group.

Take \mathfrak{m} to be 1. Then the bottom map gives an isomorphism

$$\mathbb{I}_K/K^\times \mathbb{U}_K \cong C_K$$

where C_K is just the class group of K . This realizes the class group of K as a quotient of the idele class group.

In general, for any modulus \mathfrak{m} ,

$$\mathbb{I}_K/K^\times \mathbb{U}_K(1, \mathfrak{m}) \cong \mathbb{I}_K(1, \mathfrak{m})/K(1, \mathfrak{m})\mathbb{U}_K(1, \mathfrak{m}) \cong C_K(\mathfrak{m}).$$

This realizes the ray class group modulo \mathfrak{m} as a quotient of the idele class group.

In particular, $\mathfrak{m} = 1$ was the case above. Taking $\mathfrak{m} = \prod_{v \text{ real}} v$, $P_K(1, \mathfrak{m})$ is the group of principal ideals generated by totally positive elements (also written P_K^+) and $\mathbb{U}_K(1, \mathfrak{m}) = \prod_{v \text{ real}} \mathbb{R}_{>0} \times \prod_v \mathcal{O}_v^\times$. This realizes the narrow class group of K as a quotient of the idele class group.

Remark 11.5.11: The condition on \mathfrak{m} in Proposition 11.5.9 was that $\mathbb{U}_K(1, \mathfrak{m}) \subseteq \text{Nm}_{L/K}(\mathbb{I}_L)$. We claim that we can always choose such \mathfrak{m} , such that $S(\mathfrak{m})$ consists of exactly the primes ramifying in L/K .

The condition $\mathbb{U}_K(1, \mathfrak{m}) \subseteq \text{Nm}_{L/K}(\mathbb{I}_L)$ says that $\mathcal{O}_v^\times \subseteq \text{Nm}_{L^v/K_v}(L^v)$ for all $v \nmid \mathfrak{m}$ and $I(\mathfrak{m})_v \subseteq \text{Nm}_{L^v/K_v}(L^v)$ for all $v \mid \mathfrak{m}$. Now note the following.

1. If L/K is unramified at v , i.e. L^v/K_v is unramified, then

$$\text{Nm}_{L^v/K_v}(L^{v\times}) = \pi_v^{[L^v:K_v]\mathbb{Z}} \mathcal{O}_v^\times \supseteq \mathcal{O}_v^\times.$$

This is a consequence of local class field theory (Example 14.5.1).

2. $\text{Nm}_{L^v/K_v}(L^v)$ is an open subgroup of K_v (this is the easy direction in Theorem 11.2.3) and $U_v^{(n)} := 1 + \pi_v^n \mathcal{O}_v$ is a neighborhood base of 1 in K_v .

By item 1, \mathfrak{m} doesn't need to include the places where L/K is unramified, and by item 2, for all ramified v we can choose the power of v in \mathfrak{m} large enough to force $U_v^{(n)} \subseteq \text{Nm}_{L^v/K_v}(L^{v\times})$. Then we will have $\mathbb{U}_K(1, \mathfrak{m}) \subseteq \text{Nm}_{L/K}(\mathbb{I}_L)$.

6 Global reciprocity via ideles

We now state global reciprocity in terms of ideles.

Theorem 11.6.1 (Global reciprocity, ideles). *Given a finite abelian extension L/K , there is a unique continuous² homomorphism $\phi_{L/K}$ that is compatible with the local Artin maps, i.e. the following diagram commutes³:*

$$\begin{array}{ccc} \mathbb{I}_K & \xrightarrow{\phi_{L/K}} & G(L/K) \\ \uparrow i_v & & \uparrow \\ K_v^\times & \xrightarrow{\phi_v} & G(L^v/K_v). \end{array}$$

Moreover, $\phi_{L/K}$ satisfies the following properties.

1. (Isomorphism) For every finite abelian extension L/K , ϕ_K defines an isomorphism

$$\phi_{L/K} : \mathbf{C}_K / \text{Nm}_{L/K}(\mathbf{C}_L) = \mathbb{I}_K / (K^\times \cdot \text{Nm}_{L/K}(\mathbb{I}_L)) \xrightarrow{\cong} G(L/K).$$

2. (Compatibility over all extensions) For $L \subseteq M$, L, M both finite abelian extensions of K , the following commutes:

$$\begin{array}{ccc} & G(M/K) & \\ \nearrow \phi_{M/K} & & \downarrow p_L \\ \mathbb{I}_K & \xrightarrow{\phi_{L/K}} & G(L/K) \end{array}$$

² $G(L/K)$ is given the discrete topology.

³This implies that if $v = v_{\mathfrak{p}}$ is unramified in L , then $\phi_{L/K}(i_v(\pi_v)) = \text{Frob}_{L/K}(\mathfrak{p})$. Global reciprocity is sometimes phrased in this way, though the phrasing using the local map gives a bit more information.

Thus we can define $\phi_K := \varprojlim_{L/K \text{ abelian}} \phi_{L/K}$ as a map $\mathbb{I}_K \rightarrow G(K^{ab}/K)$.

3. (Compatibility with norm map) ϕ_K is a continuous homomorphism $\mathbb{I}_K \rightarrow G(K^{ab}/K)$, and the following commutes.

$$\begin{array}{ccc} \mathbb{I}_L & \xrightarrow{\phi_L} & G(L^{ab}/L) \\ \downarrow \text{Nm}_{L/K} & & \downarrow \cdot|_{K^{ab}} \\ \mathbb{I}_K & \xrightarrow{\phi_K} & G(K^{ab}/K) \end{array}$$

Note that in the local reciprocity theorem 11.2.1, the “compatibility over all extensions” was automatic when we declared the existence of $\phi_K : K^\times \rightarrow G(K^{ab}/K)$. We stated the global reciprocity theorem a bit differently, in the above fashion for easy comparison with global reciprocity in terms of ideals 11.4.1.

Remark 11.6.2: Uniqueness and existence of $\phi_{L/K}$ is easy, and parts 2 and 3 are easy given the existence of ϕ_L . The crux of the theorem is again part 1.

For uniqueness, note that the $\phi_{L/K}$ is determined by its action on K_v^\times , since for $\mathbf{x} = (x_v)$, we must have

$$\phi_{L/K}(\mathbf{x}) = \prod_{v \in V_K} \phi_v(x_v).$$

(The product is Cauchy in the topology of \mathbb{I}_K .) This does define a continuous map on \mathbb{I}_K because $\phi_v(x_v) = 1$ whenever $x_v \in \mathcal{O}_v^\times$ and v is unramified, and this happens for all but finitely many v .

Parts 2 and 3 follow from the corresponding statements for local class field theory (see Theorem 11.2.1 and the paragraph above this remark), by how ϕ is defined to be compatible with the local maps.

The idele version of global reciprocity allows us to recast the Existence Theorem 11.4.4 in a format more similar to the Existence Theorem in 11.2.3.

Theorem 11.6.3 (Existence theorem). *For every subgroup $N \subseteq \mathbf{C}_K$ of finite index, there exists a unique abelian extension L/K such that $\text{Nm}_{L/K} \mathbf{C}_L = N$.*

Combining the two theorems, we can recast the bijective correspondence in Theorem 11.4.7 in a format more similar to local class field theory 11.2.5.

Theorem 11.6.4. *The map $L \mapsto \text{Nm}_{L/K}(\mathbf{C}_L)$ is an inclusion-reversing bijection between the set of finite abelian extensions of K and the open subgroups of finite index in \mathbf{C}_K , that switches intersections and products:*

$$\begin{aligned} L \subseteq M &\iff \text{Nm}_{L/K}(\mathbf{C}_L) \supseteq \text{Nm}_{M/K}(\mathbf{C}_M) \\ \text{Nm}_{L_1 L_2 / K}(\mathbf{C}_{L_1 L_2}) &= \text{Nm}_{L_1 / K}(\mathbf{C}_{L_1}) \cap \text{Nm}_{L_2 / K}(\mathbf{C}_{L_2}) \\ \text{Nm}_{L_1 \cap L_2 / K}(\mathbf{C}_{L_1 \cap L_2}) &= \text{Nm}_{L_1 / K}(\mathbf{C}_{L_1}) \cdot \text{Nm}_{L_2 / K}(\mathbf{C}_{L_2}). \end{aligned}$$

Similar to Theorem 11.2.4, we have the following topological isomorphism for global class field theory.

Theorem 11.6.5 (Topological isomorphism for GCFT). *Let K be a number field. Let*

$$(K_\infty^\times)^0 := \prod_{v \text{ real}} \mathbb{R}_{>0} \times \prod_{v \text{ complex}} \mathbb{C} \times \prod_{v \in V_K^0} 1.$$

The Artin map ϕ_K is surjective and induces a topological isomorphism

$$\mathbb{I}_K / \overline{K^\times (K_\infty^\times)^0} \cong G(K^{ab}/K).$$

6.1 Connecting the two formulations

We now show that the two formulations of global class field theory are equivalent, in the following sense.

Theorem 11.6.6. *We have the following implications.*

1. (Global reciprocity, ideles \implies ideals) *If Theorem 11.6.1(1) holds for a given L/K , then Theorem 11.4.1(1) holds for L/K . If Theorem 11.6.1 holds for all L/K over a specified basefield (e.g. \mathbb{Q}), then Theorem 11.4.1 holds for all such L/K .*
2. (Global reciprocity, ideals \implies (ideles) $-\varepsilon$) *If Theorem 11.6.1(1)-(2) holds for a fixed K and a family $\{L/K\}$ such that the compositum of the L^v contains K_v^{ur} for every finite place v , then Theorem 11.6.1(1)-(2) holds for the same K and $\{L/K\}$, except that the resulting map $\phi_{L/K}$ may not be compatible with ϕ_v when v is archimedean.*
3. (Global existence) *Given Theorem 11.6.1, Theorems 11.4.4 and 11.6.3 are equivalent.*
4. (Bijective correspondence) *Given Theorem 11.6.1, Theorems 11.4.7 and 11.6.4 are equivalent.*

Proof. For parts 1 and 2, we note that by Proposition 11.5.9,

$$\mathbf{C}_K / \text{Nm}_{L/K} \mathbf{C}_L = \mathbb{I}_K / K^\times \text{Nm}_{L/K} \mathbb{I}_L \cong I_K^S / P_K(1, \mathfrak{m}) \text{Nm}_{L/K}(I_L^S), \quad (11.10)$$

where by Remark 11.5.11, we can choose \mathfrak{m} to some modulus containing only ramified primes, and $S = S(\mathfrak{m})$. Thus any one of the dotted isomorphisms below gives the other isomorphism.

$$\begin{array}{ccc}
 \mathbb{I}_K / K^\times \text{Nm}_{L/K}(\mathbb{I}_L) & & \\
 \downarrow p \cong & \begin{array}{c} \nearrow \phi_{L/K} \cong \\ \searrow \psi_{L/K} \cong \end{array} & G(L/K) \\
 I_K^S / P_K(1, \mathfrak{m}) \text{Nm}_{L/K}(I_L^S) & &
 \end{array} \quad (11.11)$$

For part 1, given $\phi_{L/K}$, we define $\psi_{L/K}$ with the above diagram. Then, supposing \mathfrak{p} corresponds to the uniformizer $\pi_v \in K_{\mathfrak{p}}$,

$$\psi_{L/K}(\mathfrak{p}) = \psi_{L/K}(p(i(\pi_v))) = \phi_{L/K}(i(\pi_v)) = \phi_v(\pi_v) = \text{Frob}_{L^v/K_v}((\pi_v)) = \text{Frob}_{L/K}(\mathfrak{p}),$$

as needed. Part 2 is a more complicated; we'll give the proof below after a lemma. The “ $-\varepsilon$ ” comes from the fact that the formulation in Theorem 11.6.1 says nothing about archimedean primes.

Parts 3 and 4 now result directly from the fact that (11.10) gives a bijective correspondence between subgroups of two groups. \square

Lemma 11.6.7. *Suppose that K is a nonarchimedean local field, K^{ur} is the maximal abelian unramified extension of K , and L is an abelian extension containing K^{ur} . Let $f : K^{\times} \rightarrow G(L/K)$ be a homomorphism satisfying (1) and either (2) or (2)':*

1. *The composition $K^{\times} \xrightarrow{f} G(L/K) \rightarrow G(K^{ur}/K)$ takes α to $\text{Frob}_{K^{ur}/K}(\pi)^{v(\alpha)}$.*

2. *For any uniformizer $\pi \in K$, $f(\pi)|_{K_{\pi}} = 1$, where*

$$K_{\pi} := L^{\phi_K(\pi)}.$$

2'. *For any finite subextension K'/K of K_{π} , we have*

$$f(\text{Nm}_{K'/K}(K'^{\times}))|_{K'} = \{1\}.$$

Then f equals the reciprocity map ϕ_K .

For the proof, see Section 8.1.

Proof of Theorem 11.6.6, Part 2. Given $\psi_{L/K}$ we define $\phi_{L/K}$ using (11.11). The $\psi_{L/K}$ are compatible by Remark (11.4.2), so the $\phi_{L/K}$ are compatible (details omitted) and we can define $\phi_K = \varprojlim_{L/K} \phi_{L/K}$ where the limit is over L/K in the given family. Let L' be the compositum of the fields L .

We check the hypotheses 1 and 2' of Lemma 11.6.7. Let

$$f_v = \phi_K \circ i_v : K_v^{\times} \rightarrow G(L^v/K_v).$$

Item 1 is clear as (11.11) gives letting $v = v_{\mathfrak{p}}$, we have

$$\phi_K(i_v(\alpha))|_{K_v^{ur}} = \psi_K(\mathfrak{p}^{v(\alpha)})|_{K_v^{ur}} = \text{Frob}_{K_v^{ur}/K_v}(\alpha)^{v(\alpha)}.$$

Item 2' follows from part 3 of Theorem 11.4.1 applied to K'/K (see Remark 11.4.2): we get $\psi_{L/K}(\text{Nm}_{K'/K}(I_{K'}^S))|_{K'} = 1$ which translates into $\phi_K(i_v(\text{Nm}_{K'_v/K_v}(K_v'^{\times})))|_{K'_v} = 1$. Thus $f_v = \phi_v$ for all finite places, as needed. \square

We have proved the ideal version of global class field theory for cyclotomic extensions of \mathbb{Q} . Our plan of attack will be to show transfer this to the idele version for cyclotomic extension of \mathbb{Q} , then work on proving the idele version. Then we will be done by Theorem 11.6.6.

7 Kronecker-Weber Theorem

As a first application of class field theory, we explicitly describe the maximal abelian extensions of \mathbb{Q}_p and \mathbb{Q} .

Theorem 11.7.1 (Local Kronecker-Weber theorem). *Every abelian extension of \mathbb{Q}_p is included in a cyclotomic extension, i.e. an extension $\mathbb{Q}_p(\zeta_n)$, ζ_n a primitive n th root of unity, for some n . In other words,*

$$\mathbb{Q}_p^{ab} = \mathbb{Q}_p(\zeta_n \mid n \in \mathbb{N}).$$

Theorem 11.7.2 (Kronecker-Weber theorem). *Every abelian extension of \mathbb{Q} is included in a cyclotomic extension $\mathbb{Q}(\zeta_n)$. In other words,*

$$\mathbb{Q}^{ab} = \mathbb{Q}(\zeta_n \mid n \in \mathbb{N}).$$

Proof of Theorem 11.7.1. Consider $\mathbb{Q}_p(\zeta_k)$ where $p \nmid k$. Let U denote the group of units. As $\mathbb{Q}_p(\zeta_k)$ is unramified, local class field theory tells us

$$\mathrm{Nm}_{\mathbb{Q}_p(\zeta_k)/\mathbb{Q}_p}(\mathbb{Q}_p(\zeta_k)^\times) \cong \pi^{[\mathbb{Q}_p(\zeta_k):\mathbb{Q}_p]\mathbb{Z}}U.$$

Consider $\mathbb{Q}_p(\zeta_{p^m})$, which is totally ramified of degree $p^{m-1}(p-1)$ over \mathbb{Q}_p . Local reciprocity gives

$$\mathbb{Q}_p^\times / \mathrm{Nm}_{\mathbb{Q}_p(\zeta_{p^m})/\mathbb{Q}_p}(\mathbb{Q}_p(\zeta_{p^m})^\times) \xrightarrow{\cong} G(\mathbb{Q}_p(\zeta_{p^m})/\mathbb{Q}_p).$$

Thus both sides have the same order, $p^{m-1}(p-1)$, and we must have

$$\mathrm{Nm}_{\mathbb{Q}_p(\zeta_{p^m})/\mathbb{Q}_p}(\mathbb{Q}_p(\zeta_{p^m})^\times) = U^{(m)} := p^{\mathbb{Z}}(1 + (p^m)).$$

Suppose L/\mathbb{Q}_p is an abelian extension. Its corresponding norm group N is open of finite index in \mathbb{Q}_p , so contains

$$p^{n\mathbb{Z}}(1 + (p^m))$$

for some n, m . Choosing k large enough we may suppose $n \mid [\mathbb{Q}_p(\zeta_k) : \mathbb{Q}_p]$. Then using Theorem 11.2.5⁴,

$$N \supseteq \mathrm{Nm}(\mathbb{Q}_p(\zeta_n)^\times) \cap \mathrm{Nm}(\mathbb{Q}_p(\zeta_{p^m})^\times) = \mathrm{Nm}(\mathbb{Q}_p(\zeta_{np^m})^\times).$$

By Theorem 11.2.5, we get that $\mathbb{Q}_p(\zeta_{np^m}) \supseteq L$. □

Proof of Theorem 11.7.2. Given an abelian extension K/\mathbb{Q} , choose a modulus \mathfrak{m} so that the Artin map is defined. Every modulus for \mathbb{Q} divides $\infty(n)$ for some integer n . The ray class field of $\infty(n)$ is $\mathbb{Q}(\zeta_n)$. If \mathfrak{m} divides $\infty(n)$, then K is contained in $\mathbb{Q}(\zeta_n)$. Hence the maximal abelian extension is the union of all the $\mathbb{Q}(\zeta_n)$. □

We can similarly ask how to characterize abelian extensions of other number fields K . This is Hilbert's Twelfth Problem and Kronecker's Jugendtraum. Note that another way to phrase this theorem is the following:

⁴omitting the subscripts on norms to avoid clutter

1. \mathbb{Q}^{ab} is generated by the *torsion points* of \mathbb{Q}^\times under multiplication.
2. Let $f(z) = e^{2\pi iz}$. Then \mathbb{Q}^{ab} is generated by $f(\mathbb{Q})$:

$$\mathbb{Q}^{\text{ab}} = \mathbb{Q}(f(\mathbb{Q})).$$

We can ask: for given K , can we get K^{ab} by adjoining torsion points of some algebraic variety, and does there exist a nice function $g(z)$ parameterizing this variety, so that

$$K^{\text{ab}} \approx K(g(K))?$$

It turns out that the answer is affirmative for quadratic extensions: roughly speaking, the maximal abelian extension is generated by torsion points of elliptic curves with complex multiplication. We will give a complete solution to this problem in Chapter 17.

8 Problems

- 1.1 Why can't we define $\text{Frob}_{\mathfrak{p}} \in G(L/K)$ when \mathfrak{p} is a prime in K that is ramified in L ?
- 1.2 Fix $n \in \mathbb{N}$.
 - (a) For which primes $p \in \mathbb{Z}$ does (p) split completely in $\mathbb{Z}[\zeta_n + \zeta_n^{-1}]$? (Be careful with $p = 2$.)
 - (b) Show that the ray class field of (n) is $\mathbb{Q}(\zeta_n + \zeta_n^{-1})$.
- 1.3 (IberoAmerican Olympiad for University Students, 2010/6) Prove that, for all integers $a > 1$, the prime divisors of $5a^4 - 5a^2 + 1$ have the form $20k \pm 1$.
- 1.4 Consider the field extension $\mathbb{Q}(\sqrt[3]{d}, \zeta_3)/\mathbb{Q}$ where $d \in \mathbb{Z}$ is not a perfect cube. Let p be a prime relatively prime to $3d$. Prove that a prime p splits into n factors in $\mathbb{Q}(\sqrt[3]{d}, \zeta_3)$, where

$$n = \begin{cases} 2, & p \equiv 1 \pmod{3} \text{ and } d \text{ is a cube modulo } p \\ 3, & p \equiv 1 \pmod{3} \text{ and } d \text{ is not a cube modulo } p \\ 6, & p \equiv 2 \pmod{3}. \end{cases}$$

- 2.1 Recall that $G(\overline{K}/K)$ has profinite (Krull) topology. Topologically $W(\overline{K}/K)$ is a \mathbb{Z} -disjoint union of $G(\overline{K}/K)_0$ -cosets $G(\overline{K}/K)_0\sigma_n$, where σ_n is any lift of Frob_q^n , $n \in \mathbb{Z}$, where each $G(\overline{K}/K)_0\sigma_n$ is given the same topology as the profinite topology on $G(\overline{K}/K)_0$ via translation by σ_n .
 - (a) Show that the natural inclusion $\iota : W(\overline{K}/K) \rightarrow G(\overline{K}/K)$ is continuous and has dense image.
 - (b) Show that ι is not a topological isomorphism onto $\iota(W(\overline{K}/K))$, where the latter is equipped with the topology induced by that of $G(\overline{K}/K)$.

- 4.1 There is a correspondence between quadratic characters (homomorphisms $\chi : (\mathbb{Z}/m)^\times \rightarrow \mathbb{C}$ such that $\chi^2 = 1, \chi \neq 1$, where characters are naturally identified by restriction) and quadratic extensions,
- (a) Prove this without using the statement of class field theory.
 - (b) How does CFT give the result?
 - (c) Using CFT for quadratic extensions, prove quadratic reciprocity.
- 4.2 Characterize all quadratic extensions K/\mathbb{Q} that are contained in a $\mathbb{Z}/4$ -extension. (Ben Blum-Smith, from <http://math.stackexchange.com/questions/596195/conceptual-reason-why-a-quadratic-field-has-1-as-a-norm-if-and-only-if>)

Chapter 12

Group homology and cohomology

In this chapter we introduce the theory of group homology and cohomology. In the next chapter we'll specialize to the case of Galois groups, and then we'll use Galois cohomology to prove the theorems of class field theory. Some results in this chapter will be given without proof; for detailed proofs see Rotman [Rot09]. We assume knowledge of some basic terminology and facts from category theory and commutative algebra (covariant and contravariant functors, natural transformations, left and right exactness).

The idea of homology and cohomology—used in many different areas of mathematics—is that after applying a functor, a short exact sequence of modules may no longer be exact. Instead, we get the *long exact sequence in (co)homology*, with the (co)homology groups measuring the deviation from exactness.

Exactly what functors are we applying? In group cohomology (Section 6), we apply $\text{Hom}_G(\mathbb{Z}, \bullet)$, turning a short exact sequence of G -modules

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

into

$$0 \rightarrow A^G \rightarrow B^G \rightarrow C^G \rightarrow H^1(G, A) \rightarrow \cdots \tag{12.1}$$

where A^G is the submodule of A fixed by G . In the next chapter we will take A, B, C to be a multiplicative or additive subgroup of a field L , and $G = G(L/K)$. Then A^G is just $A \cap K$. Thus we see that the sequence (12.1) gives information about the relationship between a field K and an extension field. For example, in Kummer Theory 2, we take $C = L^{\times n}$; then $C^G = L^{\times n} \cap K$, and we can characterize $G(L/K)$ and hence L/K in terms of the n th powers of L appearing in K . This is representative a general trend in class field theory: characterize extensions of K in terms of information intrinsic to K .

We also get a sequence in group homology (Section 8), and we can splice the sequences for homology and cohomology together to get the Tate groups (Section 9). Norm groups will make their appearance here—which is how, in class field theory, we get a correspondence between norm groups and field extensions.

Finally, we assemble a toolbox of other constructions from group cohomology and homology, including cup products (Section 10), changes of group (Section 11), the corestriction

map (Section 11.5), results on cyclic groups and the Herbrand quotient (Section 12), and Tate’s theorem (Section 13). We include generalizations of cohomology to profinite groups (Section 14) and nonabelian groups (Section 15).

1 Projectives and injectives

Let \mathcal{A} be an abelian category.¹ The reader unfamiliar with category theory may assume that \mathcal{A} is the class of R -modules, since we will be primarily working with modules throughout.

Definition 12.1.1: Let \mathcal{A} be an abelian category.

1. An object $P \in \mathcal{A}$ is **projective** if for every surjection $p : M \twoheadrightarrow N$ and morphism $f : P \rightarrow N$, there exists a unique morphism $g : P \rightarrow M$ such that $f = p \circ g$:

$$\begin{array}{ccc} & & P \\ & \swarrow g & \downarrow f \\ M & \xrightarrow{p} & N \end{array}$$

Equivalently, $\text{Hom}(P, \bullet)$ is exact (or equivalently, right exact as it is always left exact).²

2. An object $I \in \mathcal{A}$ is **injective** if for every injection $i : M \hookrightarrow N$ and morphism $f : M \rightarrow I$, there exists a unique morphism $g : N \rightarrow I$, such that $f = g \circ i$:

$$\begin{array}{ccc} M & \xhookrightarrow{i} & N \\ \downarrow f & & \swarrow g \\ I & & \end{array}$$

Equivalently, $\text{Hom}(\bullet, I)$ is exact (or equivalently, just right exact).

Example 12.1.2: A free R -module (a direct sum of copies of R) is projective.

Definition 12.1.3: An abelian category $\mathcal{A} \dots$

1. **has enough injectives** if for every object $A \in \mathcal{A}$ there exists an injective object E with a monic (injective) morphism $A \hookrightarrow E$.
2. **has enough projectives** if for every object $A \in \mathcal{A}$ there exists a projective object P with an epic (surjective) morphism $P \twoheadrightarrow A$.

¹A category is an **abelian category** if it is an additive category such that every morphism has a kernel and cokernel, every monomorphism (injection) is a kernel, and every epimorphism (surjection) is a cokernel.

²The diagram is equivalent to saying that if $p : M \twoheadrightarrow N$ is surjective, then so is the map $\text{Hom}(P, M) \xrightarrow{\text{Hom}(\bullet, p)} \text{Hom}(P, N)$, i.e. $\text{Hom}(P, \bullet)$ is right exact.

Definition 12.1.4: A **projective resolution** of A is an exact sequence

$$\mathbf{P} : \dots \rightarrow P_2 \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0 \xrightarrow{\varepsilon} A \rightarrow 0$$

where each P_n is projective.

An **injective resolution** of A is an exact sequence

$$\mathbf{E} : 0 \rightarrow A \xrightarrow{\eta} E^0 \xrightarrow{d^0} E^1 \xrightarrow{d^1} E^2 \rightarrow \dots$$

where each E^n is injective.

Proposition 12.1.5: If \mathcal{A} is an abelian category with enough projectives (injectives), then every object has a projective (injective) resolution. In particular, every R -module has a projective (injective) resolution.

Proof. Build the resolution step-by-step. See Rotman [Rot09], Proposition 6.2-5. For the second part, note that the category of R -modules has enough projectives and enough injectives. \square

2 Complexes

Definition 12.2.1: A **complex** in an abelian category (for example, the category of R -modules or abelian groups) is a sequence of morphisms

$$\mathbf{C} : \dots \rightarrow C_{n+1} \xrightarrow{d_{n+1}} C_n \xrightarrow{d_n} C_{n-1} \rightarrow \dots$$

such that the composition of any two adjacent morphisms is 0:

$$d_n d_{n+1} = 0.$$

We often work with complexes only going off to the left or right (positive and negative complexes, respectively), and label them

$$\begin{array}{ccccccc} \dots & \rightarrow & C_n & \xrightarrow{d_n} & C_{n-1} & \rightarrow & \dots \rightarrow C_0 \rightarrow 0 \\ 0 & \rightarrow & C^0 & \rightarrow & \dots \rightarrow C^{n-1} & \xrightarrow{d^{n-1}} & C^n \rightarrow \dots \end{array}$$

We will want to work with complexes like they are single objects.

Theorem 12.2.2. *The class of complexes in \mathcal{A} can be made into an abelian category, $\text{Comp}(\mathcal{A})$ as follows: The objects are the complexes and the morphisms are **chain maps** $f = (f_n) : \mathbf{C} \rightarrow \mathbf{C}'$, i.e. a sequence of maps making the following commute.*

$$\begin{array}{ccccccc} \longrightarrow & C_{n+1} & \xrightarrow{d_{n+1}} & C_n & \xrightarrow{d_n} & C_{n-1} & \xrightarrow{d_{n-1}} \longrightarrow \\ & \downarrow f_{n+1} & & \downarrow f_n & & \downarrow f_{n-1} & \\ \longrightarrow & C'_{n+1} & \xrightarrow{d'_{n+1}} & C'_n & \xrightarrow{d'_n} & C'_{n-1} & \xrightarrow{d'_{n-1}} \longrightarrow \end{array}$$

Proof. See Rotman [Rot09], Proposition 5.100. □

We will be interested in cohomology and homology modules associated to chain complexes. For this, we have the following notion of what it means for chain maps to be “the same” (See Theorem 12.3.2).

Definition 12.2.3: Two chain maps $f, g : \mathbf{C} \rightarrow \mathbf{C}'$ are **homotopic** if there exist a family of morphisms $s_n : C_n \rightarrow C'_{n+1}$ such that

$$f_n - g_n = d'_{n+1}s_n + s_{n-1}d_n.$$

In Section 4 we will define the homology modules and cohomology modules from projective and injective resolutions. To show this does not depend on the choice of projective or injective resolution, we need the following theorem.

Theorem 12.2.4 (Comparison Theorem). *Let \mathcal{A} be an abelian category, and suppose we have two complexes $\mathbf{C} : \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow A \rightarrow 0$ and $\mathbf{C}' : \cdots \rightarrow P'_1 \rightarrow P'_0 \rightarrow A' \rightarrow 0$ and a map $g : A \rightarrow A'$. Then there exists a chain map f extending g :*

$$\begin{array}{ccccccc} \cdots & \longrightarrow & P_1 & \longrightarrow & P_0 & \longrightarrow & A \longrightarrow 0 \\ & & \downarrow f_1 & & \downarrow f_0 & & \downarrow g \\ \cdots & \longrightarrow & P'_1 & \longrightarrow & P'_0 & \longrightarrow & A' \longrightarrow 0. \end{array}$$

Moreover, f is unique up to homotopy.

The same is true of complexes going off to the right (reverse the arrows above).

Proof. Rotman [Rot09], Theorem 6.16. □

3 Homology and cohomology

Definition 12.3.1: Given a complex \mathbf{C} , define

$$\begin{aligned} Z_n(\mathbf{C}) &= \ker(d_n) \\ B_n(\mathbf{C}) &= \operatorname{im}(d_{n+1}) \\ H_n(\mathbf{C}) &= Z_n(\mathbf{C})/B_n(\mathbf{C}). \end{aligned}$$

H_n is called the n th **homology module**. For upper indexing, we let $Z^n(\mathbf{C}) = \ker(d^n)$, $B^n(\mathbf{C}) = \operatorname{im}(d^{n-1})$, and $H^n(\mathbf{C}) = Z^n(\mathbf{C})/B^n(\mathbf{C})$, and call H^n the n th **cohomology module**.

Think of H_n as measuring how far the complex is from being exact at C_n .

Theorem 12.3.2. *Let \mathcal{A} be an abelian category. For every integer n , H_n is an additive functor from $\operatorname{Comp}(\mathcal{A}) \rightarrow \mathcal{A}$. Moreover, homotopic chain maps induce the same map in homology.*

Proof. See Rotman [Rot09], Proposition 6.8. □

Theorem 12.3.3 (Long exact sequence). *A short exact sequence of chain complexes*

$$0 \longrightarrow \mathbf{C}' \xrightarrow{i} \mathbf{C} \xrightarrow{p} \mathbf{C}'' \longrightarrow 0$$


induces a long exact sequence of homology modules

$$\cdots \longrightarrow H'_n \xrightarrow{i_n} H_n \xrightarrow{p_n} H''_n \xrightarrow{\partial_n} H'_{n-1} \longrightarrow \cdots$$

The map ∂_n is defined by

$$\partial_n[c''_n] = [i_{n-1}^{-1}d_{n-1}p_n^{-1}c''_n] \in C'_{n-1}.$$

Proof. Let $H_n = H_n(\mathbf{C})$, $B_n = \text{im}(d^{n+1})$, and $Z_n = \text{ker}(d^n)$ for the complex \mathbf{C} , and define H'_n , H''_n , and so forth similarly. By the Snake Lemma, the gray sequence below is exact.



Note that the connecting homomorphism is exactly that in the Snake Lemma. □

4 Derived functors

4.1 Right derived functors and Ext

4.1.1 Covariant case

Given an injective resolution of B ,

$$E^B : 0 \rightarrow B \xrightarrow{\eta} E^0 \xrightarrow{d^0} E^1 \xrightarrow{d^1} E^2 \xrightarrow{d^2} \cdots,$$

applying a (covariant) functor T gives (after deleting TB)

$$0 \rightarrow TE^0 \xrightarrow{Td^0} TE^1 \xrightarrow{Td^1} TE_2 \xrightarrow{Td^2} \dots \quad (12.2)$$

We will primarily be concerned with the case where $T = \text{Hom}_R(A, \bullet)$, so the above becomes

$$0 \rightarrow \text{Hom}(A, E^0) \xrightarrow{\text{Hom}(A, d^0)} \text{Hom}(A, E^1) \xrightarrow{\text{Hom}(A, d^1)} \text{Hom}(A, E^2) \xrightarrow{\text{Hom}(A, d^2)} \dots \quad (12.3)$$

Definition 12.4.1: Let T be a covariant functor. The n th (**covariant**) **right derived functor** of T is

$$(R^n T)B := H^n(TE^B) = \frac{\ker(Td^n)}{\text{im}(Td^{n-1})},$$

i.e. it is the n th cohomology module of (12.2).

For a R -module E , define

$$\text{Ext}_R^n(A, B) := (R^n \text{Hom}_R(A, \bullet))B = H^n(\text{Hom}_R(A, E^B)),$$

i.e. it is the n th cohomology module of (12.3).

Here d^{-1} is the trivial map $0 \rightarrow E^0$. We need to show that this definition does not depend on the injective resolution chosen.

Proof of well-definedness. Suppose we have two injective resolutions of B :

$$\begin{array}{ccccccc} 0 & \longrightarrow & B & \xrightarrow{\eta} & E^0 & \xrightarrow{d^0} & E^1 & \xrightarrow{d^1} & \dots \\ & & \parallel & & \vdots & & \vdots & & \\ 0 & \longrightarrow & B & \xrightarrow{\eta'} & E'^0 & \xrightarrow{d'^0} & E'^1 & \xrightarrow{d'^1} & \dots \end{array}$$

Let $(R^n T)B = \frac{\ker(Td^n)}{\text{im}(Td^{n-1})}$ and $(R'^n T)B = \frac{\ker(Td'^n)}{\text{im}(Td'^{n-1})}$.

By the Comparison Theorem 12.2.4, there is a unique chain map f between the two resolutions, up to homotopy (the dotted lines above). Apply T to this diagram to get a chain map $Tf_n : TE^n \rightarrow TE'^n$. As H_n is a functor by Theorem 12.3.2, Tf induces a map on the cohomology modules $(R^n T)B \rightarrow (R'^n T)B$. Since we can construct a chain map g from the second to the first resolution as well, $(R^n T)B \rightarrow (R'^n T)B$ must be an isomorphism.

For the details, see [Rot09], Proposition 6.20. (The argument there is written for left derived functors, but the idea is the same.) \square

4.1.2 Contravariant case

We can define a companion functor ext_R^n that is contravariant instead of covariant. Given an projective resolution of A

$$P_A : \quad \dots \xrightarrow{d_2} P_2 \xrightarrow{d_1} P_1 \xrightarrow{d_0} P_0 \xrightarrow{\varepsilon} A \rightarrow 0,$$

applying a contravariant functor T gives

$$0 \xrightarrow{Td_{-1}=0} TP_0 \xrightarrow{Td_0} TP_1 \xrightarrow{Td_1} TP_2 \xrightarrow{Td_2} \dots \quad (12.4)$$

To define ext, let $T = \text{Hom}_R(\bullet, B)$.

Definition 12.4.2: Let T be a contravariant functor. The n th (**contravariant**) **right derived functor** of T is

$$(R^n T)A := H^n(TP_A) = \frac{\ker(Td^n)}{\text{im}(Td^{n-1})},$$

i.e. it is the n th cohomology module of (12.4).

For R -modules A, B , define

$$\text{ext}_R^n(A, B) := (R^n \text{Hom}_R(\bullet, B))A = H^n(\text{Hom}_R(P_A, B))$$

Theorem 12.4.3. For R -modules A, B ,

$$\text{Ext}_R^n(A, B) = \text{ext}_R^n(A, B).$$

This theorem says that we have two choices when we need to calculate $\text{Ext}_R^n(A, B)$, namely,

1. Find a injective resolution of B and apply $\text{Hom}(A, \bullet)$ (the Ext perspective), or
2. Find a projective (e.g. free) resolution of A and apply $\text{Hom}(\bullet, B)$ (the ext perspective).

Proof. See Rotman [Rot09], Theorem 6.67. □

4.2 Left derived functors and Tor

Next we define left derived functors and Tor analogously. Given a projective resolution of A

$$P_A : \dots \xrightarrow{d_2} P_2 \xrightarrow{d_1} P_1 \xrightarrow{d_0} P_0 \xrightarrow{\varepsilon} A \rightarrow 0,$$

applying a covariant functor T gives

$$\dots \xrightarrow{Td_2} TP_2 \xrightarrow{Td_1} TP_1 \xrightarrow{Td_0} TP_0 \xrightarrow{Td_{-1}} 0.$$

To define Tor, let $T = \bullet \otimes_R B$.

Definition 12.4.4: The n th **left derived functor** of T is

$$(L_n T)B := H_n(TP_A) = \frac{\ker(Td_{n-1})}{\text{im}(Td_n)}.$$

For A an R -module, define

$$\text{Tor}_n^R(A, B) := (L_n(\bullet \otimes_R B))A = H^n(P_A \otimes_R B)$$

$$\text{tor}_n^R(A, B) := (L_n(A \otimes_R \bullet))A = H^n(A \otimes_R P_B).$$

(Note $\text{Tor}_n^R(A, B) = \text{tor}_n^R(B, A)$.)

Note unlike the case with Ext, we need only consider covariant derived functors: Hom_R is contravariant in the first entry and covariant in the second, while \otimes_R is covariant in both entries. Similar to Theorem 12.4.3, we have the following.

Theorem 12.4.5. *For A, B R -modules,*

$$\text{Tor}_n^R(A, B) = \text{tor}_n^R(A, B).$$

Proof. See Rotman [Rot09], Theorem 6.32. □

4.3 Long exact sequences

The most important property of the derived functors is that they repair “loss of exactness” after applying the functor.

Theorem 12.4.6 (Long exact sequence). *Let $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ be a short exact sequence of G -modules.*

1. *Let T be a left exact covariant functor. Then there is a long exact sequence*

$$\begin{array}{ccccccc} 0 & \longrightarrow & (R^0T)A & \longrightarrow & (R^0T)B & \longrightarrow & (R^0T)C \xrightarrow{\partial^0} (R^1T)A \longrightarrow \dots \\ & & \parallel & & \parallel & & \parallel \\ & & TA & & TB & & TC \end{array}$$

2. *Let T be a right exact covariant functor. Then there is a long exact sequence*

$$\begin{array}{ccccccc} \dots & \longrightarrow & (L_1T)C & \xrightarrow{\partial_1} & (L_0T)A & \longrightarrow & (L_0T)B \longrightarrow (L_0T)C \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ & & TA & & TB & & TC \end{array}$$

The maps ∂^n are given by the snake lemma.

Proof. The long exact sequences exist by Theorem 12.3.3. (Note that the complexes only go off to the right/left in the two cases, respectively.) It remains to show the equalities. Take a projective resolution of A ,

$$\dots \xrightarrow{d_2} P_2 \xrightarrow{d_1} P_1 \xrightarrow{d_0} P_0 \xrightarrow{\varepsilon} A \rightarrow 0.$$

By right exactness of T , the following is exact:

$$TP_1 \xrightarrow{Td_1} TP_0 \xrightarrow{T\varepsilon} TA \longrightarrow 0.$$

Hence $(L_0T)A = TP_0/\text{im}(TP_1) \cong TA$.

The second part is similar. □

Corollary 12.4.7. *We have the long exact sequences*

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \text{Ext}_R^0(M, A) & \longrightarrow & \text{Ext}_R^0(M, B) & \longrightarrow & \text{Ext}_R^0(M, C) \xrightarrow{\partial^0} \text{Ext}_R^1(M, A) \longrightarrow \cdots \\
 & & \parallel & & \parallel & & \parallel \\
 & & \text{Hom}_R(M, A) & & \text{Hom}_R(M, B) & & \text{Hom}_R(M, C)
 \end{array}$$

and

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & \text{Tor}_1^R(C, M) & \xrightarrow{\partial_1} & \text{Tor}_0^R(A, M) & \longrightarrow & \text{Tor}_0^R(B, M) \longrightarrow \text{Tor}_0^R(C, M) \longrightarrow 0 \\
 & & & & \parallel & & \parallel \\
 & & & & M \otimes_R A & & M \otimes_R B \\
 & & & & & & \parallel \\
 & & & & & & M \otimes_R C
 \end{array}$$

Proof. $\text{Hom}_R(A, \bullet)$ is left exact and $\bullet \otimes_R B$ is right exact. □

Example 12.4.8: We have the following.

$$\begin{aligned}
 B \text{ injective} &\implies \text{Ext}_R^n(A, B) = 0 \text{ for all } A, n \geq 1 \\
 A \text{ projective} &\implies \text{Tor}_R^n(A, B) = 0 \text{ for all } B, n \geq 1.
 \end{aligned}$$

Indeed, recall that Ext is defined by taking an injective resolution of B and Tor is defined by taking a projective resolution of A , and in these cases we can take the trivial resolutions $0 \rightarrow B \rightarrow B \rightarrow 0$ and $0 \rightarrow A \rightarrow A \rightarrow 0$.

Example 12.4.9: Take $R = \mathbb{Z}$. Then a R -module is just an abelian group. Every group H has a free resolution of length 2:

$$0 \rightarrow F_1 \rightarrow F_0 \rightarrow H \rightarrow 0.$$

Thus $\text{ext}_{\mathbb{Z}}^n(H, G) = 0$ and $\text{Tor}_n^{\mathbb{Z}}(H, G) = 0$ for $n \geq 2$.

5 Homological and cohomological functors

This section is more abstract and may be skipped.

As we saw in Corollary 12.4.7 and Example 12.4.8, the key properties of Ext_R^n are roughly the following:

1. $\text{Ext}_R^n(A, B) = 0$ when B is injective and $n \geq 1$.
2. Short exact sequences give rise to long exact sequences.
3. In dimension 0, $\text{Ext}_R^0(A, B) = \text{Hom}_R(A, B)$.

We have a similar description for Tor_n^R .

We abstract the definition for Ext and Tor , by defining homological and cohomological functors. There are several reasons for doing this:

1. We want to talk about *natural transformations* between cohomological functors.
2. In the last section we showed the existence of Ext satisfying the above properties (and similarly for Tor). It turns out that these properties characterize it uniquely. Thus we can just “remember” these properties and forget the details of the construction.

There are similarly other (co)homological functors, and we sometimes want to show they are equal. To do this, it turns out we can just construct an isomorphism in dimension 0, and the rest works out by abstract nonsense. (See Theorem 12.5.2.)

Note in the above characterization of Ext we said $\text{Ext}_n^R(A, B) = 0$ for $n \geq 1$ when B is injective. This is useful because every R -module has an injective resolution. In general, though, we may want to work with a general class of objects, say χ (which in our case is the class of injective modules). The key property is that for every module A there is an injective module E and an injective morphism $A \rightarrow E$, i.e. the category of R -modules has enough injectives.

Definition 12.5.1: Let $(T^n : \mathcal{A} \rightarrow \mathcal{B})_{n \geq 0}$ be a set of additive functors on abelian categories, and let χ be a class of objects in \mathcal{A} . We say \mathcal{A} has **enough χ -objects** if every object in \mathcal{A} can be embedded in an object in χ .

Supposing \mathcal{A} has enough χ -objects, $(T^n)_{n \geq 0}$ is a **cohomological ∂ -functor** if the following hold.

1. $(T^n)_{n \geq 0}$ is **χ -coeffaceable**: $T^n(X) = 0$ for all $X \in \chi$ and $n \geq 1$.
2. For every short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ there is a long exact sequence

$$0 \rightarrow T^0(A) \rightarrow \cdots \rightarrow T^n(A) \rightarrow T^n(B) \rightarrow T^n(C) \xrightarrow{\partial^n} T^{n+1}(A) \rightarrow \cdots$$

such that the diagonal morphisms ∂^n are natural (with respect to maps between two short exact sequences).

A morphism of cohomological ∂ -functors is a natural transformation $\tau^n : T^n \rightarrow H^n$ commuting with the diagonal maps ∂^n .

There is a similar definition for effaceability and homological ∂ -functors. We can also consider $(T^n)_{n \in \mathbb{Z}}$, that is ∂ -functors extending infinitely in both directions, replacing the long exact sequence with an infinite exact sequence extending in both directions.

The following theorem gives existence and uniqueness of (co)homological ∂ -functors.

Theorem 12.5.2. 1. Suppose $\tau^0 : T^0 \rightarrow T'^0$ is a natural transformation of cohomological ∂ -functors in degree 0. Then there exists a unique morphism of cohomological ∂ -functors $\tau : T \rightarrow T'$ extending τ^0 .

2. Suppose $T^n, T^m : \mathcal{A} \rightarrow \mathcal{B}$ are two cohomological functors, and there is a natural isomorphism $T^0 \cong T'^0$. Then $T^n \cong T'^n$.

The same is true of homological ∂ -functors, and ∂ -functors extending in both directions.

Proof. See Rotman [Rot09], 6.35. □

For example, Ext_R is characterized completely by the 3 properties we gave: it is a cohomological ∂ -functor by items 1 and 2, and uniqueness comes from knowing it in dimension 0 (item 3). Ditto for Tor_R .

6 Group cohomology

To apply homology to groups, we will turn a group G into a ring, and consider modules over that ring.

Definition 12.6.1: Let R be a ring. The **group ring** $R[G]$ or RG is the ring

$$R^{\oplus G} = \left\{ \sum_{g \in G} a_g g : a_g \in R \right\}$$

with multiplication given by

$$\left(\sum_{g \in G} a_g g \right) \left(\sum_{h \in G} b_h h \right) = \sum_{g, h \in G} a_g b_h gh.$$

We will always work with $R = \mathbb{Z}$.

Note that any action of G on a \mathbb{Z} -module makes the module into a $\mathbb{Z}G$ -module. We often just abbreviate “ $\mathbb{Z}G$ -module” as “ G -module.”

Definition 12.6.2: Let G be a group and A, B be left $\mathbb{Z}G$ -modules.

1. The **diagonal action** of G on $\text{Hom}_{\mathbb{Z}}(A, B)$ is given by

$$(g\varphi)(a) = g(\varphi(g^{-1}a)).$$

2. The **diagonal action** of G on $A \otimes_{\mathbb{Z}G} B$ is given by

$$g(a \otimes b) = (ga) \otimes (gb).$$

We now apply cohomology as follows.

Definition 12.6.3: Let M be a G -module. Equip \mathbb{Z} with the trivial G -module structure. The **cohomology groups** of G with coefficients in M are defined by

$$\begin{aligned} H^n(G, M) &= \text{Ext}_{\mathbb{Z}G}^n(\mathbb{Z}, M) = H^n(\text{Hom}_{\mathbb{Z}G}(\mathbb{Z}, E^M)) \\ &= \text{ext}_{\mathbb{Z}G}^n(\mathbb{Z}, M) = H^n(\text{Hom}_{\mathbb{Z}G}(P_{\mathbb{Z}}, M)). \end{aligned}$$

Note from Theorem 12.4.3, we have two choices in finding $H^n(G, M)$: find a $\mathbb{Z}G$ -injective resolution of M , or a $\mathbb{Z}G$ -projective resolution of \mathbb{Z} .

There is a nice interpretation of $H^0(G, M)$.

Definition 12.6.4: Let L, M be G -modules and φ be a map $L \rightarrow M$. Define the **fixed point functor** by the following.

1. Action on modules:

$$M^G = \{m \in M : gm = m \text{ for all } g \in G\}.$$

2. Action on maps: Since $\varphi(L^G) \subseteq M^G$ we can define

$$\varphi^G = \varphi|_{L^G}.$$

Proposition 12.6.5: As functors,

$$H^0(G, \bullet) = \text{Hom}_{\mathbb{Z}G}(\mathbb{Z}, \bullet) = \bullet^G.$$

In particular, the fixed point functor is left exact since $\text{Hom}_{\mathbb{Z}G}(\mathbb{Z}, \bullet)$ is.

Proof. \mathbb{Z} is equipped with the trivial G -action. A G -homomorphism φ from \mathbb{Z} to M is determined by $\varphi(1)$, and $\varphi(1)$ must be a fixed point. Hence $\text{Hom}_{\mathbb{Z}G}(\mathbb{Z}, M) = M^G$ via the map $\varphi \mapsto \varphi(1)$. \square

Remark 12.6.6: This gives us another way to think about group cohomology. Given M , take an injective resolution $0 \rightarrow M \rightarrow E^0 \rightarrow E^1 \rightarrow \dots$. Applying $\text{Hom}_{\mathbb{Z}G}(\mathbb{Z}, \bullet)$ to this resolution is the same as applying \bullet^G , so we get $0 \rightarrow (E^0)^G \rightarrow (E^1)^G \rightarrow \dots$. Then $H^n(G, M)$ is the n th cohomology group of this complex.

We will need the fact that cohomology preserves products.

Proposition 12.6.7: Let G be a group and M_i be G -modules. Then

$$H^n \left(G, \prod_{i \in I} M_i \right) \cong \prod_{i \in I} H^n(G, M_i).$$

Proof. First note that the product of injective modules is an injective module: By definition a R -module I is injective iff $\text{Hom}_R(\bullet, I)$ is exact. Thus, the statement follows from the fact that $\text{Hom}_R(\bullet, \prod_i I_i) = \prod_i \text{Hom}_R(\bullet, I_i)$, and the fact that a product of exact sequences is exact.

Thus if E^{M_i} is an injective resolution for M_i , then $\prod_i E^{M_i}$ is an injective resolution for $\prod_i M_i$, and we get

$$H^n \left(G, \prod_{i \in I} M_i \right) = H^n \left(\text{Hom}_{\mathbb{Z}G} \left(\mathbb{Z}, E \prod_{i \in I} M_i \right) \right) = H^n \left(\text{Hom}_{\mathbb{Z}G} \left(\mathbb{Z}, \prod_{i \in I} E^{M_i} \right) \right) = \prod_{i \in I} H^n(G, M_i).$$

\square

7 Bar resolutions

We now describe the cohomology groups, by working with an explicit presentation of \mathbb{Z} . (We use the ext approach.) This will give practical interpretations of $H^1(G, M)$ and $H^2(G, M)$. For proofs, see Rotman [Rot09], Section 9.3.

Definition 12.7.1: Define the **bar resolution** $B(G)$ to be the exact sequence

$$\cdots \xrightarrow{d_3} B_2 \xrightarrow{d_2} B_1 \xrightarrow{d_1} B_0 \xrightarrow{d_0=\varepsilon} \mathbb{Z} \longrightarrow 0$$

where

$$B_n \cong \mathbb{Z}G^{\oplus G^n}$$

is the free abelian group with basis elements denoted by $[x_1 | \cdots | x_n]$, and

$$d_n([x_1 | \cdots | x_n]) = x_1[x_2 | \cdots | x_n] + \sum_{i=1}^{n-1} (-1)^i [x_1 | \cdots | \underbrace{x_i x_{i+1}}_i | \cdots | x_n] + (-1)^n [x_1 | \cdots | x_{n-1}]. \quad (12.5)$$

Let $U_n \subseteq B_n$ be the submodule generated by $[x_1 | \cdots | x_n]$ where at least one of the x_i equals 1, and define the **normalized bar resolution** to be the quotient complex $B^*(G) := B(G)/U(G)$.

Note in particular

$$\begin{aligned} d_3[x|y|z] &= x[y|z] - [xy|z] + [x|yz] - [x|y] \\ d_2[x|y] &= x[y] - [xy] + [x] \\ d_1[x] &= x[] - [] \\ d_0[] &= 1. \end{aligned}$$

We have $\text{Hom}_G(B_n, M) = \text{Hom}_G(\mathbb{Z}G^{\oplus G^n}, M)$, so it can be identified with the set of functions $G^n \rightarrow M$. Working out the kernels and images, we get the following.

Theorem 12.7.2. *We have the following descriptions of $H^1(G, M)$ and $H^2(G, M)$.*

1. Define a **derivation** (or crossed homomorphism) of G to be a function $G \rightarrow M$ such that

$$d(xy) = d(x) + xd(y)$$

and a **principal derivation** to be one in the form

$$d(x) = a - xa, \text{ for some } a \in M.$$

Denote the set of derivations and principal derivations by $\text{Der}(G, M)$ and $\text{PDer}(G, M)$. Then

$$H^1(G, M) \cong \text{Der}(G, M) / \text{PDer}(G, M).$$

2. We have

$$H^2(G, M) \cong \frac{\{f : G \times G \rightarrow M : f(x, y) + f(xy, z) = xf(y, z) + f(x, yz), f(x, 1) = f(1, y) = 0\}}{\{g : G \times G \rightarrow M : g(x, y) = xh(y) - h(xy) + h(x) \text{ for some } h : G \rightarrow M\}}.$$

The elements in the top set are called **factor sets**.

A particularly important case is the following.

Corollary 12.7.3. *Suppose G acts trivially on M . Then*

$$H^1(G, M) \cong \text{Hom}_{\mathbb{Z}}(G, M).$$

(On the RHS, G and M are thought of as groups.)

Proof. Because the action is trivial, a derivation is just a function with $d(xy) = d(x) + d(y)$, i.e. a homomorphism. Moreover, any principal derivation is trivial. \square

8 Group homology

Definition 12.8.1: Let A be a G -module. Equip \mathbb{Z} with the trivial G -module structure. The **homology groups** of G with coefficients in \mathbb{Z} are defined by

$$\begin{aligned} H_n(G, A) &= \text{Tor}_n^{\mathbb{Z}G}(\mathbb{Z}, A) = H_n(P_{\mathbb{Z}} \otimes_{\mathbb{Z}G} A) \\ &= \text{tor}_n^{\mathbb{Z}G}(\mathbb{Z}, A) = H_n(\mathbb{Z} \otimes_{\mathbb{Z}G} P_A). \end{aligned}$$

There is similarly a nice interpretation of $H_0(G, M)$, as well as of $H_1(G, \mathbb{Z})$. Given a group G , define the map $\varepsilon : \mathbb{Z}G \rightarrow \mathbb{Z}$ by $\varepsilon(\sum_{g \in G} a_g g) = \sum_{g \in G} a_g$, and define

$$I_G := \ker(\varepsilon) = \left\{ \sum_{g \in G} a_g g : \sum_{g \in G} a_g = 0 \right\}.$$

Proposition 12.8.2: As functors,

$$H_0(G, \bullet) = \bullet / I_G \bullet;$$

i.e. there is a natural isomorphism

$$\begin{aligned} H_0(G, A) &= \mathbb{Z} \otimes_G A \rightarrow A / I_G A \\ m \otimes a &\mapsto ma + I_G A. \end{aligned}$$

Proof. The short exact sequence $0 \rightarrow I_G \rightarrow \mathbb{Z}G \xrightarrow{\varepsilon} \mathbb{Z}$ gives exactness of

$$I_G \otimes_G A \rightarrow \mathbb{Z}G \otimes_G A \rightarrow \mathbb{Z} \otimes_G A \rightarrow 0$$

since tensoring is right exact. (G acts trivially on the \mathbb{Z} on the right.) Thus,

$$H_0(G, A) = \mathbb{Z} \otimes_G A = (\mathbb{Z}G \otimes_G A) / (I_G \otimes_G A) = A / I_G A.$$

\square

Proposition 12.8.3: There are canonical homomorphisms $H_1(G, \mathbb{Z}) \cong I_G/I_G^2 \cong G^{\text{ab}}$.

Here G^{ab} denotes the *abelianization* of G , i.e. G/G' , where G' is the derived subgroup, the (normal) subgroup generated by the commutators $aba^{-1}b^{-1}$.

Proof. The long exact sequence in homology for $0 \rightarrow I_G \rightarrow \mathbb{Z}G \xrightarrow{\varepsilon} \mathbb{Z} \rightarrow 0$ is

$$\begin{array}{ccccccccccc} H_1(G, \mathbb{Z}G) & \longrightarrow & H_1(G, \mathbb{Z}) & \xrightarrow{\partial_1} & H_0(G, I_G) & \longrightarrow & H_0(G, \mathbb{Z}G) & \longrightarrow & H_0(G, \mathbb{Z}) & \longrightarrow & 0 \\ \parallel & & & & \parallel & & \parallel & & \parallel & & \\ 0 & & & & I_G/I_G^2 & & \mathbb{Z} & & \mathbb{Z} & & \end{array}$$

The left term is 0 by Example 12.4.8 since $\mathbb{Z}G$ is free, hence projective. Thus ∂_1 is injective. From Proposition 12.8.2, we get the middle two inequalities (since $H_0(G, \mathbb{Z}G) = \mathbb{Z}G/I_G\mathbb{Z}G = \mathbb{Z}$). Surjectivity of the map $\mathbb{Z} \rightarrow \mathbb{Z}$ gives that it is actually an isomorphism, so exactness gives ∂_1 is an isomorphism. It remains to show

$$I_G/I_G^2 \cong G/G'. \quad (12.6)$$

Define a map $f : G \rightarrow I_G/I_G^2$ by letting $f(x) = (x - 1) \bmod I_G^2$. This is a homomorphism because

$$\begin{aligned} f(xy) &= xy - 1 \bmod I_G^2 \\ &= (x - 1) + (y - 1) \bmod I_G^2 && (x - 1)(y - 1) \in I_G^2 \\ &= f(x)f(y). \end{aligned}$$

Now $G' \in \ker f$ since I_G/I_G^2 is abelian ($\mathbb{Z}G$, as an additive group, is abelian), so we get a map $f : G/G' \rightarrow I_G/I_G^2$.

Now define $g : I_G \rightarrow G/G'$ by $g(x - 1) = xG'$. (Note $x - 1, x \in G \setminus \{1\}$, is a free basis for G .) We have

$$\begin{aligned} g\left(\sum_{x \in G \setminus \{1\}} m_x(x - 1) \sum_{y \in G \setminus \{1\}} m_y(y - 1)\right) &= g\left(\sum_{x, y \in G \setminus \{1\}} m_x m_y ((xy - 1) - (x - 1) - (y - 1))\right) \\ &= \prod_{x, y \in G \setminus \{1\}} (xyx^{-1}y^{-1})^{m_x m_y} G' = G' \end{aligned}$$

so g induces $g : I_G/I_G^2 \rightarrow G/G'$.

Now f and g are inverse, showing (12.6). \square

8.1 Shapiro's lemma

Shapiro's lemma will be helpful in computing (co)homology groups, especially in the guise of Corollary 12.8.8.

Definition 12.8.4: Let $S \subseteq G$ be a subgroup of finite index. Define the **induced** and **coinduced modules** to be³

$$\begin{aligned}\text{Ind}_S^G(A) &= A \otimes_{\mathbb{Z}S} \mathbb{Z}G. \\ \text{Coind}_S^G(A) &= \text{Hom}_{\mathbb{Z}S}(\mathbb{Z}G, A).\end{aligned}$$

If $S = \{1\}$ we simply write $\text{Ind}^G(A)$ or $\text{Coind}^G(A)$. An **induced module** of G is a module in the form $\text{Ind}^G(A)$; a **coinduced module** of G is a module in the form $\text{Coind}^G(A)$.

Remark 12.8.5: If G is finite, the induced and coinduced modules are canonically isomorphic via the below map, so there is no need to distinguish between them.

$$\begin{aligned}\text{Hom}_S(\mathbb{Z}G, A) &\rightarrow A \otimes_{\mathbb{Z}S} \mathbb{Z}G \\ \varphi &\mapsto \sum_{g \in G/S} \varphi(g^{-1}) \otimes_{\mathbb{Z}S} g.\end{aligned}$$

Proposition 12.8.6: If M is a coinduced G -module, and $H \subseteq G$ is a subgroup, then M is a coinduced H -module.

Proof. Write $M = \text{Hom}_{\mathbb{Z}}(\mathbb{Z}[G], A)$; we can write $\mathbb{Z}[G] = \mathbb{Z}[H] \otimes B$; then we have by adjoint associativity⁴ that $M = \text{Hom}(\mathbb{Z}[H] \otimes M, A) = \text{Hom}(\mathbb{Z}[H], \text{Hom}(M, A))$. \square

The cohomology of coinduced modules and the homology of induced modules are easy to calculate.

Lemma 12.8.7 (Shapiro's lemma). *The following hold.*

$$\begin{aligned}H^n(G, \text{Coind}_S^G(A)) &= H^n(S, A) \\ H_n(G, \text{Ind}_S^G(A)) &= H_n(S, A).\end{aligned}$$

Proof. Let $P_{\mathbb{Z}}$ be a $\mathbb{Z}G$ -projective resolution of \mathbb{Z} . Note it is also a $\mathbb{Z}S$ -projective resolution, as any $\mathbb{Z}G$ -projective module is $\mathbb{Z}S$ -projective.

By definition of cohomology group,

$$\begin{aligned}H^n(G, \text{Coind}_S^G(A)) &= H^n(\text{Hom}_{\mathbb{Z}G}(P_{\mathbb{Z}}, \text{Hom}_{\mathbb{Z}S}(\mathbb{Z}G, A))) \\ &\stackrel{(*)}{=} H^n(\text{Hom}_{\mathbb{Z}S}(P_{\mathbb{Z}} \otimes_{\mathbb{Z}G} \mathbb{Z}G, A)) = H^n(\text{Hom}_{\mathbb{Z}S}(P_{\mathbb{Z}}, A)) = H^n(S, A).\end{aligned}$$

In (*) we used adjoint associativity.

By the definition of homology group,

$$H_n(G, \text{Ind}_S^G(A)) = H_n(P_{\mathbb{Z}} \otimes_{\mathbb{Z}G} (\mathbb{Z}G \otimes_{\mathbb{Z}S} A)) = H_n(P_{\mathbb{Z}} \otimes_{\mathbb{Z}S} A) = H_n(S, A). \quad \square$$

³Be careful; in some books the definitions are reversed. We follow Serre's definition, which is the opposite of Milne's definitions.

⁴If R, R' are rings, M is a R -module, N is a (R, R') -bimodule, and P is a R' -module, then there is a canonical (R, R') -isomorphism $\text{Hom}_R(M, \text{Hom}_{R'}(N, P)) \cong \text{Hom}_{R'}(M \otimes_R N, P)$.

Corollary 12.8.8. *Suppose that $A = \bigoplus_{i \in I} A_i$, $S = \text{Stab}(A_j)$ (defined as $\{g \in G : gA_j = A_j\}$), and G permutes the submodules A_i transitively. Then*

$$H_n(G, A) = H_n(S, A_j).$$

If G is finite, then

$$H^n(G, A) = H^n(S, A_j).$$

Proof. We have $A = \text{Ind}_S^G A_j$. If G is finite then $A \cong \text{Coind}_S^G A_j$ as well. □

Corollary 12.8.9. *If M is an coinduced G -module, then $H^n(G, M) = 0$ for all $n \geq 1$.*

If M is an induced G -module, then $H_n(G, M) = 0$ for all $n \geq 1$.

Proof. By Shapiro's lemma 12.8.7,

$$\begin{aligned} M = \text{Coind}_S^G(A) &\implies H^n(G, M) = H^n(1, M) = 0 \\ M = \text{Ind}_S^G(A) &\implies H_n(G, M) = H_n(1, M) = 0. \end{aligned}$$

We used the fact that \mathbb{Z} is $\mathbb{Z}[\{1\}]$ -projective. □

9 Tate groups

By Corollary 12.4.7, given a short exact sequence of G -modules we get a long exact sequence in homology and cohomology. We splice these sequences together using the Snake Lemma to obtain a long exact sequence extending in both directions.

Definition 12.9.1: Let G be a group, S be a subgroup of finite index, and A be a G -module. Define the **norm** $N_{G/S} : A^S \rightarrow A^G$ by

$$N_{G/S}(a) = \sum_{j=1}^n t_j a,$$

where $\{t_1, \dots, t_n\}$ is a left transversal (i.e. coset representatives) of S in G . In particular, for $S = \{1\}$ the norm map is

$$N_G(a) = N(a) = \left(\sum_{g \in G} g \right) a.$$

Definition 12.9.2: Suppose G is a finite group and A is a G -module. Define the **Tate groups** by


$$H_T^q(G, A) = \begin{cases} H^q(G, A), & q \geq 1 \\ A^G / NA, & q = 0 \\ {}_N A / I_G A, & q = -1 \\ H_{-q-1}(G, A), & q \leq -2. \end{cases}$$

Here ${}_N A$ denotes $\{a \in A : Na = 0\}$.


Theorem 12.9.3. *If G is a finite group and $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is an exact sequence of G -modules, then there is a long exact sequence*

$$\cdots \rightarrow H_T^q(G, A) \rightarrow H_T^q(G, B) \rightarrow H_T^q(G, C) \rightarrow H_T^{q-1}(G, A) \rightarrow \cdots$$

Proof. It suffices to prove exactness for $q = -1$ and $q = 0$. We apply to the snake lemma to obtain the following (the top and bottom rows in the middle are the long exact sequence in homology and cohomology, respectively).



The maps N_A, N_B, N_C are the norm maps on $A, B,$ and C after associating H_0 and H^0 with their descriptions in Propositions 12.6.5 and 12.8.2:



□

9.1 Complete resolution*

⁵ The description of Tate groups in the last section is somewhat unwieldy (because you can see the glue marks...). We give a different interpretation here, where the Tate groups at 0 and -1 are less distinguished. Then we use the technique of “dimension shifting” to extend results for cohomology (or homology) groups to results for Tate groups.

Definition 12.9.4: A **complete resolution** of a group G is an exact sequence \mathbf{X}

$$\begin{array}{ccccccc} \cdots & \longrightarrow & X_1 & \longrightarrow & X_0 & \xrightarrow{d_0} & X_{-1} & \longrightarrow & X_{-2} & \longrightarrow & \cdots \\ & & & & \searrow \varepsilon & & \nearrow \eta & & & & \\ & & & & & & \mathbb{Z} & & & & \end{array}$$

where each X_q is a finitely generated G -free module, ε is surjective, and η is injective.

Proposition 12.9.5: Every finite group G has a complete resolution \mathbf{X} .

Proof. Take a G -free resolution of \mathbb{Z} and its dual ($A^* = \text{Hom}_{\mathbb{Z}}(A, \mathbb{Z})$), and splice them together.

$$\begin{array}{ccccccc} \cdots & \longrightarrow & P_1 & \longrightarrow & P_0 & \twoheadrightarrow & \mathbb{Z} & \longrightarrow & 0 \\ & & & & \searrow & & \parallel & & \\ & & & & & & \mathbb{Z} & \longrightarrow & P_0^* & \longrightarrow & P_1^* & \longrightarrow & \cdots \\ & & & & 0 & \longrightarrow & & & & & & & \end{array}$$

□

Proposition 12.9.6: Let G be a finite group, A a G -module, and \mathbf{X} a complete resolution. Then the Tate groups are exactly the cohomology groups

$$H_T^n(G, A) = H^n(\text{Hom}_G(\mathbf{X}, A)).$$

Proof. Since any two resolutions are chain-homotopic (going both ways) by the Comparison Theorem 12.2.4, it suffices to prove this for one resolution. We take a resolution as in

⁵This section will not be used and can be omitted.

Proposition 12.9.5 and apply $\text{Hom}_G(\bullet, A)$ to it. We obtain the following.

$$\begin{array}{ccccccc}
 & & -2 & & -1 & & 0 & & 1 & & \\
 \cdots & \longrightarrow & \text{Hom}_G(P_1^*, A) & \longrightarrow & \text{Hom}_G(P_0^*, A) & \longrightarrow & \text{Hom}_G(P_0, A) & \longrightarrow & \text{Hom}_G(P_1, A) & \longrightarrow & \cdots \\
 & & \downarrow \cong & & \downarrow \cong & & \parallel & & \parallel & & \\
 & \longrightarrow & P_1 \otimes_{\mathbb{Z}G} A & \xrightarrow{d^{-2}} & P_0 \otimes_{\mathbb{Z}G} A & \xrightarrow{d^{-1}} & \text{Hom}_G(P_0, A) & \xrightarrow{d^0} & \text{Hom}_G(P_1, A) & \longrightarrow & \\
 & & \searrow & & \downarrow \varepsilon \otimes \bullet & & \uparrow \varepsilon^* & & \nearrow & & \\
 & & & & \mathbb{Z} \otimes_G A & \xrightarrow{N_A} & \text{Hom}_G(\mathbb{Z}, A) & & & & \\
 & & & & \parallel & & \parallel & & & & \\
 & & & & A/I_G A & & A^G & & & &
 \end{array}$$

The isomorphisms on the left are given by the natural isomorphism

$$\begin{aligned}
 M \otimes_{\mathbb{Z}G} A &\rightarrow \text{Hom}_G(M^*, A) \\
 m \otimes a &\mapsto (f \mapsto f(m)a).
 \end{aligned}$$

The bent complex along the bottom is the complex for Tate cohomology; some diagram chasing gives that these groups are isomorphic to the cohomology groups in the middle complex. \square

9.2 Dimension shifting

Given a result or construction in dimension n , we can get the result in dimensions $n \pm 1$ by utilizing the long exact sequence 12.9.3 and the two propositions.

Proposition 12.9.7: Let G be a finite group. If M is an induced module then

$$H_T^n(G, M) = 0$$

for all n .

Proof. Since G is finite, induced and coinduced modules are the same. The statement for homology and cohomology is Corollary 12.8.9; this takes care of all $n \neq 0, -1$. For $n = 0, -1$ we calculate $H_T^n(G, M)$ directly. Writing $M = A \otimes_{\mathbb{Z}} \mathbb{Z}G$, we see that every element of m can be uniquely written as $\sum_{g \in G} a_g \otimes g$. We find that

$$\begin{aligned}
 M^G &= \left\{ a \otimes \sum_{g \in G} g : a \in A \right\} = N(M) \\
 {}_N M &= \left\{ \sum_{g \in G} a_g \otimes g : \sum_{g \in G} a_g = 0 \right\} = I_G M
 \end{aligned}$$

so $H_T^0(G, M) = H_T^{-1}(G, M) = 0$. \square

Proposition 12.9.8: Let M be a module. Then there exist (canonical) short exact sequences

$$\begin{aligned} 0 \rightarrow M \rightarrow M^* \rightarrow M^*/M \rightarrow 0 \\ 0 \rightarrow M' \rightarrow M_* \rightarrow M \rightarrow 0 \end{aligned}$$

such that M^* is coinduced and M_* is induced, and these sequences are split as abelian groups (i.e. as \mathbb{Z} -modules, but not necessarily as $\mathbb{Z}G$ -modules).

Proof. The desired maps are

$$\begin{aligned} M &\hookrightarrow \text{Coind}_{\{1\}}^G(M) \\ m &\mapsto \varphi_m(g) = gm. \\ \mathbb{Z}[G] \otimes_{\mathbb{Z}} M &\twoheadrightarrow M \\ g \otimes m &\mapsto gm \end{aligned}$$

Splitness follows from the fact that these maps have left and right inverses, respectively: $\varphi \mapsto \varphi(1)$ and $m \mapsto 1 \otimes m$. (They are only \mathbb{Z} -homomorphisms, not necessarily $\mathbb{Z}G$ -homomorphisms.) \square

Now suppose G is finite; then coinduced and induced modules coincide. Taking the long exact sequence 12.9.3 of the above short exact sequences and using Proposition 12.9.7 gives

$$\begin{aligned} H_T^n(G, M) &\cong H_T^{n-1}(G, M^*/M) \\ H_T^n(G, M) &\cong H_T^{n+1}(G, M'). \end{aligned}$$

Thus we reduce a problem about cohomology in degree n to a problem about cohomology in degree $n + 1$ or degree $n - 1$.

10 Cup products

There is a natural product defined in Tate cohomology.

Define $A \otimes B$ to be $A \otimes_{\mathbb{Z}} B$ with the structure of a G -module given by $g(a \otimes b) = ga \otimes gb$ (the diagonal action).

Theorem 12.10.1. *Let G be a finite group and A, B be G -modules. There exists a unique family of bilinear maps indexed by $(p, q) \in \mathbb{Z}^2$, together called the **cup product**,*

$$\cup : H_T^p(G, A) \times H_T^q(G, B) \rightarrow H_T^{p+q}(G, A \otimes B),$$

satisfying the following four properties.

1. *The homomorphisms are functorial in A and B .*

2. For $p = q = 0$, the cup product is induced by the map

$$A^G \otimes B^G \rightarrow (A \otimes B)^G.$$

3. If

$$\begin{aligned} 0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0 \\ 0 \rightarrow A' \otimes B \rightarrow A \otimes B \rightarrow A'' \otimes B \rightarrow 0 \end{aligned}$$

are exact⁶, and $a'' \in H_T^p(G, A'')$, $b \in H_T^q(G, B)$, then

$$(\delta a'') \cup b = \delta(a'' \cup b)$$

in $H_T^{p+q+1}(G, A' \otimes B)$. (δ is the map in the corresponding long exact sequence.)

4. If

$$\begin{aligned} 0 \rightarrow B' \rightarrow B \rightarrow B'' \rightarrow 0 \\ 0 \rightarrow A \otimes B' \rightarrow A \otimes B \rightarrow A \otimes B'' \rightarrow 0 \end{aligned}$$

are exact, and $a \in H_T^p(G, A)$, $b'' \in H_T^q(G, B'')$, then

$$a \cup (\delta b'') = (-1)^p \delta(a \cup b'')$$

in $H_T^{p+q+1}(G, A \otimes B')$.

Proof. We first define the cup product for cohomology groups and then use dimension shifting to define it for Tate groups.

We use the bar resolution⁷, so that n -chains are functions $G^n \rightarrow A$. For $p, q \geq 0$, define

$$\cup : C^p(G, A) \times C^q(G, B) \rightarrow C^{p+q}(G, A \otimes B)$$

by

$$(f \cup g)[x_1 | \cdots | x_{p+q}] = f([x_1 | \cdots | x_p]) \otimes g([x_{p+1} | \cdots | x_{p+q}]).$$

For $n = 0$, we have $(f \cup g)[\] = f[\] \otimes g[\]$ which shows property 2 is satisfied. We⁸ can laboriously verify with (12.5) that

$$d(f \cup g) = (df) \cup g + (-1)^p f \cup (dg).$$

From this we get a well-defined map

$$\cup : H^p(G, A) \times H^q(G, B) \rightarrow H^{p+q}(G, A \otimes B).$$

⁶Recall $\bullet \otimes B$ is right exact, so the content is in left exactness.

⁷We can also use the standard resolution (not defined here); in that case the map is $(f \cup g)(x_0, \dots, x_{p+q}) = f(x_0, \dots, x_p) \otimes g(x_{p+1}, \dots, x_{p+q})$.

⁸i.e. you

We can verify properties 3 and 4 by calculation.

Now we extend this definition by dimension shifting. Suppose the product is defined for $(p+1, q)$, we define it for (p, q) as follows. Write A (canonically) as a quotient of an induced module as in Proposition 12.9.8, $0 \rightarrow A' \rightarrow A_* \rightarrow A \rightarrow 0$. Since this is split, so is

$$0 \rightarrow A' \otimes B \rightarrow A_* \otimes B \rightarrow A \otimes B \rightarrow 0.$$

Since A_* is induced, so is $A_* \otimes B$ (be slightly careful about the G -action here). Thus by Theorem 12.9.3, we get $H_T^p(A) \cong H_T^{p+1}(A')$ and $H_T^{p+q}(A) \cong H_T^{p+q+1}(A' \otimes B)$ (naturally), and thus we can define the cup product

$$\begin{array}{ccc} H_T^p(A) \times H_T^q(B) & \xrightarrow{\cup} & H_T^{p+q}(A \otimes B) \\ \downarrow \cong & & \uparrow \cong \\ H_T^{p+1}(A') \times H_T^q(B) & \xrightarrow{\cup} & H_T^{p+q+1}(A' \otimes B) \end{array}$$

Similarly define it for (p, q) given $(p, q+1)$, but this time introduce a factor of $(-1)^p$ (in order to make the second condition hold). Note this is consistent with our definitions for $p, q \geq 0$, by conditions 3 and 4. It is not hard to verify that these maps are well-defined, and that conditions 3 and 4 continue to be satisfied. By the way we defined the maps, it also doesn't matter what order we define the maps in (so going from $(p+1, q+1) \rightarrow (p, q+1) \rightarrow (p, q)$ is the same as going from $(p+1, q+1) \rightarrow (p+1, q) \rightarrow (p, q)$, for instance).

Given the map for (p, q) , conditions 3 and 4 basically force us to define the map for $(p-1, q)$ and $(p, q-1)$ as above. Similarly we can dimension-shift in the opposite direction, and we get uniqueness for all (p, q) . \square

Cup products are rather nasty to work with when they aren't purely in cohomology, so if we need to do cup product computation, we work in cohomology whenever possible.

Proposition 12.10.2: The following hold:

1. Cup product is associative: For $x \in H^m(G, M)$, $y \in H^n(G, N)$, and $z \in H^p(G, P)$, $(x \cup y) \cup z = x \cup (y \cup z)$ (viewing the equation in $H^{m+n+p}(G, M \otimes N \otimes P)$).
2. Cup product is anticommutative: For $x \in H^m(G, M)$ and $y \in H^n(G, N)$, $x \cup y = (-1)^{mn} y \cup x$.

Proof. Omitted. The idea is to verify the formula in degree 0 and then dimension-shift to get the general case. \square

10.1 Cup product calculations

To compute the Artin map in class field theory, we will need to calculate the cup product of things in dimensions -2 and 2 . We will get there incrementally using dimension shifting and properties 3–4 of the cup product, first calculating the cup product on dimensions $(0, n)$ (especially $(0, 1)$), then on $(-1, 1)$, and then finally on $(-2, 1)$.

Theorem 12.10.3. *Let G be a finite group and A, B G -modules. If $a \in A^G$, let \bar{a}^0 denote its image in $H_T^0(G, A)$, and if $Na = 0$, let \bar{a}_0 denote its image in $H_T^{-1}(G, A)$. For $g \in G$ let \bar{g} denote its image in $G/G' = H_T^{-2}(G, \mathbb{Z})$.*

1. $(0, n)$. Suppose $n \geq 0$, $a \in A^G$, and $x \in H_T^n(G, B)$. Let $f_a : B \rightarrow A \otimes B$ be the map sending y to $a \otimes y$; it induces a map $H_T^n(G, A) \rightarrow H_T^n(G, A \otimes B)$. Then

$$\underbrace{\bar{a}^0}_{\in H_T^0(G, A)} \cup \underbrace{x}_{\in H_T^n(G, B)} = f_a(x) \in H_T^n(G, A \otimes B).$$

2. $(-1, 1)$. Suppose $Na = 0$, and $[f] \in H^1(G, B)$ is represented by a cocycle $f : G \rightarrow B$. Then

$$\underbrace{\bar{a}_0}_{\in H_T^{-1}(G, B)} \cup \underbrace{[f]}_{\in H_T^1(G, B)} = \overline{\left(-\sum_{t \in G} ta \otimes f(t)\right)}^0.$$

3. $(-2, 1)$. Let $s \in G$ and $[f] \in H^1(G, B)$. Then

$$\underbrace{\bar{s}}_{\in H_T^{-2}(G, \mathbb{Z})} \cup \underbrace{\bar{f}}_{\in H_T^1(G, B)} = \overline{f(s)}_0 \in H_T^{-1}(G, B).$$

Proof. We omit details of the calculations. See Serre [Ser79], pg. 176-178.

1. For $n = 0$, this follows from definition of cup product. Now use dimension shifting, with the exact sequence $0 \rightarrow B \rightarrow B^* \rightarrow B^*/B \rightarrow 0$, B^* coinduced.
2. Dimension shift from part 1 with $0 \rightarrow B \rightarrow B^* \rightarrow B^*/B \rightarrow 0$: suppose $b'' \in (B^*/B)^G$ is sent to f under the diagonal morphism. Write $\bar{a}_0 \cup \bar{f} = \bar{a}_0 \cup d(\bar{b}'') = -d(\bar{a}_0 \cup \bar{b}'')$ and use part 1.
3. Show that

$$d(\bar{s} \cup [f]) = d(\overline{f(s)}_0).$$

Evaluate the LHS using property 3 and part 2.

□

11 Change of group

We would like to be able to connect (co)homology groups corresponding to different groups G, G' and different modules over G, G' . This will allow us, for example, to define maps

$$\begin{aligned} \text{Res}^n : H^n(G, A) &\rightarrow H^n(S, A) \\ \text{Cor}_n : H_n(S, A) &\rightarrow H_n(G, A) \\ \text{Inf}^n : H^n(G/S, A^S) &\rightarrow H^n(G, A) \end{aligned} \quad S \trianglelefteq G.$$

11.1 Construction of maps

For there to be a map $H^n(G, A) \rightarrow H^n(G', A')$ we need there to be a map $G' \rightarrow G$, with some compatibility condition on the modules A, A' .

Definition 12.11.1: Let G, G' be groups, let A be a G -module and A' be a G' -module. A **cocompatible pair** is a pair (α, f) where $\alpha : G' \rightarrow G$ is a group homomorphism and $f : A \rightarrow A'$ is a \mathbb{Z} -homomorphism such that

$$f((\alpha x')a) = x'f(a)$$

for all $x' \in G'$ and $a \in A$.

$$\begin{array}{ccc} G' & \xrightarrow{\alpha} & G \\ \left. \vphantom{G'} \right\} & & \left. \vphantom{G} \right\} \\ A' & \xleftarrow{f} & A \end{array}$$

Let $((\text{Pairs}^*))$ denote the category whose objects are pairs (G, A) and whose morphisms are cocompatible (α, f) .

Define a **compatible pair** to be a pair (α, f) where $\alpha : G \rightarrow G'$ is a group homomorphism and $g : A \rightarrow A'$ is a \mathbb{Z} -homomorphism such that

$$f(xa) = (\alpha x)f(a)$$

for all $x \in G$.

$$\begin{array}{ccc} G & \xrightarrow{\alpha} & G' \\ \left. \vphantom{G} \right\} & & \left. \vphantom{G'} \right\} \\ A & \xrightarrow{f} & A' \end{array}$$

Let $((\text{Pairs}))$ denote the category whose objects are ordered pairs (G, A) and whose morphisms are compatible (α, f) .

Given a cocompatible pair, let P' be a G' -projective resolution of \mathbb{Z} and P be a G -projective resolution of \mathbb{Z} . By the Comparison Theorem 12.2.4 there is a chain map $\tau(\alpha) : P' \rightarrow P$ induced by the map $1_{\mathbb{Z}} : \mathbb{Z} \rightarrow \mathbb{Z}$ and α , unique up to homotopy. Define

$$\begin{aligned} C^n(G, A) &= \text{Hom}_{\mathbb{Z}G}(P_n, A) \rightarrow \text{Hom}_{\mathbb{Z}G}(P'_n, A') = C^n(G', A') \\ \varphi &\mapsto f \circ \varphi \circ \tau(\alpha)^n. \end{aligned}$$

Similarly, for a compatible pair, there is a chain map $\tau(\alpha) : P \rightarrow P'$ induced by $1_{\mathbb{Z}} : \mathbb{Z} \rightarrow \mathbb{Z}$ and α ; we get a map

$$\tau(\alpha)_n \otimes f : C_n(G, A) = P_n \otimes_{\mathbb{Z}G} A \rightarrow P'_n \otimes_{\mathbb{Z}G'} A' = C_n(G', A')$$

These maps descend to cohomology and homology, respectively.

Definition 12.11.2: Define the maps below using the (co)compatible pairs shown.

Name	Map on G	Map on M	Map
Restriction	$i : S \rightarrow G$	$M \xleftarrow{\cong} M$	$\text{Res}_{G/S}^n : H^n(G, M) \rightarrow H^n(S, M)$
Corestriction	$i : S \rightarrow G$	$M \xrightarrow{\cong} M$	$\text{Cor}_{S/G}^n : H_n(S, M) \rightarrow H_n(G, M)$
Inflation	$q : G \rightarrow G/S$	$M \leftrightarrow M^S$	$\text{Inf}_{S/G}^n : H^n(G/S, M^S) \rightarrow H^n(G, M)$
Conjugation	$\sigma \mapsto g\sigma g^{-1}$	$g^{-1}m \leftrightarrow m$	$H^n(G, M) \rightarrow H^n(G, M)$

For inflation, we require that $S \trianglelefteq G$ (S be a normal subgroup of G).

Proposition 12.11.3: The conjugation map $H^n(G, M) \rightarrow H^n(G, M)$ is the identity.

This is important because when we are defining maps between different cohomology groups, we can be assured that conjugation won't change it, i.e. we have a canonical map.

Proof. For $n = 0$ this is the identity map $M^G \rightarrow M^G$. Since the conjugation $H^n(G, M) \rightarrow H^n(G, M)$ is a map of cohomological functors, and the identity map $H^n(G, M) \rightarrow H^n(G, M)$ is also a map of cohomological functors, and they agree for $n = 0$, by Theorem 12.5.2(2) they must be equal for all n .

Alternatively, use dimension shifting. □

11.2 Extending maps to Tate cohomology

Right now Res^n is only defined on cohomology and Cor_n is only defined on homology. We would like to define them on Tate cohomology.

Proposition 12.11.4: Let G be a finite group. The maps Res^n and Cor_n can be defined on Tate cohomology, such that the definitions for H_T^n agree with the original definitions on cohomology and homology for $n \geq 0$ and $n \leq -1$, respectively, and such that Res and Cor are natural transformations compatible with forming the long exact sequence in homology and cohomology from a short exact sequence. Moreover, Res^n and Cor_n satisfy the following properties.

1. $\text{Cor}_{S/G}^0 : H_T^0(S, M) \rightarrow H_T^0(G, M)$ is the map $N_{G/S} : M^S/N_S M \rightarrow M^G/N_G M$.
2. $\text{Res}_{G/S}^{-1} : H_T^{-1}(G, M) \rightarrow H_T^{-1}(S, M)$ is the map $C_{G/S} : {}_N G M / I_G M \rightarrow {}_N S M / I_S M$, where $C_{G/S}$ is the **conorm** map defined by

$$C_{G/S}(a) := \sum_i t_i^{-1} a$$

where $\{t_i\}$ is a left transversal of G/S . (Equivalently, let $\{t_i\}$ be a *right* transversal and let $C_{G/S}(a) := \sum_i t_i a$.⁹)

⁹To see this, note $t_1 S = t_2 S$ iff $t_1^{-1} t_2 \in S$, iff $S t_1^{-1} = S t_2^{-1}$.

3. $\text{Cor}_{S/G}^{-2} : H_T^{-2}(S, M) \rightarrow H_T^{-2}(G, M)$ is the natural map $S^{\text{ab}} \rightarrow G^{\text{ab}}$. (See Proposition 12.8.3.)

Proof. First, the construction. We will use Theorem 12.5.2. Let χ be the class of coinduced $\mathbb{Z}G$ -modules. Note that the category of $\mathbb{Z}G$ -modules has enough coinduced $\mathbb{Z}G$ -modules, by Proposition 12.9.8. Note that $\{H_T^n(S, \bullet_S)\}$ and $\{H_T^n(G, \bullet)\}$ are cohomological ∂ -functors on the category of $\mathbb{Z}G$ -modules, with respect to χ (by M_S , we mean think of M as a S -module). Indeed, any coinduced module for G is coinduced for S by Proposition 12.8.6.¹⁰ Since

$$\text{Res}_{G/S}^0 : M^G/N_G M \rightarrow M^S/N_S M, \quad \text{Cor}_0^{S/G} : N_S M/I_S M \rightarrow N_G M/I_G M$$

are natural transformations, Theorem 12.5.2(1) applies to give unique morphisms Res and Cor extending $N_{G/S}$. (They agree in cohomology and homology with the original definitions by uniqueness in Theorem 12.5.2(1)).

Alternatively, we can extend the definitions of Res and Cor using dimension shifting (which is simpler, really).¹¹

We now calculate the maps using dimension shifting.

1. Use the short exact sequence $0 \rightarrow M' \rightarrow M^* = \mathbb{Z}G \otimes_{\mathbb{Z}} M \rightarrow M \rightarrow 0$ from Proposition 12.9.8 to get the vertical isomorphisms in the diagram on the left below. (Note as before that M^* is both G and S -(co)induced.)

$$\begin{array}{ccc} H_T^{-1}(S, M) & \xrightarrow{\text{Cor}_{S/G}^{-1}} & H_T^{-1}(G, M) & & N_S M/I_S M & \longrightarrow & N_G M/I_G M \\ \delta \downarrow \cong & & \delta \downarrow \cong & & N_S(1 \otimes \bullet) \downarrow \cong & & N_G(1 \otimes \bullet) \downarrow \cong \\ H_T^0(S, M') & \xrightarrow{\text{Cor}_{S/G}^0} & H_T^0(G, M) & & H_T^0(S, M') & \xrightarrow{?} & H_T^0(G, M). \end{array}$$

The left-hand diagram gives the right-hand diagram, after noting that δ is the map in the snake lemma in the proof of Theorem 12.9.3. From the right-hand diagram it is clear that the bottom map has to be $N_{G/S}$, because $N_{G/S} \circ N_S = N_G$.

2. From $0 \rightarrow M \rightarrow M^* \xrightarrow{f} M^*/M \rightarrow 0$ we get the commutative diagrams

$$\begin{array}{ccc} H_T^{-1}(G, M^*/M) & \xrightarrow{\text{Res}_{G/S}^{-1}} & H_T^{-1}(S, M^*/M) & & H_T^{-1}(G, M^*/M) & \xrightarrow{?} & H_T^{-1}(S, M^*/M) \\ \delta \downarrow \cong & & \delta \downarrow \cong & & N_G \circ f^{-1} \downarrow \cong & & N_S \circ f^{-1} \downarrow \cong \\ H_T^0(G, M) & \xrightarrow{\text{Res}_{G/S}^0} & H_T^0(S, M) & & M^G/N_G M & \longrightarrow & M^S/N_S M. \end{array}$$

¹⁰Note this would fail if we take χ to be the class of $\mathbb{Z}G$ -injective modules, as $\mathbb{Z}G$ -injective modules are not necessarily $\mathbb{Z}S$ -injective.

¹¹Alternatively, we can construct Cor^n explicitly as the map

$$H^n(S, M) \xrightarrow{\text{Shapiro}} H^n(G, \text{Coind}_S^G M) \rightarrow H^n(G, M)$$

where the last map is the change of group map induced by $G \cong G$ and $\text{Coind}_S^G M \rightarrow M$ given by $\phi \mapsto \sum_i t_i \varphi(t_i^{-1})$, for some transversal $\{t_i\}$ for S in G . This is just the norm map in dimension 0.

From $N_G = N_S \circ C_{G/S}$, the top map has to be $C_{G/S}$.

3. Recall the isomorphism $H_1(G^{\text{ab}}, \mathbb{Z}) \cong G^{\text{ab}}$ was defined using the horizontal maps below.

$$\begin{array}{ccccccc} H_1(S, \mathbb{Z}) & \xrightarrow[\cong]{\partial_1} & H_0(S, I_S) & \xlongequal{\quad} & I_S/I_S^2 & \longrightarrow & S/S' \\ \downarrow \text{Cor}_1 & & \downarrow \text{Cor}_0 & & & & \downarrow \\ H_1(G, \mathbb{Z}) & \xrightarrow[\cong]{\partial_1} & H_0(G, I_G) & \xlongequal{\quad} & I_G/I_G^2 & \longrightarrow & G/G' \end{array}$$

The left square commutes by functoriality of Cor and the right rectangle commutes by tracing the map in Proposition 12.8.3. \square

11.3 Further properties

Theorem 12.11.5. *Suppose H is a subgroup of G of finite index. Then $\text{Cor}^n \circ \text{Res}^n$ is multiplication by $[G : H]$.*

Proof. In degree 0, we have $\text{Cor}^0 \circ \text{Res}^0 = [G : H]$ because $N_{G/H}$ is just multiplication by $[G : H]$ on M^G . As in the proof of Proposition 12.11.3, the general case then follows from either Theorem 12.5.2 or dimension shifting. \square

Corollary 12.11.6.

1. *If G is finite, then $|G|H^n(G, M) = 1$ for any $n > 0$.*
2. *If G is finite and M is finitely generated as an abelian group, then $H^n(G, M)$ is finite.*

Proof.

1. By Theorem 12.11.5,

$$H^n(G, M) \xrightarrow{\text{Res}} H^n(1, M) \xrightarrow{\text{Cor}} H^n(G, M)$$

is multiplication by $|G|$. But $H^n(1, M) = 0$.

2. By the explicit description of $H^n(G, M)$ using the bar resolution, $H^n(G, M)$ is finitely generated. By item 1 it has finite exponent, so it must be finite. \square

Corollary 12.11.7. *Let G be a finite group and G_p its p -SSG. For any G -module M , the map*

$$\text{Res}^n : H^n(G, M) \rightarrow H^n(G_p, M)$$

is injective on the p -primary component.

Proof. Suppose that $x \in \ker(\text{Res})$. Then $[G : G_p]x = \text{Cor} \circ \text{Res}(x) = 0$. Since the order of x is a power of p but $p \nmid [G : G_p]$, we get that $x = 0$. \square

Corollary 12.11.8. *If $H_T^n(G_p, A) = 0$ for all primes p then $H_T^n(G, A) = 0$.*

We will also need to know how restriction and corestriction affect cup products.

Proposition 12.11.9: The following hold.

1. $\text{Res}(x \cup y) = \text{Res}(x) \cup \text{Res}(y)$.
2. $\text{Cor}(x \cup \text{Res}(y)) = \text{Cor}(x) \cup y$.

Proof. See Cartan-Eilenberg [CE56], Chapter 12, or Atiyah-Wall in Cassels-Frohlich [CF69], p. 107. □

11.4 Inflation-restriction exact sequence

Proposition 12.11.10: Suppose $H \trianglelefteq G$, A is a G -module, and $n > 0$. If $H^i(H, A) = 0$ for all i with $0 < i < r$, then

$$0 \rightarrow H^r(G/H, A^H) \xrightarrow{\text{Inf}} H^r(G, A) \xrightarrow{\text{Res}} H^r(H, A)$$

is exact.

Proof. We first prove the case $r = 1$. We show the following.

1. $\text{Res} \circ \text{Inf} = 0$: Change of group is functorial (easy to see from the definition), so $\text{Res} \circ \text{Inf}$ is induced by the maps $G/H \leftarrow G \leftarrow H$ and $M^H \hookrightarrow M \cong M$. The first map is 0 so $\text{Res} \circ \text{Inf} = 0$.
2. Inf is injective: Suppose $f : G/H \rightarrow A^H$ is a cocycle such that $\text{Inf}([f]) = 0$. Note $\text{Inf}([f]) = [f \circ p]$ where $p : G \rightarrow G/H$ is the projection. $\text{Inf}([f]) = 0$ means $f(s) = sa - a$ for some $a \in A$. Since f is constant on cosets, $sa - a = sta - a$ for all $t \in H$, giving $ta = a$, and $a \in A^H$. Thus $[f] = 0$ in $H^1(G/H, A^H) = 0$.
3. $\ker(\text{Res}) \subseteq \text{im}(\text{Inf})$: Suppose $f : G \rightarrow A$ is a cocycle such that $[f] \in \ker(\text{Res})$. Since $\text{Res}[f] = [f \circ i]$, this means $f(t) = ta - a$ for some $a \in A$ and all $t \in H$. Define the coboundary $g : G \rightarrow A$ by $g(s) = sa - a$ for all $s \in G$; let $f_1 = f - g$; we have $[f_1] = [f]$.

Now $f_1 = 0$ on H , and by definition of cocycle,

$$f_1(st) = f_1(s) + sf_1(t).$$

Letting t range over H , we get that $f_1(st) = f_1(s)$, i.e. f is constant on cosets of H . Letting $s \in H$ we have $f(st) = sf(t)$, so $\text{im}(f)$ is invariant under H . Thus f descends to $f : G/H \rightarrow A^H$, i.e. $f \in \text{im}(\text{Inf})$.

Now we proceed by induction. Suppose the proposition holds for $r-1$. By dimension-shifting (Proposition 12.9.8), the exact sequence

$$0 \rightarrow A \rightarrow A^* \rightarrow A^*/A \rightarrow 0 \quad (12.7)$$

with A^* coinduced gives $\partial^{n-1} : H_T^{r-1}(G, A^*/A) \xrightarrow{\cong} H_T^r(G, A)$. We now show there is a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^{r-1}(G/H, (A^*/A)^H) & \xrightarrow{\text{Inf}^{r-1}} & H^{r-1}(G, A^*/A) & \xrightarrow{\text{Res}^{r-1}} & H^{r-1}(H, A^*/A) \\ & & \downarrow \partial^{n-1} & & \downarrow \partial^{n-1} & & \downarrow \partial^{n-1} \\ 0 & \longrightarrow & H^r(G/H, A^H) & \xrightarrow{\text{Inf}^r} & H^r(G, A) & \xrightarrow{\text{Res}^r} & H^r(H, A). \end{array}$$

where all the vertical arrows are isomorphisms. We already know this for the middle arrow.

Since A^* is G -coinduced, it is H -coinduced (Proposition 12.8.6), so the right vertical arrow is an isomorphism.

Since $H^1(H, A) = 0$, taking cohomology of (12.7) gives the exact sequence

$$0 \rightarrow A^H \rightarrow (A^*)^H \rightarrow (A^*/A)^H \rightarrow 0.$$

Recall $A^* = \text{Hom}(\mathbb{Z}[G], A)$, so $(A^*)^H = \text{Hom}(\mathbb{Z}[G/H], A)$ is G/H -coinduced. Thus we get the left vertical arrow is an isomorphism.

By (cohomological) functoriality of Inf and Res , the diagram commutes. \square

11.5 Transfer

Especially important for our purposes will be the restriction map on the first homology group.

Definition 12.11.11: The map $V_{G \rightarrow S}$ defined by the diagram below

$$\begin{array}{ccc} H_1(G, \mathbb{Z}) & \xlongequal{\quad} & G/G' \\ \downarrow \text{Res}_1 & & \downarrow V_{G \rightarrow S} \\ H_1(S, \mathbb{Z}) & \xlongequal{\quad} & S/S' \end{array}$$

is called the **transfer** or **Verlagerung**.

(The map Res defined on Tate cohomology in Section 11.2 also gives a map on homology.)

Proposition 12.11.12: Let G be a group and S be a subgroup of finite index. The transfer is given by the following: Let $\{l_1, \dots, l_n\}$ be a left transversal of S in G . Then

$$\text{Res}_1(g) = \prod_{i=1}^n g_i S'$$

where the $g_i \in S$ are such that $gl_i = l_{\pi(i)}g_{\pi(i)}$ for some permutation $\pi \in S_n$.

Proof. By functoriality of Res we have the commutative diagram (cf. Proposition 12.8.3)

$$\begin{array}{ccccc}
 H_1(G, \mathbb{Z}) & \xrightarrow[\cong]{\partial_1} & H_0(G, I_G) & \xlongequal{\quad} & I_G/I_G^2 \\
 \downarrow \text{Res}_1 & & \downarrow \text{Res}_0 = C_{G/S} & & \downarrow C_{G/S} \\
 H_1(S, \mathbb{Z}) & \xrightarrow{\partial_1} & H_0(S, I_G) & \xlongequal{\quad} & I_G/I_S I_G \\
 & \searrow \cong & \uparrow & & \uparrow \\
 & & H_0(S, I_S) & \xlongequal{\quad} & I_S/I_S^2
 \end{array}$$

where the top two ∂_1 's are from the exact sequence $0 \rightarrow I_G \rightarrow \mathbb{Z}G \rightarrow \mathbb{Z} \rightarrow 0$, the bottom ∂_1 is from the exact sequence $0 \rightarrow I_H \rightarrow \mathbb{Z}H \rightarrow \mathbb{Z} \rightarrow 0$, and the lower right square is induced by the inclusion $I_H \hookrightarrow I_G$. Replacing H_1 with G^{ab} , we get

$$\begin{array}{ccc}
 G/G' & \xrightarrow{\cong} & I_G/I_G^2 \\
 \downarrow V_{G \rightarrow S} & & \downarrow C_{G/S} \\
 S/S' & \longrightarrow & I_G/I_S I_G \\
 & \searrow \cong & \uparrow \\
 & & I_S/I_S^2
 \end{array}$$

Given $g \in G/G'$, it maps to $g - 1$ in I_G/I_G^2 . We have

$$C_{G/S}(g-1) = \sum_{i=1}^n l_i^{-1}(g-1) = \sum_{i=1}^n g_i l_{\pi^{-1}(i)}^{-1} - l_i^{-1} = \sum_{i=1}^n i(g_i - 1) l_{\pi^{-1}(i)}^{-1} \equiv \sum_{i=1}^n (g_i - 1) \pmod{I_S I_G}.$$

The inverse image of this in S/S' is $\prod_{i=1}^n g_i S'$, as needed. \square

Theorem 12.11.13. *Let G be a finite group. Then the transfer map*

$$V : G^{\text{ab}} \rightarrow (G')^{\text{ab}}$$

is zero.

Proof. See Neukirch, [Neu99, VI.7.6]. The proof uses the computation in Proposition 12.11.12. \square

This will be important when we study the Hilbert class field.

12 Cohomology of cyclic groups

The cohomology of cyclic groups is especially easy to understand, and will be very useful to us: when L/K is an unramified extension of local fields, the Galois group $G(L/K) = G(l/k)$ is cyclic.

Theorem 12.12.1. *Let G be a cyclic group and x a generator. Let $\chi_x \in \text{Hom}(G, \mathbb{Q}/\mathbb{Z}) = H_T^1(G, \mathbb{Q}/\mathbb{Z})$ be the homomorphism sending x to $\frac{1}{|G|}$. Let $\delta : H_T^1(G, \mathbb{Q}/\mathbb{Z}) \rightarrow H_T^2(G, \mathbb{Z})$ be the diagonal map from the exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow 0$. The map $\bullet \cup \delta\chi_x$ gives an isomorphism*

$$H_T^r(G, M) \xrightarrow{\cong} H_T^{r+2}(G, M)$$

for all G -modules M and $r \in \mathbb{Z}$.

Hence for all $n \in \mathbb{Z}$,

$$\begin{aligned} H_T^{2n-1}(G, A) &= {}_N A / DA \\ H_T^{2n}(G, A) &= A^G / NA. \end{aligned}$$

where D is multiplication by $x - 1$.

Proof. Since \mathbb{Q} is a divisible group, so is $H^n(G, \mathbb{Q})$, by looking at the description of H^n in terms of cocycles (Section 7). Hence $\delta : H_T^1(G, \mathbb{Q}/\mathbb{Z}) \rightarrow H_T^2(G, \mathbb{Z})$ is an isomorphism and $\delta\chi_x$ is a generator of $H_T^2(G, \mathbb{Z})$.

The short exact sequence $0 \rightarrow I_G \rightarrow \mathbb{Z}G \rightarrow \mathbb{Z} \rightarrow 0$ splits because G is cyclic:

$$0 \rightleftarrows I_G \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{D} \end{array} \mathbb{Z}G \begin{array}{c} \xrightarrow{\varepsilon} \\ \xleftarrow{\quad} \end{array} \mathbb{Z} \rightleftarrows 0$$

where $\varepsilon(\sum_{g \in G} a_g g) = \sum_{g \in G} a_g$. Now $\mathbb{Z}G$ has trivial Tate cohomology by Proposition 12.9.7, so the diagonal maps in either direction are isomorphisms:

$$H_T^0(G, \mathbb{Z}) \xrightarrow[\cong]{\delta^0} H_T^1(G, I_G) \xrightarrow[\cong]{\delta^1} H_T^2(G, \mathbb{Z}).$$

Thus we can write $\delta\chi_x = \delta^0\delta^1c$ for a generator c of $H_T^0(G, \mathbb{Z}) = \mathbb{Z}/|G|\mathbb{Z}$. Then by Theorem 12.10.1(4),

$$b \cup \delta\chi_x = b \cup \delta^0\delta^1c = \delta^0\delta^1(b \cup c).$$

It suffices to show that the map $H_T^r(G, M) \xrightarrow{\bullet \cup c} H_T^r(G, M)$ is an isomorphism. But this map is just multiplication by c for $r = 0$, so it is multiplication by c for all r . Now by Proposition 12.11.6 (true for $r > 0$ and hence true for all r by dimension-shifting) $|G|H_T^r(G, M) = 0$. As c is a generator of $\mathbb{Z}/|G|\mathbb{Z}$ it is relatively prime to $|G|$; hence multiplication by c is an isomorphism on $H_T^r(G, M)$. This shows the isomorphism $H_T^r(G, M) \xrightarrow{\cong} H_T^{r+2}(G, M)$.

For the second part, note $H_T^{-1}(G, A) = {}_N A / DA$ and $H_T^0(G, A) = A^G / NA$. \square

Corollary 12.12.2. *Let G be a finite cyclic group. Suppose that $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$ is an exact sequence of G -modules. Then there is an exact hexagon*

$$\begin{array}{ccccc} & & H_T^0(G, A) & \xrightarrow{f_1} & H_T^0(G, B) & & \\ & & \nearrow f_6 & & \searrow f_2 & & \\ & H_T^1(G, C) & & & & H_T^0(G, C) & \\ & & \searrow f_5 & & \nearrow f_3 & & \\ & & H_T^1(G, B) & \xleftarrow{f_4} & H_T^1(G, A) & & \end{array} \quad (12.8)$$

Proof. We have $H_T^2(G, A) \cong H_T^0(G, A)$. □

12.1 Herbrand quotient

Definition 12.12.3: Let G be a finite cyclic group and A a finite G -module. Define the **Herbrand quotient** to be

$$h(A) = h(G, A) = \frac{|H_T^{2n}(G, A)|}{|H_T^{2n-1}(G, A)|}$$

for any n .

This is well-defined by Theorem 12.12.1.

The following key properties of the Herbrand quotient will help us in computations.

Proposition 12.12.4: Let G be a finite cyclic group. The Herbrand quotient satisfies the following.

1. If A is a finite G -module, then $h(G, A) = 1$.
2. (h is an Euler-Poincaré function) If $1 \rightarrow A \rightarrow B \rightarrow C \rightarrow 1$ is an exact sequence of G -modules, then

$$h(G, B) = h(G, A)h(G, C).$$

(If two of these are defined then the other is defined.)

3. If G acts trivially on \mathbb{Z} , then $h(G, \mathbb{Z}) = |G|$.
4. If $f : A \rightarrow B$ has finite kernel and cokernel, then $h(A) = h(B)$.

Proof. 1. We use Theorem 12.12.1 to calculate the quotient. We have the exact sequences

$$0 \longrightarrow {}_N A \longrightarrow A \xrightarrow{N} NA \longrightarrow 0 \qquad 0 \longrightarrow \ker D \longrightarrow A \longrightarrow DA \longrightarrow 0.$$

$$\qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \parallel$$

$$\qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad A^G$$

Hence

$$|NA| | {}_N A | = |A| = |A^G| |DA|,$$

giving

$$|H^1(G, A)| = |{}_N A / DA| = |A^G / NA| = |H^2(G, A)|.$$

2. Keeping the notation in the hexagon 12.8, we have

$$H^0(G, A) = |\ker f_1| \cdot \frac{|H^0(G, A)|}{|\ker f_1|} = |\operatorname{im} f_6| |\operatorname{im} f_1|.$$

We can similarly calculate the other quantities to get the result.

3. Let $|G| = n$, and $[n]$ denote multiplication by n . We have

$$h(G, \mathbb{Z}) = \frac{|H_T^0(G, \mathbb{Z})|}{|H_T^{-1}(G, \mathbb{Z})|} = \frac{|\mathbb{Z}^G/N\mathbb{Z}|}{|N\mathbb{Z}/I_G\mathbb{Z}|} = \frac{|\mathbb{Z}/n\mathbb{Z}|}{|\ker[n]|} = \frac{|G|}{1} = |G|.$$

4. The exact sequence $1 \rightarrow \ker f \rightarrow A \rightarrow B \rightarrow \text{coker } f \rightarrow 1$ gives $h(G, \ker f)h(G, B) = h(G, A)h(G, \text{coker } f)$ (split the exact sequence into 2 short exact sequences and use part 2). The result now follows from part 1. \square

13 Tate's Theorem

Theorem 12.13.1 (Tate's Theorem). *Let G be a finite group and M be a G -module. Suppose that for all subgroups $H \subseteq G$,*

1. $H^1(H, M) = 0$ and
2. $H^2(H, M)$ is cyclic of order $|H|$.

Then given a generator $u \in H^2(G, M)$, there is an isomorphism

$$H_T^r(G, \mathbb{Z}) \xrightarrow{\bullet \cup u} H_T^{r+2}(G, M)$$

for all r .

This is the main application of group cohomology to class field theory, as this will be the inverse of the Artin map: for instance, in local class field theory we have

$$\begin{aligned} H_T^{-2}(G(L/K), \mathbb{Z}) &= G(L/K)^{\text{ab}} \\ H_T^0(G(L/K), L^\times) &= (L^\times)^{G(L/K)} / \text{Nm}_{L/K}(L^\times) = K^\times / \text{Nm}_{L/K}(L^\times). \end{aligned}$$

The conditions of Tate's Theorem may seem unmotivated, but keep in mind that they are basically the key conditions satisfied in the number-theoretic setting, when G is taken to be a Galois group and M is taken to be a field (or idele group).

Class field theory was initially proved without group cohomology, but group cohomology gives a much nicer way to organize and abstract the proof. This theorem is a key part of that abstraction: isolating the key number-theoretic conditions that result in the Artin isomorphism. In proving both local and global class field theory, we will spend significant time showing that the hypothesis of Tate's Theorem holds. (The key difference in local and global class field theory is that we put in different things for M .)

Proof. Serre [Ser79], Section IX.8. \square

14 Profinite groups

In this section we study the cohomology groups when G is a profinite group. In this case topology becomes important. We will apply the results when G is an infinite Galois group.

We find that we have two ways of interpreting the resulting cohomology groups:

1. Imitate the previous construction but work in the category of topological G -modules instead. I.e. feed in “category of topological groups” into our cohomology functor.
2. Take the direct limit over finite quotients of G .

Definition 12.14.1: A **topological G -module** is a G -module that is a topological group, and such that the map

$$\begin{aligned} \varphi : G \times M &\rightarrow M \\ (g, m) &\mapsto gm \end{aligned}$$

is continuous.

We will always give M the discrete topology, so this is equivalent to the following condition:

$$M = \bigcup_{H \text{ open subgroup of } G} M^H.$$

Indeed, because M has the discrete topology, for the action to be continuous, $\pi_G(\varphi^{-1}(m))$ must be open, where $\pi_G : G \times M \rightarrow G$ is the projection. This is just the stabilizer of m , so the stabilizer of m must contain an open subgroup of G . Hence, every $m \in M$ must be contained in some M^H .

We define $H^n(G, M)$ as before, but now in the category of topological G -modules, i.e. we replace every instance of Hom_G with $\text{Hom}_G^{\text{cont}}$, since in this category the morphisms are *continuous* G -homomorphisms. Note that the category of discrete G -modules has enough injectives.

Theorem 12.14.2. *Let G be a profinite group. We have*

$$H^n(G, M) = \varinjlim H^n(G/S, M^S)$$

where the limit is over open normal subgroups S and the maps are the inflation maps

$$\text{Inf}^n : H^n(G/S, M^S) \rightarrow H^n(G/T, M^T), \quad S \supseteq T.$$

Proof. Milne [Mil08], II.4.2. □

We have a similar result if we take the limit over M .

Proposition 12.14.3: Let G be a profinite group and suppose $M = \varinjlim H^r(G, M_i)$ is a discrete G -module, and each M_i injects into M . Then

$$H^n(G, M) = \varinjlim H^n(G, M_i).$$

Proof. Milne [Mil08], II.4.4. □

15 Nonabelian cohomology

In this section we define cohomology $H^n(G, A)$ when A is *non-abelian*. (It was okay for G to be non-abelian because we saw it in the guise of $\mathbb{Z}G$, but we needed A to be in an abelian category.) The cohomological construction fails and we instead imitate the results of Theorem 12.7.2. (The description of H^1 and H^2 in Theorem 12.7.2 are useful because derivations and factor sets are used to classify a lot of things.)

We will only be able to get a “piece” of the long exact sequence. Cohomology also lacks a lot of structure: we speak not of cohomology groups, because they are now only pointed sets. We write A multiplicatively, as is the convention for nonabelian groups.

Definition 12.15.1: The category of **pointed sets** is the category whose objects are pairs (A, a) , where A is a set and $a \in A$, and such that a morphism $(A, a) \rightarrow (B, b)$ is a function $A \rightarrow B$ taking a to b .

The **kernel** of $f : (A, a) \rightarrow (B, b)$ is $f^{-1}(b)$. Thus we can define an exact sequence of pointed sets.

We now define the cohomology (pointed) sets. These will coincide with the definition in the abelian case by Theorem 12.7.2, except we only retain the structure of a pointed set.

Definition 12.15.2: Let G be a group and A a group with G -action.

1. Define

$$H^0(G, A) = A^G := \{a \in A : sa = a \text{ for all } s \in G\}.$$

The distinguished element is 1.

2. Define a **1-cocycle** to be a map $d : G \rightarrow A$ such that

$$d(xy) = d(x) \cdot xd(y)$$

and let $\text{Der}(G, A)$ be the set of 1-cocycles. Two cocycles d_1, d_2 are **cohomologous** if there exists $a \in A$ so that¹²

$$d_2(x) = a^{-1} \cdot d_1(x) \cdot xa.$$

Note this is an equivalence relation; define $H^1(G, A)$ to be the pointed set of 1-cocycles modulo equivalence. The distinguished element is the unit cocycle $d(x) \equiv 1$.

For an exact sequence of non-abelian G -modules

$$1 \rightarrow A \xrightarrow{i} B \xrightarrow{p} C \rightarrow 1$$

with $i(A) \trianglelefteq B$, define the **coboundary operator** $\delta : H^0(G, C) \rightarrow H^1(G, A)$ as follows: given $c \in C$, choose any $b \in p^{-1}(c)$ and set

$$\delta(c) = d \text{ where } d(s) = i^{-1}(b^{-1}s(b)).$$

¹²The analogue in the abelian case was $d_2(x) = -a + d_1(x) + xa$.

If furthermore $i(A)$ is in the center of B (so A is abelian), define $\Delta : H^1(G, C) \rightarrow H^2(G, A)$ as follows: for $d_c \in H^1(G, C)$, choose d_b such that $p_*d_b = d_c$, and set

$$[\Delta(d)](x, y) = d_b(s) \cdot s(d_b(t)) \cdot d_b(st)^{-1}.$$

Proof of well-definedness. Note the coboundary operator is defined by imitating the construction in the snake lemma.

$$\begin{array}{ccc}
 & & C^G \\
 & & \downarrow \\
 A & \xrightarrow{\quad i \quad} & B \xrightarrow{\quad p \quad} C \\
 \downarrow d_1 & & \downarrow d_1 \\
 \text{Der}(G, A) & \xrightarrow{\quad i \quad} & \text{Der}(G, B)
 \end{array}
 \qquad
 \begin{array}{ccc}
 & & c \\
 & & \downarrow \\
 b & \xrightarrow{\quad p \quad} & c \\
 \downarrow d_1 & & \\
 (s \mapsto i^{-1}(b^{-1}s(b))) & \xrightarrow{\quad i \quad} & (s \mapsto b^{-1}s(b))
 \end{array}$$

We need to show that $s \mapsto b^{-1}s(b)$ is actually a cocycle; its image is in A because $s(b) \equiv b^{-1} \pmod{i(A)}$ by exactness; show that the cohomology class is independent of the choice of b .

The second part is similar. Everything is easy to prove so we omit it. See Serre [Ser79], Appendix to Chapter VII. □

Theorem 12.15.3 (Exact sequence in nonabelian cohomology). *Let $1 \rightarrow A \xrightarrow{i} B \xrightarrow{p} C \rightarrow 1$ be an exact sequence of non-abelian G -modules. Then the following is exact.*

$$\begin{array}{ccccccccccc}
 1 & \longrightarrow & H^0(G, A) & \xrightarrow{i_0} & H^0(G, B) & \xrightarrow{p_0} & H^0(G, C) & \xrightarrow{\delta} & H^1(G, A) & \xrightarrow{i_1} & H^1(G, B) & \xrightarrow{p_1} & H^1(G, C) \\
 & & & & & & & & & & & & \downarrow \Delta \\
 & & & & & & & & & & & & H^2(G, A)
 \end{array}$$

(with the last map present if A is in the center of B).

Chapter 13

Introduction to Galois cohomology

We will apply group (co)homology as follows: Take a Galois extension L/K and let $G := G(L/K)$. Take as a G -module a multiplicative or additive subgroup S of L . The special case that G is cyclic will come up often, since if L/K is an unramified extension of local fields, then G is cyclic. Furthermore, the norm map N_G has a natural interpretation:

1. If $S \subseteq L^\times$ then for $a \in S$,

$$N_G(a) = \prod_{\sigma \in G} \sigma(a) = \text{Nm}_{L/K}(a).$$

2. If $S \subseteq L^+$ then for $a \in S$,

$$N_G(a) = \sum_{\sigma \in G} \sigma(a) = \text{tr}_{L/K}(a).$$

In Section 2 we give an application to Kummer theory (characterizing certain abelian extensions L of K in terms of $L^{\times n} \cap K$). Kummer theory will allow us to prove the linear independence of n th roots.

Finally, we give two interpretations of Galois cohomology groups.

1. $H^1(G(L/K), \text{Aut}(V))$ parameterizes algebraic structures defined over K that become isomorphic in L (Section 3). This is called *descent*.
2. $H^2(G(L/K), L^\times)$ parameterizes classes of K -algebras “split” over L (Section 4), i.e. it is the *Brauer group*.

1 Basic results

We prove two fundamental theorems on the cohomology of L^\times and L^+ .

Theorem 13.1.1 (Hilbert's Theorem 90). (†) *Let L/K be a Galois extension with Galois group G . Then*

$$H^1(G, L^\times) = \{1\}.$$

Moreover, if $G = \langle \sigma \rangle$ is cyclic and $u \in L^\times$, then the following are equivalent.

1. $\text{Nm}_{L/K}(u) = 1$.
2. *There exists $v \in L^\times$ such that $u = \sigma(v)v^{-1}$.*

We will often abbreviate $H^1(G(L/K), L^\times)$ as $H^1(L/K)$.

Proof. First suppose G is finite. Let $c : G \rightarrow L^\times$ be a 1-cocycle; we have $c_{\sigma\tau} = \sigma(c_\tau)c_\sigma$. Consider the function

$$b(e) := \sum_{\tau \in G} c_\tau \tau(e).$$

By linear independence of the characters $\tau \in G$, b is not identically zero; hence there exists $e \in L^\times$ so that $b(e) \neq 0$. Operating by σ on both sides and using the cocycle condition gives

$$\sigma(b(e)) = \sum_{\tau \in G} \sigma(c_\tau)(\sigma\tau)(e) = \sum_{\tau \in G} c_{\sigma\tau} c_{\sigma^{-1}}(\sigma\tau)(e) = c_\sigma^{-1} b(e) \quad (13.1)$$

and $c_\sigma = b(e)\sigma(b(e))^{-1}$, so c is a coboundary.

The infinite case follows from the finite case and Theorem 12.14.2.

For the second part, note that $H^1(G, L^\times) = \ker(N)/\text{im}(D) = 0$ gives $\ker(N) = \text{im}(D)$. Here N is the norm map $\text{Nm}_{L/K}$ and D is the map $\sigma - 1$, i.e. the map $v \mapsto \frac{\sigma(v)}{v}$. \square

Next we think of L as an additive group.

Theorem 13.1.2. *Let L/K be a finite Galois extension. Then*

$$H^r(G, L^+) = 0, \quad r > 0.$$

Proof. From the normal basis theorem 5.4.3, there exists $\alpha \in L$ such that $\{\sigma\alpha : \sigma \in G\}$ is a basis for L over K . We get that $K[G] \cong L$ as G -modules by the map

$$\sum_{\sigma \in G} a_\sigma \sigma \mapsto \sum_{\sigma \in G} a_\sigma \sigma\alpha.$$

Since $K[G] \cong \text{Ind}_{\{1\}}^G(K)$,

$$H^r(G, L^+) \cong H^r(\{1\}, K) = 0$$

by Shapiro's Lemma 12.8.7. \square

2 Kummer theory

We use Galois cohomology to prove the following.

Theorem 13.2.1 (Kummer theory). *Suppose K is a field containing a primitive n th root of 1. Then there is a bijection between*

1. *Finite abelian extensions of K of exponent dividing n (i.e. for any σ in the Galois group $G(L/K)$, $\sigma^n = 1$).*
2. *Subgroups of K^\times containing $K^{\times n}$ as a subgroup of finite index (i.e. subgroups of $K^\times/K^{\times n}$).*

This correspondence is given by

$$\begin{aligned} L &\mapsto K^\times \cap L^{\times n} \\ K[B^{\frac{1}{n}}] &\leftrightarrow B. \end{aligned}$$

Moreover,

$$[L : K] = [K^\times \cap L^{\times n} : K^\times] \tag{13.2}$$

(Note in the reverse map, which n th roots we take doesn't matter because K contains n th roots of unity.)

In the course of proving this theorem, we will show the following useful proposition.

Proposition 13.2.2: Let K be a field containing a primitive n th root of 1 and L/K an abelian extension with Galois group G . Then there is a natural isomorphism

$$\begin{aligned} K^\times \cap L^{\times n} / K^{\times n} &\cong H^1(G, \mu_n) = \text{Hom}(G, \mu_n) \\ a &\mapsto \left(\sigma \mapsto \frac{\sigma\left(a^{\frac{1}{n}}\right)}{a^{\frac{1}{n}}} \right). \end{aligned}$$

In particular, there is a natural isomorphism

$$K^\times / K^{\times n} \cong H^1(G(K^s/K), \mu_n) = \text{Hom}(G(K^s/K), \mu_n).$$

Proof. Let $G = G(L/K)$, and denote the forward map by $B(L) = K^\times \cap L^{\times n}$. The key step is showing that (13.2) holds; we do this by interpreting $K^\times \cap L^{\times n}$ as a 0th cohomology module. The inclusions $L \supseteq K(B(L)^{\frac{1}{n}})$ and $B(K(B^{\frac{1}{n}})) \supseteq B$ are easily seen to hold (Step 2), so (13.2) will give that equality holds (Steps 3-4).

Step 1: By Theorem 12.4.6, the short exact sequence of G -modules

$$1 \rightarrow \mu_n \rightarrow L^\times \xrightarrow{x \mapsto x^n} L^{\times n} \rightarrow 1$$

induces the long exact sequence

$$1 \rightarrow H^0(G, \mu_n) \rightarrow H^0(G, L^\times) \rightarrow H^0(G, L^{\times n}) \rightarrow H^1(G, \mu_n) \rightarrow H^1(G, L^\times) \rightarrow \dots$$

We need not go further because Hilbert's Theorem 90 (Theorem 13.1.1) tells us

$$H^1(G(L/K), L^\times) = 1.$$

Next, note that $H^0(G, H)$ is simply the subgroup of H fixed by G , and that the subfield of L fixed by G is K . As $\mu_n \subset K$, G acts trivially on μ_n and $H^1(G, \mu_n) = \text{Hom}(G, \mu_n)$ by Corollary 12.7.3. The sequence becomes

$$1 \rightarrow \mu_n \rightarrow K^\times \xrightarrow{x \mapsto x^n} K^\times \cap L^{\times n} \rightarrow \text{Hom}(G, \mu_n) \rightarrow 1,$$

giving an isomorphism

$$K^\times \cap L^{\times n} / K^{\times n} \cong \text{Hom}(G, \mu_n).$$

The map is $\partial^1(a) = \left(\sigma \mapsto \frac{\sigma(a^{\frac{1}{n}})}{a^{\frac{1}{n}}} \right)$, as shown by tracing through the construction in Theorem 12.3.3. This proves Proposition 13.2.2.

$$\begin{array}{ccc}
 & & K^\times \cap L^{\times n} \\
 & & \downarrow \\
 \mu_n & \xrightarrow{i} & L^\times \xrightarrow{x \mapsto x^n} L^{\times n} \\
 \downarrow d_1 & & \downarrow d_1 \\
 \text{Der}(G, \mu_n) & \xrightarrow{i} & \text{Der}(G, L^\times)
 \end{array}
 \qquad
 \begin{array}{ccc}
 & & a \\
 & & \downarrow \\
 a^{\frac{1}{n}} & \xrightarrow{x \mapsto x^n} & a \\
 \downarrow d_1 & & \downarrow d_1 \\
 \left(\sigma \mapsto \frac{\sigma(a^{\frac{1}{n}})}{a^{\frac{1}{n}}} \right) & \xrightarrow{i} & \left(\sigma \mapsto \frac{\sigma(a^{\frac{1}{n}})}{a^{\frac{1}{n}}} \right)
 \end{array}$$

We claim that $|\text{Hom}(G, \mu_n)| = |G|$. Indeed, by the structure theorem for abelian groups, G decomposes as $(\mathbb{Z}/n_1\mathbb{Z}) \times \dots \times (\mathbb{Z}/n_m\mathbb{Z})$ where $n_1, \dots, n_m \mid n$. To choose a homomorphism for G means choosing images for the generators of $\mathbb{Z}/n_1\mathbb{Z}, \dots, \mathbb{Z}/n_m\mathbb{Z}$; there are n_1, \dots, n_m possibilities, respectively, for a total of $|G|$.

Then

$$[L : K] = |G(L/K)| = [K^\times \cap L^{\times n} : K^\times].$$

This shows (13.2).

Step 2: Next note the following two inclusions.

1. $K[B(L)^{\frac{1}{n}}] \subseteq L$: Anything in $(K^\times \cap L^{\times n})^{\frac{1}{n}}$ is in the form $(\beta^n)^{\frac{1}{n}}$ and hence in L .
2. $B(K[B^{\frac{1}{n}}]) \supseteq B$: Anything in B is in the form $(b^{\frac{1}{n}})^n$ and hence in $K^\times \cap K(B^{\frac{1}{n}})^{\times n}$.

Step 3: We show that $K[B(L)^{\frac{1}{n}}] = L$. By the inclusions in step 2,

$$[L : K] \geq [K[B(L)^{\frac{1}{n}}] : K] \stackrel{(13.2)}{=} [B(K[B(L)^{\frac{1}{n}}]) : K^\times] \geq [B(L) : K^\times].$$

But $[L : K] = [B(L) : K^\times]$ by (13.2), so equality holds everywhere. The first equality gives the conclusion.

Step 4: We show that $B(K[B^{\frac{1}{n}}]) = B$. We apply step 1 with $L = K[B^{\frac{1}{n}}]$ to get the isomorphism

$$B(L) = K^\times \cap L^{\times n} / K^{\times n} \xrightarrow{\cong} \text{Hom}(G, \mu_n)$$

$$a \mapsto \left(\sigma \mapsto \frac{\sigma(a^{\frac{1}{n}})}{a^{\frac{1}{n}}} \right).$$

Now $B \subseteq B(L)$ gets mapped to a subgroup $H' \subseteq \text{Hom}(G, \mu_n)$, which can be identified with $\text{Hom}(G/H, \mu_n)$ ¹. But as the $a^{\frac{1}{n}}$ generate L over K and the fixed field of G is K , $\bigcap_{h \in H'} \ker h = 1$. Thus $H = \{1\}$. Hence $|B(L)| = |G| = |B|$, and $B = B(L)$. \square

Corollary 13.2.3 (*n*th roots are linearly independent). *Let S be a set of nonzero integers so that $\frac{a}{b}$ is not a perfect *n*th power for any distinct $a, b \in S$. Then the elements*

$$\sqrt[n]{s}, \quad s \in S$$

are linearly independent over \mathbb{Q} .

Proof. Step 1: It suffices to show that for distinct primes p_1, \dots, p_k , we have

$$[\mathbb{Q}(\sqrt[n]{p_1}, \dots, \sqrt[n]{p_k}) : \mathbb{Q}] = n^k. \quad (13.3)$$

Then a basis for this extension over \mathbb{Q} is formed by taking products of basis elements for the $\mathbb{Q}(\sqrt[n]{p_j})$:

$$\left\{ \sqrt[n]{p_1^{a_1} \cdots p_k^{a_k}} : 0 \leq a_j < n \right\}. \quad (13.4)$$

However, the radicands are exactly the representatives of elements in $\mathbb{Q}^\times / \mathbb{Q}^{\times n}$. The elements of S are all represented by distinct elements of (13.4) modulo \mathbb{Q}^\times , so the theorem will follow. (To deal with $s \in S$ negative, note if s is negative then $\sqrt[n]{s}$ is not in \mathbb{R} .)

We want to use Kummer theory to conclude (13.3). However, \mathbb{Q} only has square roots of unity (± 1), so we have to consider all other roots separately. We may as well assume $2 \mid n$.

¹ The subgroups of G are in bijective correspondence with the subgroups of $\text{Hom}(G, \mu_n)$ via the map

$$H \xrightarrow{\Phi} \{h \in \text{Hom}(G, \mu_n) : H \subseteq \ker h\} \cong \text{Hom}(G/H, \mu_n)$$

$$\bigcap_{h \in H'} \ker h \xleftarrow{\Psi} H'$$

Indeed, clearly $\Psi(\Phi(H)) \supseteq H$, and we have equality since for every $g \in G \setminus H$ we can find $h \in \text{Hom}(G, \mu_n)$ with kernel containing H , so that $h(g) \neq 1$. Since $\text{Hom}(G, \mu_n) \cong G$, they have the same number of subgroups, and this is a bijection.

Step 2: We first show

$$[\mathbb{Q}(\sqrt{p_1}, \dots, \sqrt{p_k}) : \mathbb{Q}] = 2^k. \quad (13.5)$$

Let B be the subgroup of \mathbb{Q}^\times generated by p_1, \dots, p_k and $\mathbb{Q}^{\times 2}$. By Theorem 13.2.1,

$$[\mathbb{Q}(B^{\frac{1}{2}}) : \mathbb{Q}] = [B : \mathbb{Q}^{\times 2}] = 2^k,$$

as needed.

Step 3: We now adjoin n th roots of unity such that we can apply Kummer theory for n th roots. Let N be a positive integers such that $n \mid N$ and $\mathbb{Q}(\sqrt{p_1}, \dots, \sqrt{p_k}) \subseteq \mathbb{Q}(\zeta_N)$ (every quadratic extension is contained in a cyclotomic extension; we can take $N = 4p_1 \cdots p_k n$).

However, if we look at $K := \mathbb{Q}(\zeta_N)$, what if elements that aren't n th powers in \mathbb{Q} become n th powers? Fortunately, this doesn't happen for $n \neq 2$. We show that for even $n \neq 2$ and $m \in \mathbb{Q}$ not a perfect $\frac{n}{2}$ th power,

$$\sqrt[n]{m} \notin \mathbb{Q}(\zeta_N). \quad (13.6)$$

By taking roots, we may assume that m is not a perfect d th power for any $d \mid n$.

Note $L := \mathbb{Q}(\sqrt[n]{m}, \zeta_n)$ is a Galois extension of \mathbb{Q} since it is the splitting field of $X^n - m$. Note $X^n - m$ is irreducible over \mathbb{Q} because the constant term of any proper factor must be in the form $m^{\frac{j}{n}} \notin \mathbb{Q}$ where $0 < j < n$. Hence there exists $\tau \in G(L/\mathbb{Q})$ sending $\sqrt[n]{m}$ to $\zeta_n \sqrt[n]{m}$. Let $\sigma \in G(L/\mathbb{Q})$ denote complex conjugation. Then

$$\begin{aligned} \sigma\tau(\sqrt[n]{m}) &= \sigma(\zeta_n \sqrt[n]{m}) = \zeta_n^{-1} \sqrt[n]{m} \\ \tau\sigma(\sqrt[n]{m}) &= \tau(\sqrt[n]{m}) = \zeta_n \sqrt[n]{m}. \end{aligned}$$

Hence $G(L/\mathbb{Q})$ is not abelian. Since all cyclotomic extensions are abelian, L cannot be contained in an abelian extension, giving (13.6).

Let C be the subgroup of $\mathbb{Q}(\zeta_N)^\times$ generated by $\sqrt{p_1}, \dots, \sqrt{p_k}$ and $\mathbb{Q}(\zeta_N)^{\times \frac{n}{2}}$. We showed above that $\sqrt[n]{m} \notin \mathbb{Q}(\zeta_N)^{\times \frac{n}{2}}$ for any m not a perfect $\frac{n}{2}$ th power so $[C : \mathbb{Q}(\zeta_N)^{\times \frac{n}{2}}] = \left(\frac{n}{2}\right)^k$. By Kummer Theory,

$$[\mathbb{Q}(\zeta_N, \sqrt[n]{p_1}, \dots, \sqrt[n]{p_k}) : \mathbb{Q}(\zeta_N)] = [\mathbb{Q}(C^{\frac{n}{2}}) : K] = [C : \mathbb{Q}(\zeta_N)^{\times \frac{n}{2}}] = \left(\frac{n}{2}\right)^k.$$

Since $\mathbb{Q}(\sqrt{p_1}, \dots, \sqrt{p_k}) \subseteq \mathbb{Q}(\zeta_N)$ we get

$$[\mathbb{Q}(\sqrt[n]{p_1}, \dots, \sqrt[n]{p_k}) : \mathbb{Q}(\sqrt{p_1}, \dots, \sqrt{p_k})] = \left(\frac{n}{2}\right)^k. \quad (13.7)$$

Combining (13.5) and (13.7) gives (13.3), as needed. \square

3 Nonabelian Galois cohomology

Because of the definition of H^1 in Section 15, we find that we can often interpret $H^1(G(L/K), A)$ as parameterizing certain algebraic structures, specifically a set of them defined over K that

become isomorphic in L . (This is known as *descent* because it answers the question, how many ways can an algebraic structure (or in general, a variety) “descend” from L to K ?) In general,

$$\begin{aligned} H^1(G(L/K), \{\text{automorphisms preserving } V \text{ over } K\}) \\ \cong \{K\text{-isomorphism classes that are } L\text{-congruent to } V\} \quad (13.8) \end{aligned}$$

In this section, we will see several examples where A is an algebraic group. We could also take A to be an abelian variety (see Silverman [Sil86], Theorem X.2.2, for instance).

In particular, we find in the next section that a special cohomology group classifies algebra structures over K : the Brauer group.

First, we need the following nonabelian generalization of Hilbert’s Theorem 90 (13.1.1).

Theorem 13.3.1 (Generalization of Hilbert’s Theorem 90). *For any finite Galois extension L/K , letting $G = G(L/K)$,*

$$H^1(G, \text{GL}_n(L)) = H^1(G, \text{SL}_n(L)) = 1.$$

Proof. As in Theorem 13.1.1, given a 1-cocycle $c : G \rightarrow \text{GL}_n(L)$, consider the function

$$\begin{aligned} b : \text{GL}_n(L) &\rightarrow \mathcal{M}_n(L) \\ b(A) &:= \sum_{\tau \in G} c_\tau \tau(A). \end{aligned}$$

Note that unlike in the proof of Theorem 13.1.1, we not only have to choose A to be nonzero, but also invertible.

Also define b on L^n in the same way:

$$\begin{aligned} b : L^n &\rightarrow L^n \\ b(\mathbf{x}) &:= \sum_{\tau \in G} c_\tau \tau(\mathbf{x}). \end{aligned}$$

We show that $\{b(\mathbf{x}) : \mathbf{x} \in L^n\}$ generate L^n as a vector space over L .² Suppose a linear functional $f : L^n \rightarrow L$ vanishes on all the $b(\mathbf{x})$. Then for every $\alpha \in L$,

$$0 = f(b(\alpha\mathbf{x})) = \sum_{\tau \in G} f(c_\tau \tau(\alpha)\tau(\mathbf{x})) = \sum_{\tau \in G} \tau(\alpha) f(c_\tau \tau(\mathbf{x})).$$

By linear independence of characters, we get that all the coefficients of the $\tau(\alpha)$ must be 0, i.e. $f(c_\tau \tau(\mathbf{x}))$ for all c_τ, \mathbf{x} . But $c_\tau \in \text{GL}_n(L)$ is invertible, so f must vanish identically on L^n . We’ve shown that every linear functional vanishing on $\{b(\mathbf{x})\}$ vanishes on L^n ; therefore $\text{span}_L \{b(\mathbf{x})\} = L^n$.

Thus we can choose $\mathbf{x}_1, \dots, \mathbf{x}_n$ such that $\mathbf{y}_j = b(\mathbf{x}_j)$ form a basis for L^n over L . Let A be the matrix sending the canonical basis \mathbf{e}_j to the \mathbf{x}_j . Then (note τ acts trivially on the e_j)

$$b(A)\mathbf{e}_j = b(A\mathbf{e}_j) = \mathbf{y}_j$$

²Note b is not a L -linear transformation; it is a K -linear transformation.

so $b(A)$ is invertible.

The the rest of the proof of Theorem 13.1.1 goes through: we have as in (13.1) that

$$c_\sigma = b(A)\sigma(b(A))^{-1},$$

i.e. c is a coboundary. This shows $H^1(G, \text{GL}_n(L)) = 1$.

For the second part, the exact sequence

$$1 \rightarrow \text{SL}_n(L) \rightarrow \text{GL}_n(L) \xrightarrow{\det} L^\times \rightarrow 1$$

gives the long exact sequence 12.15.3

$$\begin{array}{ccccccc} H^0(G, \text{GL}_n(L)) & \xrightarrow{\det} & H^0(G, L^\times) & \longrightarrow & H^1(G, \text{SL}_n(L)) & \longrightarrow & H^1(G, \text{GL}_n(L)) \\ \parallel & & \parallel & & & & \parallel \\ \text{GL}_n(K) & \xrightarrow{\det} & K^\times & & & & 0. \end{array}$$

As the map on the left is surjective, we get $H^1(G, \text{SL}_n(L)) = 1$. □

We have now established (13.8) when V is a vector space: all vector spaces that become isomorphic in L have the same dimension to begin with so are isomorphic in K , so the right-hand side of (13.8) is $\{1\}$, and if $V = K^n$, $\text{GL}_n(L)$ is the group of automorphisms preserving V over L , and Theorem 13.3.1 shows the right-hand side of (13.8) is $\{1\}$. We now extend this to other algebraic structures.

To encode an algebraic structure, we consider vector spaces and tensors.

Example 13.3.2:

Let V be a finite-dimensional vector space. The space $V^{\otimes p} \otimes V^{*\otimes q}$ encodes...

p	q	Structure
1	0	vectors
0	1	linear functionals
1	1	linear operators
0	2	bilinear forms
1	2	algebra structures

We focus on the case $p = 1, q = 2$. Given a tensor $\sum_i v_i \otimes f_i \otimes g_i \in V \otimes V^{*\otimes 2}$, define a (not necessarily commutative or associative) algebra structure on V by

$$v \cdot w = \sum_i f_i(v)g_i(w)v_i.$$

Conversely, any algebra structure can be encoded in this way: Take a basis $\{v_i\}$ for V and a dual basis f_i for V^* , and encode the structure by $\sum_{i,j}(v_i \cdot v_j) \otimes f_i \otimes g_j$.

Definition 13.3.3: Let V be a vector space over K and $x \in V^{\otimes p} \otimes V^{*\otimes q}$ be a tensor of type (p, q) . Two pairs (V, x) and (V', x') are isomorphic if there is a K -linear isomorphism

$$f : V \rightarrow V'$$

such that $f(x) = x'$. Here, f sends

$$x_1 \otimes \cdots \otimes x_p \otimes f_1 \otimes \cdots \otimes f_q \mapsto f(x_1) \otimes \cdots \otimes f(x_p) \otimes (f_1 \circ f^{-1}) \otimes \cdots \otimes (f_q \circ f^{-1}). \quad (13.9)$$

Given (V, x) defined over K , we can consider it over L by extending scalars; denote the resulting pair by $(V_L = V \otimes_K L, x_L)$.

We say that (V, x) and (V', x') are L -isomorphic if (V_L, x_L) and (V'_L, x'_L) are isomorphic. Let $E_{V,x}(L/K)$ denote the L -isomorphism classes of pairs that are K -equivalent to (V, x) . If L/K is Galois, let $s \in G(L/K)$ act on $v \otimes \alpha \in V \otimes_K L = V_L$ by $s(v \otimes_K \alpha) := v \otimes_K s(\alpha)$ and let s act on A_L by conjugation:

$$f^s := s \circ f \circ s^{-1}.$$

Theorem 13.3.4 (Descent for tensors). *Let L/K be a Galois extension, $G = G(L/K)$, and let A_L be the group of L -automorphisms of (V_L, x_L) . Define the map*

$$\begin{aligned} \theta : E_{V,x}(L/K) &\rightarrow H^1(G, A_L) \\ (V', x') &\mapsto (d : \sigma \mapsto f^{-1} \circ f^\sigma = f^{-1} \circ \sigma \circ f \circ \sigma^{-1}) \end{aligned}$$

where $f : (V_L, x_L) \rightarrow (V'_L, x'_L)$ is any L -automorphism. Then θ is a bijection.

Proof. We show the following.

1. θ is well-defined. First, $\theta(V', x')$ is a cocycle as

$$d(\sigma t) = f^{-1} \sigma t f t^{-1} \sigma^{-1} = (f^{-1} \sigma f \sigma^{-1}) [\sigma (f^{-1} t f t^{-1}) \sigma^{-1}] = d(\sigma) \circ d(t)^\sigma.$$

(See Definition 12.15.2.) Next, we show $\theta(V', x')$ does not depend on the choice of f : Let $d_f(\sigma) = f^{-1} \sigma f \sigma^{-1}$ and $d_g(s) = g^{-1} \sigma g \sigma^{-1}$. Then

$$d_g(\sigma) = g^{-1} \sigma g \sigma^{-1} = g^{-1} f (f^{-1} \sigma f \sigma^{-1}) \sigma f^{-1} g \sigma^{-1} = (f g^{-1})^{-1} d_f(\sigma) (f g^{-1})^\sigma$$

so d_f and d_g are cohomologous.

2. θ is injective. Suppose $\theta(V'_1, x'_1) = \theta(V'_2, x'_2)$. We can choose the isomorphisms f_1 and f_2 such that $f_1^{-1} f_1^\sigma = f_2^{-1} f_2^\sigma$ for all $\sigma \in G(L/K)$. Then $(f_2 f_1^{-1})^\sigma = f_2 f_1^{-1}$ for all $\sigma \in G(L/K)$, i.e. $f_2 f_1^{-1}$ is an isomorphism defined over K . Thus (V'_1, x'_1) and (V'_2, x'_2) are K -isomorphic.
3. θ is surjective. Let c_σ be a 1-cocycle of G with values in A_L . Since $A_L \subseteq \text{GL}(V_L)$, by Theorem 13.3.1 there exists $f \in \text{GL}(V_L)$ such that

$$c_\sigma = f^{-1} \circ f^\sigma$$

Let f operate on $V^{\otimes p} \otimes V^{*\otimes q}$ as in (13.9) and let $x' = f(x)$. As $x \in V_K^{\otimes p} \otimes V_K^{*\otimes q}$ and c_σ fixes K , we have

$$\sigma(x') = f^\sigma(\sigma(x)) = f^\sigma(x) = f \circ c_\sigma(x) = f(x) = x'.$$

Thus x' is rational over K (i.e. in $V_K^{\otimes p} \otimes V_K^{*\otimes q}$), and (V, x') maps to c_σ .

□

Note that since we always take an isomorphism $V \rightarrow V'$, we can really consider all the vector spaces to be the “same,” and just vary the tensors x . If we consider $V = V'$, then we abbreviate $f : (V_L, x_L) \rightarrow (V'_L, x'_L)$ by $f : x \rightarrow x'$.

Example 13.3.5: We can use Galois cohomology to classify quadratic forms over a field K . Let Φ be a quadratic form (which corresponds to a bilinear form and can be represented by a tensor of type $(0, 2)$), and $O_L(\Phi)$ be the orthogonal group of Φ , i.e. linear transformations that preserve Φ . Then $H^1(G(L/K), O_L(\Phi))$ classifies the quadratic forms over K that are L -isomorphic to Φ .

4 Brauer group

The Brauer group characterizes algebras over a field K . We already know a simple way of making algebras: just consider the algebra of $n \times n$ matrices, $\mathcal{M}_n(K)$. Thus, we will essentially “mod out” by these when constructing the group.

As we will see, there is an isomorphism to a second cohomology group. Thus, we can apply results about algebras over K to Galois cohomology, or conversely, apply Galois cohomology to get information on algebras over K .

First, we need some results from noncommutative algebra. We refer the reader to Cohn [Coh03], Chapter 5, or Milne [Mil08], Chapter IV.1–2, for the proofs.

4.1 Background from noncommutative algebra

Definition 13.4.1: An **algebra** over a field K is a ring A with K in its center³. Its dimension is the dimension of A as a K -vector space, denoted $[A : K]$. *In this chapter we assume all algebras to be finite-dimensional as K -vector spaces.*

An algebra over K is

1. **simple** if it has no proper two-sided ideals.
2. **central** if its center is K .

An algebra is a **division algebra** if every nonzero element has an inverse.

³The center of a ring R is the set of elements commuting with all elements of R .

Example 13.4.2: The algebra of $n \times n$ matrices $\mathcal{M}_n(K)$ is a central simple algebra over K .

Definition 13.4.3: Let A be an algebra over K . We use “ A -module” to mean any finitely generated left A -module V ; the map $A \rightarrow \text{End}(V)$ is called a **representation** of A . The module (or representation) is **faithful** if $av = 0$ for all $v \in V$ implies $a = 0$, i.e. $A \hookrightarrow \text{End}(V)$ is injective. A module is **simple** if it doesn’t contain a proper A -submodule, and **indecomposable** if it is not the direct sum of two proper A -submodules. (Note that simple implies indecomposable, but not vice versa.) A module is **semisimple** if it is the direct sum of simple A -modules.⁴

We say A is semisimple if it is semisimple as a module.

We need some basic results from noncommutative algebra.

Definition 13.4.4: Let $B \subseteq A$ be a subalgebra. Define the **centralizer** of B to be the elements of A commuting with B :

$$C(B) := \{a \in A : ab = ba \text{ for all } b \in B\}.$$

Theorem 13.4.5 (Double centralizer theorem). *Let A be a K -algebra, and V a faithful semisimple A -module. Consider A as a subalgebra of $\text{End}_K(V)$. Then*

$$C(C(A)) = A.$$

Proof. Milne [Mil08], Theorem IV.1.3, or Etingof [Eti], Theorem 4.54. □

Theorem 13.4.6 (Wedderburn’s structure theorem). *An algebra A is semisimple iff it is isomorphic to the direct sum of matrix algebras over division algebras.*

If A is an algebra over an algebraically closed field K and K , then any semisimple algebra over K is isomorphic to a direct sum of matrix algebras over K .

Proof. Milne [Mil08], Theorem IV.1.15.

For the second part, we need to show the only division algebra over an algebraically closed field K is K itself. Suppose D is a division algebra and $\alpha \in D$. As $[D : K]$ is finite-dimensional, $K(\alpha)$ is a finite extension of K . Hence $\alpha \in K$, giving $D = K$. □

Theorem 13.4.7 (Noether-Skolem theorem). *Let $f, g : A \rightarrow B$ be homomorphisms, where A is a simple K -algebra and B is a central simple K -algebra. Then there exists $b \in B$ such that*

$$f(a) = b \cdot g(a) \cdot b^{-1}$$

for all $a \in A$, i.e. f, g differ by an inner automorphism of B .

In particular, taking $A = B$ and $g = 1$, all automorphisms of a central simple K -algebra are inner (come from conjugation). In particular, this is true for $\mathcal{M}_n(K)$.

⁴Equivalently, the radical of A is trivial. If it is semisimple the factors in the decomposition are unique up to isomorphism (Jordan-Hölder).

4.2 Central simple algebras and the Brauer group

We now define the Brauer group.

Definition 13.4.8: Let A and B be simple algebras over K . We say A and B are similar and write $A \sim B$ if

$$A \otimes_K \mathcal{M}_m(K) \cong B \otimes_K \mathcal{M}_n(K)$$

for some m, n .

The **Brauer group** Br_K is the set of similarity classes of central simple algebras over K , with multiplication defined by

$$[A][B] = [A \otimes_K B].$$

The Brauer group $\text{Br}_{L/K}$ is the subgroup of classes of central simple algebras over K that are **split** over L , i.e. such that $A \otimes_K L$ is a matrix algebra.

Proof (sketch) that this is a group. We need to check that...

1. The tensor product of two central simple algebras is central simple. By Wedderburn's Theorem 13.4.6 we can write the algebras as $A = M_m(D)$ and $B = M_{m'}(D')$, where D, D' are division algebras. One can show $A \otimes_K D'$ is simple; hence it equals $M_n(D'')$ for some D'' . Then $A \otimes_K B \cong M_{m'n}(D'')$ is simple. It is central because $C(A \otimes_K B) = C(A) \otimes_K C(B) = K$.

2. “ \sim ” is an equivalence relation. If $A \sim B$ and $B \sim C$, then $A \otimes_K \mathcal{M}_m(K) \cong B \otimes_K \mathcal{M}_n(K)$, $B \otimes_K \mathcal{M}_{n'}(K) \cong C \otimes_K \mathcal{M}_p(K)$ for some m, n, n', p . Then

$$A \otimes_K \mathcal{M}_{mn'}(K) \cong A \otimes_K \mathcal{M}_m(K) \otimes_K \mathcal{M}_{n'}(K) \cong C \otimes_K \mathcal{M}_n(K) \otimes_K \mathcal{M}_p(K) \cong C \otimes_K \mathcal{M}_{np}(K).$$

3. “ \sim ” is preserved under the operation \otimes . If $A_i \otimes_K \mathcal{M}_{m_i}(K) \cong B_i \otimes_K \mathcal{M}_{n_i}(K)$ for $i = 1, 2$, then $A_1 \otimes_K A_2 \otimes_K \mathcal{M}_{m_1 m_2}(K) \cong B_1 \otimes_K B_2 \otimes_K \mathcal{M}_{n_1 n_2}(K)$.

4. A has an inverse. Letting A^{opp} be the opposite algebra, we find that

$$A \otimes_K A^{\text{opp}} \cong \mathcal{M}_n(K), \quad n = [A : K].$$

5. The operation is commutative and associative. This follows since tensor product is commutative and associative.

□

By Wedderburn's Structure Theorem 13.4.6, each (central) simple algebra is $M_n(D) \cong M_n(K) \otimes_K D$ for some (central) division algebra D , so every similarity class is represented by a central division algebra. Thus to determine the Brauer group it suffices to classify central division algebras.

Example 13.4.9: We have

$$\mathrm{Br}_{\mathbb{R}} = \{\mathbb{R}, \mathbb{H}\}$$

where \mathbb{H} denotes the quaternions: the algebra with basis $1, \mathbf{i}, \mathbf{j}, \mathbf{k} = \mathbf{ij}$ and relations $\mathbf{i}^2 = 1$, $\mathbf{j}^2 = 1$, and $\mathbf{ij} = -\mathbf{ji}$.

Indeed, by Frobenius's Theorem, the only finite-dimensional (associative) division algebras over \mathbb{R} are \mathbb{R} , \mathbb{C} , and \mathbb{H} , and only \mathbb{R} and \mathbb{H} have center equal to \mathbb{R} .

Proposition 13.4.10: For any algebraically closed field K ,

$$\mathrm{Br}_{\overline{K}} = 0.$$

Proof. By Wedderburn's Theorem 13.4.6, all central simple algebras over K are $\mathcal{M}_n(K)$ for some n . □

4.3 Subfields and splitting of central simple algebras

An important way of studying a central simple algebra is to look at its subfields.

Theorem 13.4.11 (Double centralizer theorem, generalization). *Let A be a central simple K -algebra and B be a simple K -subalgebra. Let $C = C(B)$. Then C is simple, $C(C) = A$, and*

$$[B : K][C : K] = [A : K].$$

Proof. See Milne [Mil08], Theorem IV.3.1. □

Corollary 13.4.12. *Let A be central simple over K , and L be a subfield with $K \subseteq L \subseteq A$. Then the following are equivalent.*

1. $L = C(L)$.
2. $[A : K] = [L : K]^2$.
3. L is the maximal commutative K -subalgebra of A .

Proof. Milne [Mil08], Corollary IV.3.4. □

The following describes the fields over which a central simple algebra splits.

Corollary 13.4.13. *Let A be central simple over K . A finite extension field M splits A iff there exists an algebra $B \sim A$ containing M with $[B : K] = [L : K]^2$. In particular, any subfield L of A of degree $\sqrt{[A : K]}$ splits A .*

If D is a division algebra of degree n^2 over K , and L is a field of degree n over K (equivalently a maximal commutative subfield of D), then L splits D , i.e. $D \cong \mathcal{M}_n(L)$.

Proof. Milne [Mil08], IV.3.6, and 3.7. □

Theorem 13.4.14. *Every central division algebra over K is split over some finite Galois extension L/K . Therefore*

$$\mathrm{Br}_K = \mathrm{Br}_{\overline{K}/K} = \bigcup_{L/K \text{ finite Galois}} \mathrm{Br}_{L/K}.$$

Proof. When K is perfect, this follows directly from Corollary 13.4.13. The general case requires a separate argument; see Milne [Mil08], IV.3.10. \square

Similar to the commutative case, we can define a valuation on division algebras.

Proposition 13.4.15: Let D be a division algebra of rank n^2 over a local field K . Then D admits a discrete valuation extending the valuation on K , such that for any $a \in (0, 1)$, $\|x\|_D := a^{v(x)}$ defines a norm on D . The set of integral elements $\{x : v(x) \geq 0\}$ is a subring of D .

5 Brauer group and cohomology

5.1 The Brauer group is a second cohomology group

Definition 13.5.1: Let $\mathrm{Br}_{L/K,n}$ denote the subset of $\mathrm{Br}_{L/K}$ consisting of $[A]$ where $A \otimes_K L \cong \mathcal{M}_n(L)$. Note that $\mathrm{Br}_{L/K} = \bigcup_{n \in \mathbb{N}} \mathrm{Br}_{L/K,n}$.

Theorem 13.5.2 (Cohomological interpretation of Brauer group). *There are canonical bijections*

$$\theta_n : \mathrm{Br}_{L/K,n} \rightarrow H^1(G, \mathrm{PGL}_n(K))$$

and canonical isomorphisms

$$\begin{aligned} \delta : \mathrm{Br}_{L/K} &\rightarrow H^2(L/K) \\ \delta : \mathrm{Br}_K &\rightarrow H^2(K) \end{aligned}$$

where $H^2(K) := H(\overline{K}/K) = \varinjlim_{L/K \text{ finite Galois}} H^2(L/K)$.

Proof. We can represent elements of $\mathrm{Br}_{L/K,n}$ as algebras of dimension n^2 over K , that are L -isomorphic to the algebra $\mathcal{M}_n(L)$. By Example 13.3.2, we can encode the algebra $\mathcal{M}_n(L)$ by a tensor of type (1,2). By Theorem 13.3.4,

$$\mathrm{Br}_{L/K,n} \cong H^1(G, \mathrm{Aut}(\mathcal{M}_n(L))). \tag{13.10}$$

By the Noether-Skolem Theorem 13.4.7, every automorphism of $\mathcal{M}_n(L)$ is conjugation by an element of $\mathrm{GL}_n(K)$. Since the matrices that act trivially by conjugation are just the scalar matrices, we have the short exact sequence

$$1 \rightarrow L^\times \rightarrow \mathrm{GL}_n(L) \rightarrow \mathrm{Aut}(\mathcal{M}_n(L)) \cong \mathrm{PGL}_n(L) \rightarrow 1. \tag{13.11}$$

Along with (13.10) this proves the first part.

The long exact sequence 12.15.3 of (13.11) gives

$$0 = H^1(G, \mathrm{GL}_n(L)) \rightarrow H^1(G, \mathrm{PGL}_n(L)) \xrightarrow{\Delta_n} H^2(G, L^\times),$$

where the LHS follows from Theorem 13.3.1. Let $\delta_n = \Delta_n \circ \theta_n$; then δ_n is an injective map. We show the following.

1. The δ_n for different n combine compatibly into an injective group homomorphism $\delta : \mathrm{Br}(L/K) \rightarrow H^2(L/K)$: We need to show

$$\delta_{nn'}(A \otimes A') = \delta_n(A)\delta_{n'}(A')$$

for any $A \in \mathrm{Br}_{L/K,n}$ and $A' \in \mathrm{Br}_{L/K,n'}$.

First, note that if a, a' are tensors encoding the algebras A, A' on $V \otimes V^{*\otimes 2}$ and $V' \otimes V'^{*\otimes 2}$, then $x \otimes x'$ encodes the algebra $A \otimes A'$ on $(V \otimes V') \otimes (V \otimes V')^{*\otimes 2}$. Let x, x' encode $\mathcal{M}_n(K)$ and $\mathcal{M}_{n'}(K)$, so that $x \otimes x'$ encodes $\mathcal{M}_{nn'}(K)$. If $f : x \rightarrow a$ and $f' : x' \rightarrow a'$ are L -linear maps, then we have the L -linear map on $\mathcal{M}_{nn'}(L)$,

$$f \otimes f' : x \otimes x' \rightarrow a \otimes a'.$$

Now $\theta_n, \theta_{n'}$ map A and A' to $c_\sigma = f^{-1} \circ f^\sigma$ and $c'_\sigma = f'^{-1} \circ f'^\sigma$. Suppose c_σ and c'_σ are represented by conjugation by S_σ and S'_σ , respectively. Now $\theta_{nn'}$ maps $A \otimes A'$ onto $d_\sigma = (f \otimes f')^{-1} \circ (f \otimes f')^\sigma$, which corresponds to conjugation by $S_\sigma \otimes S'_\sigma$. Then by the description of Δ in Theorem 12.15.2, we see that

$$\delta_{nn'}(A \otimes A') = \{a_{\sigma,\tau} = i_{nn'}^{-1}[(S_\sigma \otimes S'_\sigma)\sigma(S_\tau \otimes S'_\tau)(S_{\sigma\tau} \otimes S'_{\sigma\tau})^{-1}]\} = \delta_n(A)\delta_{n'}(A')$$

where $i_{nn'}$ is the inclusion map $L^\times \rightarrow \mathrm{GL}_{nn'}(L)$. Under the inverse of $i_{nn'} = i_n \otimes i_{n'}$, tensor product becomes simply the product.

2. δ is surjective. It suffices to show Δ_n is surjective, where $n = [L : K]$.⁵ Take an 2-cocycle $a_{\sigma,\tau} \in H^2(G, L^\times)$. We need to show that

$$a_{\sigma,\tau} = S_\sigma \sigma(S_\tau) S_{\sigma\tau}^{-1}$$

for some values of $S_\sigma \in \mathrm{GL}_n(L)$. We identify L^n with the group algebra $L[G]$, and let $S_\sigma \in \mathrm{GL}(L[G])$ be the map sending τ to $a_{\sigma,\tau}\sigma\tau$ (it is invertible as $a_{\sigma,\tau} \in L^\times$). Then we calculate for every $u \in G \subset L[G]$,

$$\begin{aligned} [S_\sigma \sigma(S_\tau)]u &= [a_{\sigma,\tau u} \sigma(a_{\tau,u})] \sigma\tau u \\ [a_{\sigma\tau} S_{\sigma\tau}]u &= [a_{\sigma,\tau} a_{\sigma\tau,u}] \sigma\tau u. \end{aligned}$$

The right-hand sides are equal since $a_{\sigma,\tau}$ is a cocycle. Hence

$$a_{\sigma,\tau} = S_\sigma \sigma(S_\tau) S_{\sigma\tau}^{-1}$$

is in the image of Δ_n .

⁵Incidentally, this shows that every equivalence class of algebras is represented by one of dimension at most $[L : K]^2$. This is consistent with results of the previous section.

3. δ gives an isomorphism $\text{Br}_K \cong H^2(K)$: This follows from Theorem 13.4.14, the following easy-to-check commutative diagram (which holds for any $K \subseteq L \subseteq M$),

$$\begin{array}{ccc} H^2(L/K) & \xrightarrow{\text{Inf}} & H^2(M/K) \\ \delta \downarrow & & \delta \downarrow \\ \text{Br}_{L/K} & \xrightarrow{\quad} & \text{Br}_{M/K}, \end{array}$$

and taking the direct limit of the maps $\text{Br}_{L/K} \rightarrow H^2(L/K)$.

□

Remark 13.5.3: Milne [Mil08] makes this correspondence more explicit. The relationship between the two approaches can be seen by choosing a basis for the tensor product $V \otimes V^{*\otimes 2}$; the coefficients are called the *structure constants* of the algebra. (We followed Serre; note that the isomorphism in Serre is the opposite of the isomorphism in Milne.)

5.2 Exact sequence of Brauer groups

The importance of the Brauer group in class field theory is given by the following proposition.

Theorem 13.5.4. *Let $M/L/K$ be Galois extensions. Then there is an exact sequence*

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^2(L/K) & \longrightarrow & H^2(M/K) & \longrightarrow & H^2(M/L) \\ & & \parallel & & \parallel & & \parallel \\ & & \text{Br}_{L/K} & & \text{Br}_{M/K} & & \text{Br}_{M/L}. \end{array}$$

For any Galois extension L/K there is an exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^2(L/K) & \longrightarrow & H^2(K) & \longrightarrow & H^2(L) \\ & & \parallel & & \parallel & & \parallel \\ & & \text{Br}_{L/K} & & \text{Br}_K & & \text{Br}_L. \end{array}$$

Proof. Since $H^1(L/K) = 0$ by Hilbert's Theorem 90 (13.1.1), the inflation-restriction exact sequence 12.11.10 with $G = G(M/K)$ and $H = G(M/L)$ gives

$$0 \rightarrow H^2(L/K) \xrightarrow{\text{Inf}} H^2(M/K) \xrightarrow{\text{Res}} H^2(M/L).$$

The equality with the Brauer groups follows from Theorem 13.5.2.

Taking the direct limit over all finite Galois extensions M/K gives the second result. □

6 Problems

- 2.1 (Artin-Schreier) Let L/K be a Galois extension of degree p , with K/\mathbb{F}_p a finite extension. Prove that $L = K(\alpha)$ for some α such that $\alpha^p - \alpha \in K$. (Hint: Consider a short exact sequence as in the proof of Kummer theory. However, use the map $x \mapsto x^p - x$ instead of $x \mapsto x^p$, and consider additive instead of multiplicative groups.)

Chapter 14

Local class field theory

We now prove the main theorems of class field theory using cohomology. Throughout this chapter, K , L , etc. will denote nonarchimedean local fields, unless specified otherwise.¹ The main steps are the following.

1. Construct the invariant map $H^2(K^{\text{ur}}/K) \rightarrow \mathbb{Q}/\mathbb{Z}$. (Proposition 14.2.1)
 - (a) Show that $H^2(G(K^{\text{ur}}/K), U_{K^{\text{ur}}}) = 0$. (Theorem 14.1.1)
 - (b) From the decomposition $K^{\text{ur}\times} = U_{K^{\text{ur}}} \times \mathbb{Z}$ and step 1, we get $H^2(G(K^{\text{ur}}/K), K^{\text{ur}\times}) \cong H^2(G(K^{\text{ur}}/K), \mathbb{Z})$. (Note the projection $K^{\text{ur}\times} \rightarrow \mathbb{Z}$ is the valuation map $v_{K^{\text{ur}}}$.) Relate $H^2(G(K^{\text{ur}}/K), \mathbb{Z})$ to \mathbb{Q}/\mathbb{Z} using the long exact sequence in cohomology associated to $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow 0$.
2. Now show that there is an isomorphism $\text{Br}_K := H^2(\overline{K}/K) \cong H^2(K^{\text{ur}}/K)$ (Theorem 14.3.1). Thus we can restrict attention to unramified extensions of K and use step 1. Unramified extensions are easier to deal with! There are two approaches:

- (a) By Theorem 13.5.4 there is an exact sequence

$$0 \rightarrow H^2(K^{\text{ur}}/K) \rightarrow \text{Br}_K \rightarrow \text{Br}_{K^{\text{ur}}}.$$

Show that $\text{Br}_{K^{\text{ur}}} = 0$ by considering central simple algebras over local fields.

- (b) Study the cohomology of U_L when L/K is cyclic to conclude that the Herbrand quotient $h(U_L)$ is 1. From this get $h(L^\times) = [L : K]$. From this calculation and Hilbert's Theorem 90 (13.1.1), compute²

$$\begin{aligned} |H^1(L/K)| &= 1, \\ |H^2(L/K)| &= [L : K]. \end{aligned}$$

Conclude that $H^2(L/K)$ is cyclic of order $[L : K]$ and hence included in $H^2(K^{\text{ur}}/K)$, for any finite L/K .

¹Local class field theory for \mathbb{R} and \mathbb{C} is trivial and left to the reader. (The only nontrivial field extension is \mathbb{C}/\mathbb{R} .)

²This is the input for *abstract class field theory* according to Neukirch [Neu99].

3. Combining the first two steps, we get the invariant map $\text{inv}_K : \text{Br}_K \rightarrow \mathbb{Q}/\mathbb{Z}$. Show that this is compatible with restriction and hence that $(G(\overline{K}/K), \overline{K})$ is a *class formation*. Note inv_K restricts to $H^2(L/K) \rightarrow \frac{1}{[L:K]}\mathbb{Z}$; supposing its image is generated by $u_{L/K}$, Tate's Theorem 12.13.1 gives an isomorphism

$$\begin{array}{ccc} H_T^{-2}(G(L/K), \mathbb{Z}) & \xrightarrow[\cong]{\bullet \cup u_{L/K}} & H_T^0(G, L^\times) \\ \parallel & & \parallel \\ G(L/K)^{\text{ab}} & & K^\times / \text{Nm}_{L/K}(L^\times) \end{array}$$

that sends $\text{Frob}_{L/K}$ to $[\pi]$ when L/K is unramified. Taking a direct limit, we get a map $K^\times \rightarrow G(K^{\text{ab}}/K)$. Note we only get a map from G^{ab} (norm limitation).

4. Study the Hilbert symbol to prove the existence theorem (See Sections 6–7).

Unfortunately it is quite difficult to trace through the maps to find out what the Artin map actually is—for this Lubin-Tate Theory is better.

1 Cohomology of the units

For an unramified extension, the cohomology of the units is trivial.

Theorem 14.1.1 (Cohomology of units). *Suppose L/K is a finite unramified extension of local fields with Galois group G . Let U_L be the group of units of L . Then*

$$H_T^r(G, U_L) = 1$$

for any r . Hence $H^n(G(K^{ur}/K), U_{K^{ur}}) = 0$ for $n > 0$.

Proof. We will show that

$$H_T^1(G, U_L) = H_T^0(G, U_L) = 1.$$

Then it follows from Proposition 12.12.1 that all the Tate groups are trivial. The second part follows from taking the direct limit.

We have

$$L^\times = U_L \times \pi^\mathbb{Z} \cong U_L \times \mathbb{Z} \tag{14.1}$$

where π is a uniformizer for L . Since L/K is unramified, we can choose $\pi \in K$. Then G acts trivially on π , so acts trivially on \mathbb{Z} in the decomposition above. Thus (14.1) gives a decomposition of L^\times as a G -module (not just as a group). We have by Hilbert's Theorem 90 (Theorem 13.1.1) and the fact that cohomology respects products (Proposition 12.6.7) that

$$0 = H^1(G, L^\times) = H^1(G, U_L) \times H^1(G, \mathbb{Z}).$$

Hence $H^1(G, U_L) = 1$.

It remains to show $H_T^0(G, U_L) = 1$. To do this, let \mathfrak{m} be the maximal ideal of L , $U_L^{(m)} := 1 + \mathfrak{m}^m$, and consider the filtration

$$U_K^{(0)} := U_K \supset U_K^{(1)} \supset U_K^{(2)} \supset \dots$$

Proposition 14.1.2 and 14.1.3 below show that each quotient has trivial cohomology:

$$H_T^0(G, U_L^{(i)}/U_L^{(i+1)}) = 1.$$

Then Lemma 14.1.4 gives that $H_T^0(G, U_L) = 1$, as needed. \square

Proposition 14.1.2: Let K be a complete field with discrete valuation, \mathfrak{m} be the associated maximal ideal, and $U_K^{(m)} := 1 + \mathfrak{m}^m$. Then we have isomorphisms

$$\begin{array}{ccc} U_K/U_K^{(1)} \xrightarrow{\cong} k^\times & & U_K^{(m)}/U_K^{(m+1)} \xrightarrow{\cong} k^+ \\ u \mapsto u \pmod{\mathfrak{m}} & & 1 + a\pi^m \mapsto a \pmod{\mathfrak{m}} \end{array}$$

that preserve Galois action.

Proof. This is Proposition 9.4.8. \square

Proposition 14.1.3: Let l/k be an extension of finite fields and $G := G(l/k)$. Then

$$\begin{aligned} H_T^r(G, l^\times) &= \{1\} \\ H_T^r(G, l^+) &= \{0\} \end{aligned}$$

for all $r \in \mathbb{Z}$. Moreover, the maps $\text{Nm}_{l/k} : l \rightarrow k$ and $\text{tr}_{l/k} : l \rightarrow k$ are surjective.

Proof. By Hilbert's Theorem 90 (13.1.1), $H^1(G, l^\times) = 0$. Since G is cyclic and l is finite, by Proposition 12.12.4, $h(l^\times) = 1$, giving $H^2(G, l^\times) = 0$. Again since G is cyclic, by Theorem 12.12.1, all the Tate groups are 0.

From Theorem 13.1.2, $H_T^r(G, l^+) = 0$ for $r \geq 0$.

For the second statement, just note

$$\begin{aligned} \{1\} &= H_T^0(G, l^\times) = (l^\times)^G / N_G(l^\times) = k^\times / \text{Nm}_{l/k}(l^\times) \\ \{0\} &= H_T^0(G, l^+) = l^G / N_G(l) = k / \text{tr}_{l/k}(l). \end{aligned} \quad \square$$

Lemma 14.1.4. Let G be a finite group and M be a G -module. Let

$$M = M^0 \supseteq M^1 \supseteq \dots$$

be a decreasing sequence of G -submodules and suppose $M = \varprojlim M/M^i$ (i.e. M is complete with respect to this filtration). If $H^q(G, M^i/M^{i+1}) = 0$ for all i , then $H^q(G, M) = 0$.

Proof. Let f be a q -cocycle of M . Since $H^q(G, M/M^1) = 0$, the long exact sequence of $0 \rightarrow M^1 \rightarrow M \rightarrow M/M^1$ gives $H^q(G, M^1) \twoheadrightarrow H^q(G, M)$ and we can write $f = g_0 + f_1$, where $g_0 = \delta h_0$ is a coboundary in M and f_1 is a q -cocycle in M^1 . Given $f_n \in H^q(G, M^n)$, we can write

$$f_n = \delta h_n + f_{n+1}$$

where h_n is a $(q-1)$ -cocycle of M^n and f_{n+1} is a q -cocycle of M^{n+1} . Then

$$f = \delta(h_1 + h_2 + \cdots),$$

the infinite series being defined in $H^{q-1}(G, M)$ since h_n is a cochain with values in M^n , and M is complete with respect to this filtration. \square

This proves Theorem 14.1.1. We record the following corollary, for easy reference.

Corollary 14.1.5. *Suppose L/K is a finite extension of local fields. Then*

$$U_K \subseteq \text{Nm}_{L/K} U_L.$$

Proof. If L/K is Galois, then this follows since by Theorem 14.1.1

$$U_K / \text{Nm}_{L/K} U_L = H_T^0(G(L/K), U_L) = \{1\}$$

so the norm map $U_L \rightarrow U_K$ is surjective.

For general extensions L/K , consider the Galois closure and use transitivity of norms. \square

2 The invariant map

2.1 Defining the invariant maps

Proposition 14.2.1: For any finite unramified Galois extension of local fields L/K there is a canonical isomorphism

$$\text{inv}_{L/K} : H^2(L/K) \xrightarrow{\cong} \frac{1}{[L : K]} \mathbb{Z}/\mathbb{Z}.$$

Taking the direct limit gives an injective map

$$\text{inv}_{K^{\text{ur}}/K} : H^2(K^{\text{ur}}/K) \rightarrow \mathbb{Q}/\mathbb{Z}.$$

Proof. Consider the short exact sequence

$$1 \rightarrow U_L \rightarrow L^\times \xrightarrow{v_L} \mathbb{Z} \rightarrow 0.$$

Since $H_T^n(G, U_L) = 0$ for all n by Theorem 14.1.1, taking the long exact sequence gives

$$\cancel{H^2(G, U_L)} \xrightarrow{0} H^2(L/K) \xrightarrow{\cong} H^2(G, \mathbb{Z}) \rightarrow \cancel{H^3(G, U_L)} \xrightarrow{0}$$

We relate $H^2(G, \mathbb{Z})$ to a lower cohomology group by considering the short exact sequence

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow 0.$$

Note $H^n(G, \mathbb{Q})$ is torsion for any $n > 0$ by Corollary 12.11.6. Since \mathbb{Q} is a divisible group, so is $H^n(G, \mathbb{Q})$, by looking at the description of H^n in terms of cocycles (Section 7). Hence $H^n(G, \mathbb{Q}) = 0$ for any $n > 0$. Taking the long exact sequence of the above we get

$$\cancel{H^1(G, \mathbb{Q})} \xrightarrow{0} H^1(G, \mathbb{Q}/\mathbb{Z}) \xrightarrow{\cong} H^2(G, \mathbb{Z}) \rightarrow \cancel{H^2(G, \mathbb{Q})} \xrightarrow{0}$$

Thus we get a map

$$\text{inv}_{L/K} : H^2(L/K) \xrightarrow{\cong} H^2(G, \mathbb{Z}) \xleftarrow{\cong} H^1(G, \mathbb{Q}/\mathbb{Z}) \stackrel{12.7.3}{\cong} \text{Hom}(G, \mathbb{Q}/\mathbb{Z}) \xrightarrow{\cong} \frac{1}{[L:K]} \mathbb{Z}/\mathbb{Z}. \quad (14.2)$$

where the last is defined by taking the Frobenius element σ of G and mapping $f \mapsto f(\sigma)$. (Note G is cyclic and σ generates G ; the Frobenius is a canonical choice.)

Now define $\text{inv}_{K^{\text{ur}}/K} = \varinjlim_{L/K \text{ finite Galois unramified}} \text{inv}_{L/K}$, taking the direct limit under inflation. Since inflation is functorial, the first two maps in (14.2) commute with it. Identifying $H^1(G, \mathbb{Q}/\mathbb{Z}) \cong \text{Hom}(G, \mathbb{Q}/\mathbb{Z})$, inflation sends a map $G(L/K) \rightarrow \mathbb{Q}/\mathbb{Z}$ to $G(M/K) \rightarrow G(L/K) \rightarrow \mathbb{Q}/\mathbb{Z}$. Moreover, $\text{Frob}_{L/K}$ is the projection of $\text{Frob}_{M/K}$ to $G(L/K)$. Hence $\text{Inf}_{M/L}$ commutes with the inclusion map $\frac{1}{[L:K]} \mathbb{Z}/\mathbb{Z} \hookrightarrow \frac{1}{[M:K]} \mathbb{Z}/\mathbb{Z}$, and the $\text{inv}_{L/K}$ form a compatible system under inflation. \square

Remark 14.2.2: Let K be any nonarchimedean complete field (not necessarily local) with residue field k . Then

$$H^n(L/K) = H^n(l/k) \times H^n(G(L/K), \mathbb{Q}/\mathbb{Z}).$$

Indeed, Proposition 14.1.2 and Theorem 13.1.2 still give

$$H_T^r(G, U_L^{(i)}/U_L^{(i+1)}) \cong H_T^r(G, l^+) = 0$$

for $i \geq 1$. This gives $H_T^r(G, U_L^{(1)}) = 0$ by Lemma 14.1.4. From the long exact sequence associated to

$$1 \rightarrow U_L^{(1)} \rightarrow U_L \rightarrow U_L/U_L^{(1)} \cong l^\times \rightarrow 1$$

we get

$$H^n(L/K) \cong H^n(G, U_L) \times H^n(G, \mathbb{Z}) = H^n(G, l^\times) \times H^n(G, \mathbb{Z}).$$

In the case of a local field, l was finite so $H^n(G, l^\times) = 1$.

2.2 Compatibility of the invariant maps

We show that the invariant maps are compatible, in the following sense.

Theorem 14.2.3. *Let L/K be a Galois extension of local fields, and $n = [L : K]$. Then*

$$\text{inv}_{K^{\text{ur}}/L} \circ \text{Res}_{K/L} = n \text{inv}_{K^{\text{ur}}/K}$$

Proof. To do this we have to unravel all those steps we took to define $\text{inv}_{K^{\text{ur}}/K} \dots$. We first prove this for two special cases.

1. L/K is unramified. Let $G = G(K^{\text{ur}}/K)$ and $S = G(K^{\text{ur}}/L)$. We claim the following commutes.

$$\begin{array}{ccccccc} H^2(K^{\text{ur}}/K) & \longrightarrow & H^2(G, \mathbb{Z}) & \longleftarrow & H^1(G, \mathbb{Q}/\mathbb{Z}) & \xrightarrow{\gamma} & \mathbb{Q}/\mathbb{Z} \\ \downarrow \text{Res} & & \downarrow \text{Res} & & \downarrow \text{Res} & & \downarrow n \\ H^2(K^{\text{ur}}/L) & \longrightarrow & H^2(S, \mathbb{Z}) & \longleftarrow & H^1(S, \mathbb{Q}/\mathbb{Z}) & \xrightarrow{\gamma} & \mathbb{Q}/\mathbb{Z}. \end{array}$$

For the squares involving Res, this follows from naturality of Res. For the last square, identify $H^1(G, \mathbb{Q}/\mathbb{Z}) = \text{Hom}(G, \mathbb{Q}/\mathbb{Z})$; Res becomes simply restriction of homomorphisms. Recall that γ was defined taking the Frobenius $\text{Frob}(K^{\text{ur}}/K) \in G(K^{\text{ur}}/K)$ and sending $f \in H^1(G, \mathbb{Q}/\mathbb{Z}) = \text{Hom}(G, \mathbb{Q}/\mathbb{Z})$ to $f(\sigma)$, and we have

$$\text{Frob}_{K^{\text{ur}}/K}^n = \text{Frob}_{K^{\text{ur}}/L}$$

by Proposition 11.1.4.

2. L/K is totally ramified. Note that $G = G(K^{\text{ur}}/K) = G(K^{\text{ur}}L/L) = G(L^{\text{ur}}/L)$ in this case, from the description of K^{ur} in Theorem 8.2.6. We show the following commutes:

$$\begin{array}{ccccccc} H^2(K^{\text{ur}}/K) & \longrightarrow & H^2(G, \mathbb{Z}) & \longleftarrow & H^1(G, \mathbb{Q}/\mathbb{Z}) & \xrightarrow{\gamma} & \mathbb{Q}/\mathbb{Z} \\ \downarrow \text{Res} & & \downarrow n & & \downarrow n & & \downarrow n \\ H^2(K^{\text{ur}}/L) & \longrightarrow & H^2(G, \mathbb{Z}) & \longleftarrow & H^1(G, \mathbb{Q}/\mathbb{Z}) & \xrightarrow{\gamma} & \mathbb{Q}/\mathbb{Z}. \end{array}$$

The first square commutes by commutativity of

$$\begin{array}{ccc} K^{\text{ur}\times} & \xrightarrow{v_K} & \mathbb{Z} \\ \downarrow & & \downarrow n \\ L^{\text{ur}\times} & \xrightarrow{v_L} & \mathbb{Z}. \end{array}$$

(and of course, naturality of cohomology). Here v_K and v_L are the valuation maps, i.e. the projections $K^\times \cong U_K \times \mathbb{Z} \rightarrow \mathbb{Z}$ and $L^\times \cong U_L \times \mathbb{Z} \rightarrow \mathbb{Z}$.

The general case follows by considering $L/L^{L/K}$ (totally ramified) and $L^{L/K}/K$ (unramified). (See Theorem 2.7.2.) \square

3 $H^2(\overline{K}/K) \cong H^2(K^{\text{ur}}/K)$

We prove the following.

Theorem 14.3.1. *The inclusion (inflation) map*

$$H^2(K^{\text{ur}}/K) \rightarrow H^2(\overline{K}/K)$$

is an isomorphism.

For short we write $H^2(K) := H^2(\overline{K}/K)$.

3.1 First proof (Brauer group)

First proof. By Proposition 13.5.4 there is an exact sequence

$$0 \rightarrow H^2(K^{\text{ur}}/K) \rightarrow H^2(K) \rightarrow H^2(K^{\text{ur}}) = \text{Br}_{K^{\text{ur}}} \rightarrow 0$$

The last term is zero by Theorem 14.3.2 below. Thus we get $H^2(K^{\text{ur}}/K) \cong H^2(K)$, as needed. \square

Theorem 14.3.2. *Let K be a local field. Then $\text{Br}_{K^{\text{ur}}} = 0$.*

We need two lemmas.

Lemma 14.3.3. *Suppose D is a central division algebra of rank $n^2 > 1$ over a field K , and the residue field k is perfect. Then there exists a commutative subfield L of D properly containing K , unramified over K .*

Lemma 14.3.4. *Keep the same hypotheses as Lemma 14.3.3. There is a subfield of D of degree n unramified over K .*

Note this is a maximal subfield by Corollary 13.4.13.

Proof of Lemma 14.3.3. Suppose by way of contradiction that every commutative subfield L of D properly containing K is ramified. Then the extension of residue fields l/k must be trivial (see Theorem 2.7.2). Let $a \in D$ be integral and $\pi \in D$ be a uniformizer for D . (See Proposition 13.4.15.) Since $l = k$, there exists $b \in K$ such that $b \equiv a \pmod{\pi}$, and we can write $a = b + \pi b_1$ for some $b_1 \in \mathcal{O}_D$, where \mathcal{O}_D is the ring of integers in D . Iterating this with b_1 , we find

$$a = b + \pi b_1 + \cdots + \pi^{n-1} b_{n-1} + \pi^n b_n$$

where $b_1, \dots, b_{n-1} \in \mathcal{O}_K$ and $b_n \in \mathcal{O}_D$. Thus a is in the closure of $K(\pi)$. But $K(\pi)$ is closed (it is a finite-dimensional vector space over K), so $a \in K(\pi)$, i.e. $D = K(\pi)$ and D is commutative, a contradiction. \square

Proof of Lemma 14.3.4. Induct on n . The case $n = 1$ is clear. Let $n \geq 2$. By Lemma 14.3.3 there exists a proper unramified extension K'/K inside D . Let $D' = C(K')$. Since $D' \subseteq D$, D' must be a division algebra (a finite dimensional integral domain must contain inverses). Let its center be K'' . The maximal commutative subfield of D' then has dimension $\sqrt{[D' : K'']}$ over K'' , or dimension $\sqrt{[D' : K'']}[K'' : K] = \sqrt{[D' : K][K'' : K]}$ over K . This is at most n , since the field is also contained in D . But $\sqrt{[D' : K][K' : K]} = n$ by the double centralizer theorem 13.4.11, so we must have $K'' = K$. Thus D' is a division algebra with center K' . Its degree over K' is less than n^2 , so by the induction hypothesis, D' has a maximal commutative subfield L containing K' , of degree $\sqrt{[D' : K']}$, and unramified over K' , hence over K . We calculate

$$[L : K] = [L : K'][K' : K] = \sqrt{[D' : K']}[K' : K] = \sqrt{[D' : K][K' : K]} = \sqrt{[D : K]}$$

where we used Theorem 13.4.11 in the last step. This finishes the induction step. \square

Proof of Theorem 14.3.2. Suppose D is a central division algebra over K^{ur} of rank n^2 . Then lemma 2 furnishes a subfield of K^{ur} of degree n , unramified over K^{ur} . Hence $n = 1$, and D is trivial. Thus $\text{Br}_{K^{\text{ur}}} = 0$. This proves Theorem 14.3.2 and hence Theorem 14.3.1. \square

3.2 Second proof (Herbrand quotient calculation)

3.2.1 Herbrand quotient calculation

We first need the following lemma.

Lemma 14.3.5. *Given a local field L , there exists an open subgroup V of U_L with trivial cohomology, i.e. $H^q(G, V) = 0$ for all q .*

Proof. The idea is to compare a multiplicative G -module V with an additive G -module (more accurately, compare the filtration of V), and use the same argument as in Theorem 13.1.2.³

By the normal basis theorem, L^+ has a normal basis $\{\sigma(\alpha) : \sigma \in G\}$, i.e. it is free over $K[G]$. Let $A = \sum_{\sigma \in G} \mathcal{O}_K \sigma(\alpha)$.⁴ By multiplying α by a power of π_K we may assume that $\alpha \in \mathcal{O}_L$. Suppose that

$$\pi_K^n \mathcal{O}_L \subseteq A \subseteq \mathcal{O}_L.$$

Let $M = \pi_K^{n+1} A$, $V = 1 + M$ and $V^{(i)} = 1 + \pi_K^i M$. Note that

$$M \cdot M \subseteq \pi_K^{2n+2} A \cdot A \subseteq \pi_K \pi_K^{n+1} \pi_K^n \mathcal{O}_L \subseteq \pi_K \pi_K^{n+1} A \subseteq \pi_K M.$$

³If $\text{char}(L) = 0$ there is a faster proof: Note that e^x is a topological isomorphism from a neighborhood of 0 in the additive group L to a neighborhood of 1 in the multiplicative group \mathcal{O}_L . Moreover, it preserves the action of G because the fact that G acts continuously on L gives

$$e^{\sigma x} = \sum_{n=0}^{\infty} \frac{(\sigma x)^n}{n!} = \sum_{n=0}^{\infty} \frac{\sigma(x^n)}{n!} = \sigma e^x.$$

Now Theorem 13.1.2 applies directly.

⁴Warning: A is a $\mathcal{O}_K[G]$ -module; we don't know it is an \mathcal{O}_L -module.

This shows that

1. V is a subgroup: Indeed, $(1 + M)(1 + M) \subseteq 1 + M + M \cdot M \subseteq 1 + M$ by the above.
2. $V^i/V^{i+1} \cong A/\pi_K A$ as G -modules. Indeed, if $m_1, m_2 \in M$, then for some $m_3 \in M$, we have

$$(1 + \pi_K^i m_1)(1 + \pi_K^i m_2) = 1 + \pi_K^i(m_1 + m_2) + \pi_K^{2i} \pi_K m_3 \equiv 1 + \pi_K^i(m_1 + m_2) \pmod{\pi_K^{i+1} M}.$$

Hence

$$H^q(G, V^{(i)}/V^{(i+1)}) = H^q(G, M/\pi_K M) = 0$$

for each q , since $M/\pi_K M$ is an induced module over G (and has trivial cohomology by Shapiro's Lemma 8.1). (By construction $M/\pi_K M = \text{Ind}^G[(\pi_K^{n+1} \alpha \mathcal{O}_K)/(\pi_K^{n+2} \alpha \mathcal{O}_K)]$.) Lemma 14.1.4 applied to V finishes the proof. \square

Proposition 14.3.6: Suppose L/K is cyclic of degree n . Then

$$\begin{aligned} h(U_L) &= 1. \\ h(L^\times) &= n. \end{aligned}$$

Proof. Choose V as in Lemma 14.3.5. Since V is open, U_L/V is finite. By Proposition 12.12.4(1), $h(U_L/V) = 1$. Hence

$$h(U_L) = h(V)h(U_L/V) = 1.$$

By Proposition 12.12.4(3), $h(\mathbb{Z}) = |G| = n$. Since $L^\times = U_L \times \pi_L^{\mathbb{Z}}$ we get

$$h(L^\times) = h(U_L)h(\mathbb{Z}) = n.$$

\square

Theorem 14.3.7 (Class field axiom for local class field theory). *Let L/K be a cyclic extension of degree n . Then*

$$\begin{aligned} |H^1(L/K)| &= 1 \\ |H^2(L/K)| &= n. \end{aligned}$$

Proof. The first follows directly from Hilbert's Theorem 90 (13.1.1). For the second, we have $|H^2(L/K)| = h(L^\times)|H^1(L/K)| = n$ using Proposition 14.3.6. \square

We want to show that $|H^2(L/K)| = n$ for all Galois extensions L/K , and in fact $H^2(L/K)$ is cyclic of order n . We proceed in 2 steps.

3.2.2 First inequality

We show that for all Galois extensions L/K , $|H^2(L/K)| \geq [L : K]$. In fact, we show the following.

Lemma 14.3.8. *Let L/K be a Galois extension of local fields of degree n . Then $H^2(L/K)$ contains a subgroup canonically isomorphic to $\frac{1}{n}\mathbb{Z}/\mathbb{Z}$.*

Proof. We prove this using Theorem 14.2.3, which relates the invariant maps on K^{ur}/K and L^{ur}/L . By Theorem 13.5.4, we have the exact sequence $0 \rightarrow H^2(L/K) \rightarrow H^2(K) \rightarrow H^2(L)$. Inflation and restriction commute by functoriality of change of group, so we have the commutative diagram with exact columns

$$\begin{array}{ccc}
 0 & & 0 \\
 \downarrow & & \downarrow \\
 H^2(L/K) & \longleftarrow & \ker(\text{Res}) \\
 \downarrow & & \downarrow \\
 H^2(K) & \xleftarrow{\text{Inf}} & H^2(K^{\text{ur}}/K) \\
 \downarrow \text{Res} & & \downarrow \text{Res} \\
 H^2(L) & \xleftarrow{\text{Inf}} & H^2(K^{\text{ur}}/L).
 \end{array} \tag{14.3}$$

By Theorem 14.2.3, the map $H^2(K^{\text{ur}}/K) \rightarrow H^2(L^{\text{ur}}/L)$ corresponds to the multiplication-by- $[L : K]$ map after identifying both sides with a subgroup of \mathbb{Q}/\mathbb{Z} through the respective invariant maps. Hence $\ker(\text{Res}) = \frac{1}{n}\mathbb{Z}/\mathbb{Z}$. The top map exists and is an injection because the other two are (4-lemma). Hence $\frac{1}{n}\mathbb{Z}/\mathbb{Z} \hookrightarrow H^2(L/K)$, as needed. \square

3.2.3 Second inequality

Next we show $|H^2(L/K)| \leq [L : K]$, so $|H^2(L/K)| = [L : K]$.

Lemma 14.3.9. *Let L/K be a Galois extension of local fields of degree n . Then $H^2(L/K) \cong \frac{1}{n}\mathbb{Z}/\mathbb{Z}$.*

Proof. We already know that $|H^2(L/K)| = [L : K]$ for L/K cyclic (Theorem 14.3.7). We prove that $|H^2(L/K)| = [L : K]$ by induction on the degree.

By Corollary 9.4.12, $G(L/K)$ is solvable. Thus, if $G(L/K)$ is not cyclic, it has a normal subgroup $G(L/K')$. By Theorem 13.5.4 we have an exact sequence

$$0 \rightarrow H^2(K'/K) \rightarrow H^2(L/K) \rightarrow H^2(L/K')$$

so

$$|H^2(L/K)| \leq |H^2(K'/K)| \cdot |H^2(L/K')| = [K' : K][L : K'] = [L : K].$$

By Lemma 14.3.8, equality holds. \square

3.2.4 Finishing the proof

Second proof of Theorem 14.3.1. Take any element $a \in H^2(\overline{K}/K)$; it is in $H^2(L/K)$ for some finite Galois L/K . The top injection in (14.3) is an isomorphism by Lemma 14.3.9, and we get $a \in H^2(K^{\text{ur}}/K)$. \square

4 Class formations

The preceding sections show that

$$(G(\overline{K}/K), \{G(L/K) : L/K \text{ finite Galois}\}, \overline{K})$$

is a *class formation*. That is, it satisfies the basic axioms that allow us to obtain the conclusions of class field theory. With the abstraction of class formations, when we prove global class field theory, we only have to verify the axioms and we will get the desired conclusions in the same way as in local class field theory.

4.1 Class formations in the abstract

Definition 14.4.1: An **abstract Galois group** is a group G with a family of subgroups of finite index $\{G_L\}_{L \in X}$ such that

1. (Closure under intersection) If $L_1, L_2 \in X$, then there exists M such that

$$G_{L_1} \cap G_{L_2} = G_M.$$

2. (Closure under superset) If $G_L \subseteq G' \subseteq G$ are subgroups, then $G' = G_{K'}$ for some K' .
3. (Closure under conjugation) For every $s \in G$ and $L \in X$ there exists L' so that

$$sG_Ls^{-1} = G_{L'}.$$

This definition is motivated by the fact that these are the key properties of Galois groups.

Proposition 14.4.2: A topological Galois group $G(\Omega/K_0)$ with all its closed subgroups, is an abstract Galois group.

Proof. By the fundamental theorem of infinite Galois theory 5.8.4, the closed subgroups of $G(\Omega/K_0)$ are exactly those in the form $G(\Omega/K)$ with $K_0 \subseteq K \subseteq \Omega$. The above properties correspond to the following facts from Galois theory.

1. $G(\Omega/K) \cap G(\Omega/L) = G(\Omega/KL)$.
2. The subgroups of $G(\Omega/K_0)$ containing $G(\Omega/L)$ correspond to intermediate extensions between K_0 and L .

$$3. sG(\Omega/K)s^{-1} = G(\Omega/sK). \quad \square$$

We transfer some terminology about Galois groups to the abstract case.

Definition 14.4.3: Let $(G, \{G_L\}_{L \in X})$ be an abstract Galois group. The elements of X are called fields. The field K_0 with $G_{K_0} = G$ is called the basefield. For $G_M \subseteq G_L$, define $[M : L]$ to be $[G_L : G_M]$; we say M/L is a Galois extension if $G_M \trianglelefteq G_L$, and write

$$G(M/L) = G_L/G_M$$

(called the ‘‘Galois group’’ of M/L). We say M/L is abelian, etc. if $G(M/L)$ is abelian, etc.

The field M such that $G_{L_1} \cap G_{L_2} = G_M$ is called the composite of L_1 and L_2 , and denoted by L_1L_2 ; the field L' such that $sG_Ls^{-1} = G_{L'}$ is denoted by sL .

Note every extension M/L is contained in a Galois extension: Since $[G_L : G_M]$ is finite G_M has finitely many conjugates sG_Ms^{-1} in G_L ; by the axioms $G_{M'} = \bigcap_s sG_Ms^{-1}$ for some M' , called the Galois closure of M/L .

Definition 14.4.4: A **formation** is a triple $(G, \{G_K\}_{K \in X}, A)$ where $(G, \{G_K\}_{K \in X})$ is an abstract Galois group and A is a discrete topological G -module (see Definition 12.14.1). Let $A_K := A^{G_K}$.

Define the norm $\text{Nm}_{L/K} : A_L \rightarrow A_K$ by letting $\text{Nm}_{L/K}(a) = \prod_{\sigma \in G(L'/K)/G(L/K)} \sigma(a)$ for any L' Galois over K .

For L/K Galois, we define $H^n(L/K) := H^n(G(L/K), A_L)$. We can define inflation, restriction, and corestriction maps in the natural way, with $\text{Res}_{K/L} = \text{Res}_{G_K/G_L}$, and so forth.

Definition 14.4.5: A **class formation** is a formation $(G, \{G_K\}_{K \in X}, A)$ with a homomorphism $\text{inv}_{L/K} : H^2(L/K) \rightarrow \mathbb{Q}/\mathbb{Z}$ for each Galois extension L/K , such that the following hold.

1. $H^1(L/K) = 0$ for every cyclic extension of prime degree.
2. $\text{inv}_{L/K}$ is an isomorphism from $H^2(L/K)$ to $\frac{1}{[L:K]}\mathbb{Z}/\mathbb{Z}$.
3. (Compatibility under inflation) For any finite extension M/L ,

$$\text{inv}_{M/K} \circ \text{Inf}_{M/L} = \text{inv}_{L/K}.$$

Hence we can define $\text{inv}_K : \varinjlim_L H^2(L/K) \rightarrow \mathbb{Q}/\mathbb{Z}$. (This axiom implies that inflations are injective on H^2 , so we can think of $H(K) := \varinjlim_L H^2(L/K)$ as $\bigcup_L H^2(L/K)$.)

4. (Compatibility with restriction) For any finite Galois extension L/K ,

$$\text{inv}_L \circ \text{Res}_{K/L} = [L : K] \text{inv}_K.$$

Define the **fundamental unit** of L/K to be

$$u_{L/K} = \text{inv}_K^{-1} \left(\frac{1}{[L : K]} \right).$$

Proposition 14.4.6: Assume a formation satisfies axiom 1. Then for every Galois extension L/K ,

$$H^1(L/K) = 0.$$

Proof. First we show this when $[L : K]$ is a prime power p^n . Induct on the degree. The base case is given. Every p -group has a subgroup of index p , so there is $K \subset K' \subset L$ such that $G(K'/K)$ has order p . By the inflation-restriction exact sequence 12.11.10, we get

$$0 \rightarrow \cancel{H^1(K'/K)} \xrightarrow{\text{Inf}} H^1(K/L) \xrightarrow{\text{Res}} \cancel{H^1(L/K)} \rightarrow 0$$

the first and last terms are 0 by axiom 1 and by the induction hypothesis. So $H^1(K/L) = 0$.

For general L/K , this shows $H^1(G(L/K)_p, A_L) = 0$, so the result follows from Corollary 12.11.8. \square

Proposition 14.4.7: Assume a formation satisfies axiom 2. Transferring the action of Res, Cor, and Inf to the subgroups of \mathbb{Q}/\mathbb{Z} , we get the following diagram:

$$\begin{array}{c}
 M \\
 \downarrow [M:L] \\
 L \\
 \downarrow [L:K] \\
 K
 \end{array}
 \quad
 \begin{array}{ccc}
 H^2(M/L) & \xrightarrow{\text{inv}_L} & \frac{1}{[M:L]} \mathbb{Z}/\mathbb{Z} \\
 \uparrow \text{Cor}_{L/K} & \text{Res}_{K/L} & \uparrow i \\
 H^2(M/K) & \xrightarrow{\text{inv}_K} & \frac{1}{[M:K]} \mathbb{Z}/\mathbb{Z}
 \end{array}
 \quad
 \begin{array}{c}
 L \\
 \downarrow \\
 K
 \end{array}
 \quad
 \begin{array}{ccc}
 & & H^2(L/K) \xrightarrow{\text{inv}_K} \frac{1}{[L:K]} \mathbb{Z}/\mathbb{Z} \\
 & \swarrow \text{Inf}_{M/L} & \nwarrow i
 \end{array}$$

(Note $\text{Cor}_{L/K} \circ \text{Res}_{K/L} = [L : K]$.) Moreover (passing to the limit), the following hold.

1. For every extension L/K ,

$$\text{Res}_{K/L} : H^2(K) \rightarrow H^2(L)$$

is surjective.

2. For every extension L/K ,

$$\text{Cor}_{L/K} : H^2(L) \hookrightarrow H^2(K)$$

is injective, and

$$\text{inv}_K \circ \text{Cor}_{L/K} = \text{inv}_L.$$

3. For every $s \in G$, letting $s^* : H^2(K) \rightarrow H^2(sK)$,

$$\text{inv}_{sK} \circ s^* = \text{inv}_K.$$

Proof. The surjectivity of $\text{Res}_{K/L}$ in the diagram comes directly from the injectivity of inv_K and $\text{inv}_L \circ \text{Res}_{K/L} = [L : K] \text{inv}_K$.

For the action of $\text{Cor}_{L/K}$, note

$$\text{inv}_K \circ \text{Cor}_{L/K} \circ \text{Res}_{K/L} = \text{inv}_K \circ [L : K] = \text{inv}_L \circ \text{Res}_{K/L}$$

where the first follows from Theorem 12.11.5 and the second from the axiom. Surjectivity of $\text{Res}_{K/L}$ gives $\text{inv}_K \circ \text{Cor}_{L/K} = \text{inv}_L$, as needed.

Items 1 and 2 now follow from taking the direct limit.

For 3, let the basefield be K_0 ; note the map $s^* : H^2(K_0) \rightarrow H^2(sK_0) = H^2(K_0)$ is the identity by Proposition 12.11.3, so $\text{inv}_{sK_0} \circ s^* = \text{inv}_{K_0}$. For arbitrary $x \in H^2(K)$, by surjectivity of $\text{Res}_{K/L}$ we can write $x = \text{Res}_{K/L}(x_0)$. Since Res and s^* commute (transport of structure),

$$\text{inv}_{sK}(s^*x) = \text{inv}_{sK}(s^* \text{Res}_{K_0/K} x_0) = \text{inv}_{sK} \text{Res}_{sK/sK_0}(s^*x_0) = [sK : sK_0] \text{inv}_{sK_0}(x_0) = \text{inv}_K(x).$$

□

The reciprocity law follows from the properties of class formations.

Theorem 14.4.8 (Abstract reciprocity law). *Let $(G, \{G_K\}_{K \in X}, \{A_K\}, \text{inv}_{L/K})$ be a class formation. Then there is an isomorphism*

$$\begin{array}{ccc} H_T^{-2}(G(L/K), \mathbb{Z}) \xrightarrow[\cong]{u_{L/K} \cup \bullet} & H_T^0(G, A_L) & \\ \parallel & \parallel & \\ G(L/K)^{ab} & A_K / \text{Nm}_{L/K}(A_L) & \end{array}$$

Here $\text{Nm}_{L/K}$ means N_{G_K/G_L} . Denote the reverse map by $\phi_{L/K}$.

Proof. The identifications are from Theorem 12.8.3 and Definition 12.9.2. Axioms 1 and 2 for class formation give that the two conditions of Tate's Theorem 12.13.1 are satisfied. □

This map is hard to calculate directly because cup products on negative Tate cohomology are hard to deal with. The following helps us by transferring the cup products to nonnegative Tate groups.

Theorem 14.4.9. *Keep the above hypothesis. Then for any $\chi \in \text{Hom}^{\text{cont}}(G(L/K), \mathbb{Q}/\mathbb{Z}) = H^1(G, \mathbb{Q}/\mathbb{Z})$ and $a \in A_K$,*

$$\chi(\phi_{L/K}(a)) = \text{inv}_K(\bar{a} \cup \delta\chi).$$

Here \bar{a} denotes the image of a in $H_T^0(G(L/K), A_L) = A_L / \text{Nm}_{L/K} A_L$, and δ is the diagonal morphism corresponding to the exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow 0$.

Note this characterizes the reciprocity map since knowing the image of an element of an abelian group under all homomorphisms to \mathbb{Q}/\mathbb{Z} is equivalent to knowing the element itself.⁵

Proof. Suppose $\chi(\phi_{L/K}(a)) = \frac{r}{n}$.

By the definition of the Artin map as the inverse of $u_{L/K} \cup \bullet$, we have

$$\bar{a} = u_{L/K} \cup \phi_{L/K}(a).$$

We now calculate the following (for easy reference, we note which cohomology groups the elements are in).

$$\begin{aligned}
 \underbrace{\bar{a}}_0 \cup \underbrace{\delta\chi}_2 &= \underbrace{[u_{L/K} \cup \phi_{L/K}(a)]}_2 \cup \underbrace{\delta\chi}_2 \\
 &= \underbrace{u_{L/K}}_2 \cup \underbrace{[\phi_{L/K}(a) \cup \delta\chi]}_2 && \text{associativity} \\
 &= \underbrace{u_{L/K}}_2 \cup \underbrace{[\delta(\phi_{L/K}(a) \cup \chi)]}_1 && \text{Theorem 12.10.1(4)} \\
 &= \underbrace{u_{L/K}}_2 \cup \underbrace{\delta(\chi(\phi_{L/K}(a)))}_0 && \text{Theorem 12.10.3(3)} \\
 &= \underbrace{u_{L/K}}_2 \cup \underbrace{\delta\left(\frac{r}{n}\right)}_0 \\
 &= \underbrace{u_{L/K}}_2 \cup \underbrace{r}_0 && (14.4) \\
 &= ru_{L/K} && \text{Theorem 12.10.3(1)} \\
 \text{inv}_K(\bar{a} \cup \delta\chi) &= \frac{r}{n} = \chi(\phi_{L/K}(a)).
 \end{aligned}$$

In (14.4), we use the map in the snake lemma to calculate $\delta\left(\frac{r}{n}\right)$: it pulls back to $\frac{r}{n} \in \mathbb{Q} \cong H_T^{-1}(G, \mathbb{Q})$; the norm maps it to $r = n \cdot \frac{r}{n} \in \mathbb{Q} \cong H_T^0(G, \mathbb{Q}) \supseteq H_T^0(G, \mathbb{Z})$. In the second-to-last line, we note that $\bullet \cup r$ is simply multiplication by r in dimension 0, so Theorem 12.10.3(1) tells us it is multiplication by r in dimension 2 as well. \square

We need several naturality properties of the reciprocity map.

⁵It may seem odd to calculate $\chi \circ \phi_{L/K}$ instead of $\phi_{L/K}$ directly but keep in mind that for general L/K , $\text{Frob}_{L/K}(\mathfrak{p})$ is only defined to be a *conjugacy class*, and it is natural to look at the action of characters on conjugacy classes because characters are class functions.

Theorem 14.4.10. *Let $M/L/K$ be Galois extensions. The following are commutative.*

$$\begin{array}{ccc}
 A_L & \xrightarrow{\text{Cor}^0 = \text{Nm}_{L/K}} & A_K \\
 \phi_{M/L} \downarrow & & \phi_{M/K} \downarrow \\
 G(M/L)^{ab} & \xrightarrow[\text{natural}]{\text{Cor}^{-2}} & G(M/K)^{ab}
 \end{array}
 \qquad
 \begin{array}{ccc}
 A_K & \xrightarrow{\text{Res}^0 = i} & A_L \\
 \phi_{M/K} \downarrow & & \downarrow \phi_{M/L} \\
 G(M/K)^{ab} & \xrightarrow[\text{V}]{\text{Res}^{-2}} & G(M/L)^{ab}
 \end{array}$$

$$\begin{array}{ccc}
 A_K & \xrightarrow{s^*} & A_{sK} \\
 \phi_{L/K} \downarrow & & \downarrow \phi_{sL/sK} \\
 G(L/K)^{ab} & \xrightarrow{s^*} & G(sL/sK)
 \end{array}
 \qquad
 \begin{array}{ccc}
 A_K & & \\
 \phi_{M/K} \downarrow & \searrow \phi_{L/K} & \\
 G(M/K)^{ab} & \longrightarrow & G(L/K)^{ab}.
 \end{array}$$

Proof. First note that the maps in the first diagram are corestrictions and the maps in the second diagram (on the right) are restrictions by Proposition 12.11.4.

From axiom 4 of Proposition 14.4.5, we have

$$\text{Res}_{K/L}(u_{M/K}) = u_{M/L}.$$

We will use Proposition 12.11.9, about the commutativity of cup products with restriction and corestriction. The first diagram follows from

$$\text{Cor}_{L/K}^0(x \cup u_{M/L}) = \text{Cor}_{L/K}^0(x \cup \text{Res}_{K/L}(u_{M/K})) = \text{Cor}_{L/K}^2(x) \cup u_{M/K}, \quad x \in G(M/L)^{ab}.$$

The second diagram follows from

$$\text{Res}_{K/L}^0(x \cup u_{M/K}) = \text{Res}_{K/L}^{-2}(x) \cup u_{M/L}.$$

The third diagram follows from the fact that the map $s^* : AL \rightarrow A_{sK}$ takes $u_{L/K}$ to $u_{sL/sK}$.

For the last diagram, let χ be a character on $G(L/K)$, which gives a character $\chi_{M/K}$ on $G(M/K)$ using the projection $G(M/K) \rightarrow G(L/K)$. By Theorem 14.4.9 we have, for any character χ ,

$$\chi_{M/K}(\phi_{M/K}(a)) = \text{inv}_K(\bar{a}_{M/K} \cup \delta\chi_{M/K}) = \text{inv}_K(\bar{a}_{L/K} \cup \delta\chi) = \chi(\phi_{L/K}(a))$$

where $\bar{a}_{M/K}, \bar{a}_{L/K}$ are the images in $H_T^0(M/K)$ and $H_T^0(L/K)$, respectively. \square

The fourth diagram means that the maps $\phi_{L/K}$ are compatible, so we can define

$$\phi_K = \varprojlim_L \phi_{L/K} : A \rightarrow G^{ab}.$$

(Note $A = \cup A^H$.)

Theorem 14.4.11 (Norm limitation). *Let $(G, \{G_K\}, \{A_K\}, \text{inv}_{L/K})$ be a class formation. Let L/K be an extension and E/K be the largest abelian subextension. Then*

$$\text{Nm}_{L/K} A_L = \text{Nm}_{E/K} A_E.$$

Proof. Let L^{gal} be the Galois closure of L . Transitivity of norms (just look at the definition of norm...) gives us \subseteq . Conversely, suppose $a \in \text{Nm}_{E/K} A_E$. Let $G = G(L^{\text{gal}}/K)$ and $H = G(L'/L)$. Since E is the largest abelian subextension of L^{gal} abelian over K and contained in L , the subgroup of G fixing it is $G'H$. We have the commutative diagram

$$\begin{array}{ccc}
 A_L & \xrightarrow{\phi_{L^{\text{gal}}/L}} & H/H' \\
 \downarrow \text{Nm}_{L/K} & & \downarrow i \\
 A_K & \xrightarrow{\phi_{L^{\text{gal}}/K}} & G/G' \\
 & \searrow \phi_{E/K} & \downarrow \\
 & & G/G'H
 \end{array}$$

where i is induced by inclusion. Because $a \in \text{Nm}_{E/K} A_E$, $\phi_{E/K}(a) = 1$ in $G/G'H$. Thus $\phi_{L^{\text{gal}}/K}(a) \in G'H/G'$, and $\phi_{L^{\text{gal}}/K}(a)$ is in the image of i and hence $i \circ \phi_{L'/L}$, and there exists $b \in A_L$ such that $\phi_{L^{\text{gal}}/K}(a) = i(\phi_{L^{\text{gal}}/L}(b))$. Then

$$\phi_{L^{\text{gal}}/K}(a) = i(\phi_{L^{\text{gal}}/L}(b)) = \phi_{L^{\text{gal}}/K}(\text{Nm}_{L/K}(b)).$$

This means $\frac{a}{\text{Nm}_{L/K}(b)} \in \ker(\phi_{L^{\text{gal}}/K}) = \text{Nm}_{L^{\text{gal}}/K}(A_{L'})$; say it equals $\text{Nm}_{L^{\text{gal}}/K}(c)$. Then

$$a = \text{Nm}_{L/K}(b \text{Nm}_{L^{\text{gal}}/L}(c)) \in \text{Nm}_{L/K}(A_L),$$

as needed. □

Definition 14.4.12: A subgroup S of A_K is a **norm group** if there exists an extension L/K such that $S = \text{Nm}_{L/K}(A_L)$.

Theorem 14.4.13 (Bijective correspondence). *Let $(G, \{G_K\}, \{A_K\}, \text{inv}_{L/K})$ be a class formation. Then there is a bijective correspondence between finite abelian extensions of K and the set of norm groups of A_K , given by*

$$L \mapsto \text{Nm}_{L/K}(A_L).$$

Furthermore, this is an inclusion-reserving bijection that takes intersections to products and products to intersections:

$$\begin{aligned}
 L \subseteq M &\iff \text{Nm}_{L/K}(A_L) \supseteq \text{Nm}_{M/K}(A_M) \\
 \text{Nm}_{L \cdot L'/K}(A_{L \cdot L'}) &= \text{Nm}_{L/K}(A_L) \cap \text{Nm}_{L'/K}(A_{L'}) \\
 \text{Nm}_{L \cap L'/K}(A_{L \cap L'}) &= \text{Nm}_{L/K}(A_L) \cdot \text{Nm}_{L'/K}(A_{L'}).
 \end{aligned}$$

Finally, every subgroup of A_K containing a norm group is a norm group.

Proof. Abbreviate $\text{Nm}_{L/K}(A_L)$ by N_L .

First we show $N_{LL'} = N_L \cap N_{L'}$. By reciprocity,

$$N_L \cap N_{L'} = \ker(\phi_{L/K}) \cap \ker(\phi_{L'/K}) \stackrel{(*)}{=} \ker(\phi_{LL'/K}) = N_{LL'}$$

where $(*)$ comes from compatibility of the ϕ and the fact that the map $G(LL'/K) \rightarrow G(L/K) \times G(L'/K)$ is injective.

If $L \subseteq M$, then $N_L \supseteq N_M$ from transitivity of norms. Conversely, if $N_L \supseteq N_M$, then by the above $N_L = N_L N_M = N_{LM}$. Thus $[A_K : N_L] = [A_K : N_{LM}]$, and reciprocity gives $[L : K] = [LM : K]$, i.e. $LM = L$, i.e. $L \subseteq M$. Thus, $L \mapsto N_L$ is an inclusion-reversing bijection.

Next we show that every subgroup containing a norm group is a norm group. Suppose $N_L \subseteq N$; we show N is a norm group. We have that $\phi_{L/K}$ maps N isomorphically onto $G(L/K')$, where $K' = L^{\phi_{L/K}(N)}$, the fixed field of $\phi_{L/K}(N)$. Consider the following commutative diagram from Theorem 14.4.10:

$$\begin{array}{ccc} A_K & \xrightarrow{\phi_{L/K}} & G(L/K) \\ & \searrow \phi_{K'/K} & \downarrow \\ & & G(K'/K). \end{array}$$

From this we find

$$N = \ker(\phi_{K'/K}) = N_{K'}$$

as needed.

Finally, we show $N_{L \cap L'} = N_L \cdot N_{L'}$. Note $L \cap L'$ is the largest extension contained in both L and L' , while $N_L \cdot N_{L'}$ is the smallest group containing both N_L and $N_{L'}$, and it is a norm group by the above. Since $L \mapsto N_L$ is an inclusion-reversing bijection, we must have $N_{L \cap L'} = N_L \cdot N_{L'}$. \square

4.2 Class formations for local class field theory

As promised, we apply the results of the last section to $(G(\overline{K}/K), \overline{K})$ where K is a local field. (In the global case we will set A to be the ideles instead.)

Theorem 14.4.14. *Let L be a local field. Then*

$$(G(\overline{K}/K), \{G(L/K) : L/K \text{ finite Galois}\}, \overline{K})$$

is a class formation.

Proof. We verify the axioms of class formations.

1. $H^1(L/K) = 0$ for every cyclic extension of prime degree, by Hilbert's Theorem 90 (13.1.1).

2. Take the composition of the isomorphism $H^2(K) \cong H^2(K^{\text{ur}}/K)$ of Theorem 14.3.1 with the invariant map $H^2(K^{\text{ur}}/K) \rightarrow \mathbb{Q}/\mathbb{Z}$ to get

$$\text{inv}_K : H^2(K) \rightarrow \mathbb{Q}/\mathbb{Z}.$$

The maps $\text{inv}_{L/K} : H^2(L/K) \hookrightarrow H^2(K) \rightarrow \mathbb{Q}/\mathbb{Z}$ are isomorphisms onto their image, which must be $\frac{1}{[L:K]}\mathbb{Z}/\mathbb{Z}$.

Now we verify that

$$\text{inv}_L \circ \text{Res}_{K/L} = n \text{inv}_K, \quad n = [L : K].$$

This follows from the following commutative diagram. From Theorem 14.2.3, the right square commutes; from the fact that inflation commutes with restriction (by functoriality), the left square commutes.

$$\begin{array}{ccccc} H^2(K) & \xleftarrow[\cong]{\text{Inf}} & H^2(K^{\text{ur}}/K) & \xrightarrow{\text{inv}_{K^{\text{ur}}/K}} & \mathbb{Q}/\mathbb{Z} \\ \downarrow \text{Res}_{K/L} & & \downarrow \text{Res}_{K/L} & & \downarrow n \\ H^2(L) & \xleftarrow[\cong]{\text{Inf}} & H^2(L^{\text{ur}}/L) & \xrightarrow{\text{inv}_{L^{\text{ur}}/L}} & \mathbb{Q}/\mathbb{Z}. \end{array}$$

(Note that the target of the restriction in the middle is $H^2(K^{\text{ur}}/L)$, which is a subgroup of $H^2(L^{\text{ur}}/L)$.) □

Applying results about class field theory, we get the main results of local class field theory, restated below.

Theorem (Local reciprocity law, Theorem 11.2.1). *For any nonarchimedean local field K , there exists a unique homomorphism*

$$\phi_K : K^\times \rightarrow G(K^{\text{ab}}/K),$$

called the **local Artin (reciprocity) map** with the following properties.

1. (Relationship with Frobenius map) For any prime element π of K and any finite unramified extension L of K , $\phi_K(\pi)$ acts on L as $\text{Frob}_{L/K}(\pi)$.
2. (Isomorphism) Let p_L be the projection $G(K^{\text{ab}}/K) \rightarrow G(L/K)$. For any finite abelian extension L/K , ϕ_K induces an isomorphism $\phi_{L/K} : K^\times / \text{Nm}_{L/K}(L^\times) \rightarrow G(L/K)$ making the following commute:

$$\begin{array}{ccc} K^\times & \xrightarrow{\phi_K} & G(K^{\text{ab}}/K) \\ \downarrow & & \downarrow p_L \\ K^\times / \text{Nm}_{L/K}(L^\times) & \xrightarrow[\cong]{\phi_{L/K}} & G(L/K). \end{array}$$

3. (Compatibility with norm map) For any $K \subseteq K'$, the following diagram commutes.

$$\begin{array}{ccc} K'^{\times} & \xrightarrow{\phi_{K'}} & G(K'^{ab}/K') \\ \downarrow \text{Nm}_{K'/K} & & \downarrow \bullet|_{K^{ab}} \\ K^{\times} & \xrightarrow{\phi_K} & G(K^{ab}/K) \end{array}$$

Proof. By Theorem 14.4.14, $(G(\overline{K}/K), \{G(L/K) : L/K \text{ finite Galois}\}, \overline{K})$ is a class formation. By the Abstract Reciprocity Law applied to $A_K = K$, we thus have an isomorphism $K^{\times}/\text{Nm}_{L/K} L^{\times} \xrightarrow{\cong} G(L/K)^{ab}$. These maps are compatible by the first and fourth diagrams in Theorem 14.4.10.

Next we show that $\phi_K(\pi)$ acts on L as $\text{Frob}_{L/K}$. For the first, we use Theorem 14.4.9, which says

$$\chi(\phi_{L/K}(\pi)) = \text{inv}_K(\overline{\pi} \cup \delta\chi).$$

We calculate the invariant map on $\overline{\pi} \cup \delta\chi$, recalling that the map $H^1(G, \mathbb{Q}/\mathbb{Z}) \rightarrow \mathbb{Q}/\mathbb{Z}$ is evaluation at the Frobenius:

$$H^2(L/K) \longrightarrow H^2(G, \mathbb{Z}) \xleftarrow{\delta} H^1(G, \mathbb{Q}/\mathbb{Z}) \longrightarrow \mathbb{Q}/\mathbb{Z}$$

$$\overline{\pi} \cup \delta\chi \longrightarrow v(\pi) \cup \delta\chi = 1 \cup \delta\chi \xleftarrow{\quad} 1 \cup \chi \longrightarrow \chi(\text{Frob}_{L/K}).$$

Thus $\chi(\phi_{L/K}(\pi)) = \chi(\text{Frob}_{L/K})$ for all characters χ on $G(L/K)$, and $\phi_{L/K}(\pi) = \text{Frob}_{L/K}$.

We will prove uniqueness in Section 8.1 □

Proof of norm limitation, Theorem 11.2.6. This follows directly from Theorem 14.4.14 and Theorem 14.4.11. □

5 Examples

Before we move on to the existence theorem, we seek to understand the reciprocity map a bit better.

5.1 Unramified case

The reciprocity map is easiest to understand for unramified extensions.

Example 14.5.1: Suppose L/K is an unramified extension of local fields of degree n (possibly infinite). Then the reciprocity map is

$$\begin{aligned} \phi_{L/K} : K^{\times}/\text{Nm}_{L/K}(L^{\times}) &\cong K^{\times}/\pi^{n\mathbb{Z}}U_K \rightarrow G(L/K) \\ a &\mapsto \text{Frob}_{L/K}^{v(a)}. \end{aligned}$$

Proof. There are many ways to see this. We know that any uniformizer maps to $\text{Frob}_{L/K}$. But the uniformizers generate K^\times , so $\phi_{L/K}$ must be the map $a \mapsto \text{Frob}_{L/K}^{v(a)}$. As $\text{Frob}_{L/K}$ has order n , the kernel is $\pi^{n\mathbb{Z}}U_K$.

Alternatively, in the proof of Theorem 11.2.1 above, run the argument with arbitrary a instead of π . \square

5.2 Ramified case

To understand the reciprocity map on ramified extensions, we have the following.

Proposition 14.5.2: For any Galois extension of local fields L/K ,

$$\phi_{L/K}(U_K) \subseteq I(L/K),$$

where $I(L/K)$ is the inertia group.

Proof. By Theorem 2.7.2, $L^{I(L/K)}/K$ is the maximal unramified subextension of L/K , so $U_K \subseteq \ker(\phi_{L^{I(L/K)}/K})$ from Example 14.5.1. This means that $\phi_{L/K}(U_K)$ projects trivially on $G(L^{I(L/K)}/K)$, i.e. $\phi_{L/K}(U_K) \subseteq I(L/K)$. \square

In fact, the reciprocity map relates filtration on the unit group U_K with the filtration on ramification groups (cf. Section 4.2), so Proposition 14.5.2 is just the beginning of the story.

Theorem 14.5.3. *The reciprocity map transforms the filtration*

$$K^\times / \text{Nm}_{L/K}(L^\times) \supseteq U_K / \text{Nm}_{L/K}(U_L) \supseteq U_K^{(1)} / \text{Nm}_{L/K}(U_L^{\psi(1)}) \supseteq \dots$$

into the filtration

$$G(L/K) \supseteq G^0 = I(L/K) \supseteq G(L/K)^1 \supseteq \dots$$

Proof. This uses more about local fields and local symbols than we'll prove. See Serre [Ser79], Chapter XV or Neukirch [Neu99], V.§6. \square

Example 14.5.4: For the totally ramified extension $\mathbb{Q}_p(\zeta_{p^\infty})/\mathbb{Q}_p$, the reciprocity map sends

$$p^{\mathbb{Z}}(1 + (p^r)) \mapsto G(\mathbb{Q}_p(\zeta_{p^\infty})/\mathbb{Q}_p(\zeta_{p^r})).$$

The RHS is the r th upper ramification group G^r .

Explicit computation of the reciprocity map in the ramified case is difficult without Lubin-Tate Theory.

6 Hilbert symbols

To prove the existence theorem, we need to show that every closed subgroup of G occurs as a norm group, i.e. as the kernel of some Artin map $\phi_{L/K}$. To do this, we explicitly construct field extensions L/K that give these norm groups. We will construct Kummer extensions, extensions that come from adjoining an n th root. We focus on these extensions for several reasons.

1. Recall that we don't have a way to directly calculate the action of $\phi_{L/K}$. Instead, we calculate indirectly by Theorem 14.4.9: If we know $\chi(\phi_{L/K}(a))$ for all characters on $G(L/K)$, then we have determined $\phi_{L/K}(a)$.

An easy source of characters comes from Kummer Theory 13.2.2, since the group of characters is isomorphic to a cyclic group.⁶

2. We want to show that certain subgroups of norm groups are also norm groups. After verifying several topological properties of ϕ_K , we can reduce this to a statement about p th powers/roots of norm groups. In the abstract existence theorem 14.7.2, properties 1 and 3 are easy to check; they are basically the reductions that allow property 2 to be sufficient.

Recally from Proposition 13.2.2 that $K^\times/K^{\times n} \cong \text{Hom}(G(K^s/K), \mu_n)$. Thus the characters we get are in bijection with elements of $K^\times/K^{\times n}$. We can also consider $a \in K^\times$ as inside $K^\times/K^{\times n}$, and this gives us a sort of "duality": the Kummer pairing. We will see eventually that this is the source of reciprocity laws (Section 1), so these symbols are good for more than just proving the existence theorem.

We assume throughout that K contains a n th root of unity, and $\text{char}(K) \nmid n$.

Definition 14.6.1: Let $G = G(K^s/K)$. Define the local symbol

$$\begin{aligned}
 (\cdot, \cdot)_n : H^1(G, \mathbb{Q}/\mathbb{Z}) \times \underbrace{H^0(G, K^{s\times})}_{K^\times} &\rightarrow H^2(G, K^{s\times}) = \text{Br}_K \\
 (\chi, b) &= \bar{b} \cup \delta\chi
 \end{aligned}$$

Here δ is with respect to the exact sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{Z}/\mathbb{Q} \rightarrow 0$ and \bar{b} is the image of $K^{s\times}$ in $H_T^0(G, K^{s\times})$.

We will drop the subscript n when the context is clear.

Since cup product is bilinear and δ is linear, (\cdot, \cdot) is bilinear. If K is local, by Theorem 14.4.9, we have for any Galois L/K and any character χ on $G(L/K)$,

$$\text{inv}_K(\chi, \phi_{L/K}(a)) = \text{inv}_K(a \cup \delta\chi) = \chi(\phi_{L/K}(a)). \quad (14.5)$$

As promised, we now transfer this action to $K^\times/K^{\times n}$.

⁶Artin-Schreier theory, from exercise 13.2.1, is another source of characters.

Definition 14.6.2: Suppose K is a local field, and let $G = G(K^s/K)$. For $a \in K^\times$, define the character as in Proposition 13.2.2 by

$$\chi_a(\sigma) = \frac{\sigma(a^{\frac{1}{n}})}{a^{\frac{1}{n}}}, \quad \chi_a \in H^1\left(G, \frac{1}{n}\mathbb{Z}/\mathbb{Z}\right) \cong H^1(G, \mu_n),$$

where $G = G(L/K)$ and $L = K(a^{\frac{1}{n}})$. Here we choose a root of unity ζ to make a correspondence $\frac{1}{n}\mathbb{Z}/\mathbb{Z} \cong \mu_n$.

Define the **Hilbert symbol** by

$$\begin{aligned} K^\times \times K^\times &\rightarrow \text{Br}_K[n] \cong \frac{1}{n}\mathbb{Z}/\mathbb{Z} \cong \mu_n \\ (a, b) &:= (\chi_a, b) = b \cup \delta\chi_a. \end{aligned}$$

If K is a global field, let $(a, b)_v$ denote the Hilbert symbol where a, b are considered as elements of K_v .

Note that the image is in $\frac{1}{n}\mathbb{Z}/\mathbb{Z}$, not just in \mathbb{Q}/\mathbb{Z} , because $n\chi_a = 0$.

We'll abuse notation and not make a clear distinction between $\text{Br}_K[n] \cong \frac{1}{n}\mathbb{Z}/\mathbb{Z} \cong \mu_n$, where $\text{Br}_K[n]$ denotes the n -torsion subgroup of Br_K . The first isomorphism is given by inv_K and the second by $\frac{1}{n} \leftrightarrow \zeta$. We transfer the χ_a from being defined on μ_n to $\frac{1}{n}\mathbb{Z}/\mathbb{Z}$, then transfer back from $\text{Br}_K[n] \cong \frac{1}{n}\mathbb{Z}/\mathbb{Z}$ to μ_n at the end, so we may as well use the formula (14.5) for the χ_a treated in $H^1(G, \mu_n)$.

The following relates the Hilbert symbol to the Artin map.

Proposition 14.6.3: We have

$$(a, b) = \frac{[\phi_{L/K}(b)](\sqrt[n]{a})}{\sqrt[n]{a}}$$

where $L = K(\sqrt[n]{a})$.

Proof. Formula (14.5) gives (remember we're identifying $\text{Br}_K \cong \frac{1}{n}\mathbb{Z}/\mathbb{Z} \cong \mu_n$; by abuse of notation we drop the "inv $_K$ " because it is an isomorphism)

$$(a, b) = (\chi_a, \phi_{L/K}(b)) = \chi_a(\phi_{L/K}(b)) = \frac{[\phi_{L/K}(b)](\sqrt[n]{a})}{\sqrt[n]{a}}$$

where L is any field Galois over K , containing $\sqrt[n]{a}$. □

Theorem 14.6.4. *The Hilbert symbol descends to a nondegenerate skew-symmetric bilinear map*

$$K^\times/K^{\times n} \times K^\times/K^{\times n} \rightarrow \mu_n$$

satisfying the following.

1. $(a, b) = 1$ iff $b \in \text{Nm}_{K(a^{\frac{1}{n}})/K}(K(a^{\frac{1}{n}})^\times)$.

2. If $a \in K^\times$, $x \in K^\times$, and $x^n - a \neq 0$, then

$$(a, x^n - a) = 1.$$

In particular, $(a, -a) = 1 = (a, 1 - a)$.

Proof. Everything that went into defining $(,)$ was linear in either variable (cup products, evaluation homomorphisms, snake lemma morphism), so $(,)$ gives a bilinear map $K^\times \times K^\times \rightarrow \mu_n$.

Suppose χ is an element of order n . Then its kernel $\ker(\chi)$ has index n in $G(K^s/K)$. Under the Artin map this corresponds to a extension L_χ of degree n , such that $\ker(\chi) = \phi_K(\text{Nm}_{L_\chi/K}(L_\chi^\times))$. Then

$$(\chi, b) = \chi(\phi_K(b)) = 0 \iff \phi_K(b) \in \ker(\chi)$$

iff $b \in \text{Nm}_{L_\chi/K}(L_\chi^\times)$.

We apply this to $\chi = \chi_a$. Note that χ has order $[K(a^{\frac{1}{n}}) : K]$ and $\chi_a(G(K^s/K(a^{\frac{1}{n}}))) = 0$. Hence $\phi_K(\text{Nm}_{K(a^{\frac{1}{n}})/K}(K(a^{\frac{1}{n}})^\times)) \subseteq \ker \chi_a$. By comparing indices in $G(K^s/K)$, equality holds, giving the first item.

For the second item, note that

$$x^n - a = \prod_{j=0}^{n-1} (x - \zeta_n^j a^{\frac{1}{n}})$$

(for any choice of n th root). The factors in the product can be grouped into conjugates over K , so $x^n - a$ is a norm from $K(a^{\frac{1}{n}})/K$. Then $(a, x^n - a) = 1$ from the first item. Setting $x = 0, 1$ gives $(a, -a) = 1$ and $(a, 1 - a) = 1$.

To show skew-symmetry, note from item 2 and bilinearity that

$$1 = (ab, -ab) = (a, -a)(a, b)(b, a)(b, -b) = (a, b)(b, a).$$

To show nondegeneracy, suppose $b \in K^\times$ such that $(a, b) = 1$ for all $a \in K^\times$; we show $b \in K^{\times n}$. The condition $(a, b) = 1$ translates into $\chi_a(\phi_K(b)) = 1$ for all a . Now the image of ϕ_K is dense in $G(L/K)^{\text{ab}}$ (because it is surjective for every finite extension L/K , and $G(L/K)$ has the profinite topology). Hence $\chi_a = 0$. This means $a^{\frac{1}{n}} \in K$, i.e. $a \in K^{\times n}$. \square

Corollary 14.6.5. *Suppose K is a local field, $K(a^{\frac{1}{n}})/K$ is unramified, and b is a unit in K . Then $(a, b) = 1$.*

If K is a global field, then $(a, b)_v = 1$ in K_v unless either a or b is not a unit in K_v , or $K(a^{\frac{1}{n}})/K$ is ramified (which happen at finitely many places).

Proof. Since $K(a^{\frac{1}{n}})/K$ is unramified, $U_K \subset \text{Nm}_{K(a^{\frac{1}{n}})/K}(K(a^{\frac{1}{n}})^\times)$. The result now follows from Theorem 14.6.4.

The second part says that $(a, b)_v = 1$ if a, b are units in K_v and $K(a^{\frac{1}{n}})/K$ is unramified, which is clear from part 1. \square

Remark 14.6.6: In fact, $(a, b) = i(\chi_a \cup \chi_b)$ where $i : H^2(G, \mathbb{Z}/n\mathbb{Z}) \rightarrow \text{Br}_K$. (See Serre, p. 207.) This explains the symmetry better but takes more work to prove.

7 Existence theorem

We show that the existence theorem follows from several further (topological) axioms on formations. We then prove that in local class field theory, these axioms are satisfied.

7.1 Existence theorem in the abstract

First, a definition.

Definition 14.7.1: Let $(G, \{G_K\}_{K \in X}, A)$ be a class formation. The **universal norm group** D_K of K is the intersection of all norm groups of A_K :

$$D_K = \bigcap_{L/K} \text{Nm}_{L/K}(A_L).$$

Theorem 14.7.2 (Abstract existence). *Suppose that $(G, \{G_K\}_{K \in X}, A)$ is a formation satisfying the following conditions.*

1. *For every extension L/K , the norm homomorphism has closed image and compact kernel.*
2. *Let $[p]$ denote the map $x \mapsto px$ on A . For every prime p , there exists a field K_p such that for K containing K_p , $\ker([p]|_{A_K})$ is compact and $\text{im}([p]|_{A_K})$ contains D_K .*
3. *There exists a compact subgroup U_K of A_K such that every closed subgroup of finite index in A_K containing U_K is a norm group.*

Then a subgroup of A_K is a norm group iff it is closed of finite index.

If the conclusion holds, $nA_K \subseteq D_K$ for every K , because nA_K is closed of finite index and hence a norm group. Conversely, $D_K \subseteq \bigcap_{n \geq 1} nA_K$ because every norm group N has finite index so n kills A_K/N for some n . Furthermore, D_K must be divisible: else we could find a norm group $N \supseteq D_K$, and n such that $nN \not\supseteq D_K$, even though nN is still of finite index. (Note we write A_K additively here, but in class field theory, $A_K = K$ and nA_K actually means A_K^n .) The most important condition is item 2, because it will give us these two conditions. This gives us a large set of norm groups, and items 1 and 3 (which are more topological in nature) will give us the rest of the desired norm groups.

Proof. Step 1: Suppose axiom 1 holds. We show that for every extension L/K , $\text{Nm}_{L/K}(D_L) = D_K$.

By transitivity of norms, $\text{Nm}_{L/K}(D_L) \subseteq D_K$.

Conversely, suppose $a \in D_K$. Since $a \in D_K$, for any extension M/L , A_M contains an element b such that $\text{Nm}_{M/K}(b) = \text{Nm}_{L/K} \text{Nm}_{M/L}(b) = a$. Thus

$$S_M := \text{Nm}_{L/K}^{-1}(a) \cap \text{Nm}_{M/L}(A_M)$$

is nonempty. Since Nm has compact kernel, the first group is compact; since Nm has closed image, the second group is closed; thus S_M is compact. Since the S_M for all M/L form a directed system of compact subsets, $S = \bigcap_M S_M$ is nonempty. Any element of S is an element of $\text{Nm}_{L/K}^{-1}(a) \cap D_L$. This shows $a \in \text{Nm}_{L/K}(D_L)$.

Step 2: Suppose axioms 1 and 2 hold. We show D_K is divisible and

$$D_K = \bigcap_{n \geq 1} nA_K.$$

First we show that for every prime p , $pD_K = D_K$. Let L be a field containing K_p , $a \in D_K$, and set

$$S_L = [p]^{-1}(a) \cap \text{Nm}_{L/K} A_L.$$

Since $[p]^{-1}(a)$ is compact (as $\ker([p]|_{A_K})$ is compact by axiom 2) and $\text{Nm}_{L/K} A_L$ is closed, S_L is compact. Now this set is nonempty: since $a \in D_K = \text{Nm}_{L/K} D_L$ by step 1, we can write $a = \text{Nm}_{L/K} x$, $x \in D_L$. By axiom 2, $x = py$ with $y \in A_K$, so $b := \text{Nm}_{L/K} y \in S_L$. Then $\bigcap_{L \supseteq K_p} S_L$ is nonempty as in step 1. Hence $a \in pD_K$.

This shows $pD_K = D_K$, and we get $D_K = \bigcap_{n \geq 1} nD_K \subseteq \bigcap_{n \geq 1} nA_K$.

For the other direction, note that na is the norm of any extension of degree n , so $\bigcap_{n \geq 1} nA_K \subseteq D_K$.

Step 3: Assume all the axioms. We prove the theorem.

First, note that any norm group is closed by axiom 1, and has finite index by the reciprocity law 14.4.8. Indeed, by transitivity of norm, it suffices to consider Galois extensions, and the reciprocity law says $\text{Nm}_{L/K}(A_L)$ has index equal to $G(L/K)^{\text{ab}}$.

Conversely, suppose S is a closed subgroup of finite index n . We will find a norm subgroup contained in S and then apply Theorem 14.4.13. Since A_K/S has order n , we get $D_K \subseteq nA_K \subseteq S$, so

$$\bigcap_{N \text{ norm group}} (N \cap U_K) = D_K \cap U_K \subseteq S.$$

Since $N \cap U_K$ are compact (N is closed and U_K is compact) and S is open (closed subgroups of finite index are also open), there exists N such that

$$N \cap U_K \subseteq S.$$

Note $U_K + (N \cap S)$ is closed of finite index in A_K because N, S are closed of finite index; we show we can replace U_K with $U_K + (N \cap S)$ above:

$$N \cap (U_K + (N \cap S)) \subseteq S.$$

Suppose $a \in U_K$ and $a' \in N \cap S$ such that $a + a' \in N$. Then $a \in N$, but $N \cap U_K \subseteq S$ so $a \in S$ as well. Thus $a + a' \in S$, as needed.

Now $N \cap (U_K + (N \cap S))$ is closed of finite index containing U_K , so is a norm group by axiom 3. By Theorem 14.4.13, we get S is also a norm group. \square

7.2 Existence theorem for local class field theory

Proof of Theorem 11.2.3. We verify that the class formation for LCFT satisfies the three axioms of Theorem 14.7.2.

1. To see that the norm map is closed, note that

$$\text{Nm}_{L/K}(L^\times) \cap U_K = \text{Nm}_{L/K}(U_L)$$

because an element is a unit iff its norm is a unit. As U_L is compact and $\text{Nm}_{L/K}$ is continuous (Proposition 8.1.6), $\text{Nm}_{L/K}(U_L)$ is compact and hence closed. Now $\text{Nm}_{L/K}(L^\times)$ is a union of translates of U_L , therefore closed as well.

The kernel of $\text{Nm}_{L/K}$ is a closed subset of U_L , hence compact.

2. Take K_p containing all p th roots of unity. The kernel of the p th power map is the p th roots of unity, which is a compact set. Suppose $K \supseteq K_p$, and let $b \in D_K$ be a universal norm. Then $(a, b) = 1$ for all a by Theorem 14.6.4. Since the p th power Hilbert symbol is nondegenerate on $K^\times/K^{\times p}$, $a \in K^{\times p}$. Thus $D_K \subseteq K^{\times p}$.
3. Take U_K to be the group of units of K^\times . The closed subgroups of finite index containing U_K are just $\pi^{n\mathbb{Z}}U_K$ for $n \neq 0$; these are the norm groups of unramified extensions of degree n by Proposition 14.5.1. (Note these extensions exist—just adjoin appropriate roots of unity.) \square

Proof of Theorem 11.2.5. This follows from Theorem 14.4.13, Theorem 14.4.14 (class formation for LCFT), and the existence theorem just proved. \square

Note the existence theorem gives the following.

Corollary 14.7.3. *The universal norm group D_K is $\{1\}$.*

Proof. All open subgroups of finite index are norm groups by the Existence Theorem 11.2.3. The intersection of all open subgroups of finite index is $\{1\}$, as $\bigcap_{m,n}(1+(\pi^m))\pi^{n\mathbb{Z}} = \{1\}$. \square

8 Topology of the local reciprocity map

We now prove that ϕ_K gives a topological isomorphism $K^\times \rightarrow W(L/K)$.

Proof of Theorem 11.2.4. By Proposition 14.5.2, $\phi_{L/K}(U_K) \subseteq I(L/K)$, so we have the commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & U_K & \longrightarrow & K^\times & \xrightarrow{v} & \mathbb{Z} & \longrightarrow & 1 \\ & & \downarrow \phi_{L/K} & & \downarrow \phi_{L/K} & & \downarrow & & \\ 1 & \longrightarrow & I(L/K) & \longrightarrow & G(L/K) & \longrightarrow & G(l/k) & \longrightarrow & 1. \end{array}$$

where the rightmost vertical map sends 1 to the p th power Frobenius ($p = |k|$). The vertical maps factor as

$$\begin{array}{ccccccc}
 1 & \longrightarrow & U_K / \text{Nm}_{L/K}(U_L) & \longrightarrow & K^\times / \text{Nm}_{L/K}(L^\times) & \xrightarrow{v} & \mathbb{Z}/f\mathbb{Z} \longrightarrow 1 \\
 & & \cong \downarrow \phi_{L/K} & & \cong \downarrow \phi_{L/K} & & \cong \downarrow \\
 1 & \longrightarrow & I(L/K) & \longrightarrow & G(L/K) & \longrightarrow & G(l/k) \longrightarrow 1.
 \end{array} \tag{14.6}$$

where $f = [l : k]$. Recall $\phi_K = \varprojlim_L \phi_{L/K}$. The intersection of all norm groups is $\{1\}$ by Corollary 14.7.3, so ϕ_K is injective on K^\times .

In forming $\phi_K = \varprojlim_L \phi_{L/K}$, we are really considering the embedding

$$K^\times \hookrightarrow \widehat{K^\times} := \varprojlim_L K^\times / \text{Nm}_{L/K}(L^\times) \xrightarrow{\cong} G(K^{\text{ab}}/K).$$

Decomposing $K^\times / \text{Nm}_{L/K}(L^\times)$ as in (14.6), we have that

1. $\varprojlim_L U_K / \text{Nm}_{L/K}(U_L) \cong U_K$ since U_K is compact, hence complete, so $U_K \cong I(K^{\text{ab}}/K)$.
2. $\varprojlim_L \mathbb{Z}/f\mathbb{Z} = \widehat{\mathbb{Z}}$.

Thus $K^\times \hookrightarrow \widehat{K^\times}$ is the embedding $U_K \times \pi^\mathbb{Z} \hookrightarrow U_K \times \pi^{\widehat{\mathbb{Z}}}$.

Recalling that $W(L/K)$ is the inverse image of $\text{Frob}^\mathbb{Z} \subseteq G(\bar{k}/k)$, we get $\phi_{L/K} : K^\times \rightarrow W(L/K)$ is a topological isomorphism. In summary, we have the diagram

$$\begin{array}{ccccccc}
 1 & \longrightarrow & U_K & \longrightarrow & K^\times & \longrightarrow & \pi^\mathbb{Z} \longrightarrow 1 \\
 & & \cong \downarrow \phi_K & & \cong \downarrow \phi_K & & \cong \downarrow \\
 1 & \longrightarrow & I(K^{\text{ab}}/K) & \longrightarrow & W(K^{\text{ab}}/K) & \longrightarrow & \text{Frob}^\mathbb{Z} \longrightarrow 1 \\
 & & \searrow & & \downarrow & & \downarrow \\
 & & & & G(K^{\text{ab}}/K) & \longrightarrow & \text{Frob}^{\widehat{\mathbb{Z}}} = G(\bar{k}/k) \longrightarrow 1
 \end{array}$$

□

8.1 Uniqueness of the reciprocity map

Finally, we prove uniqueness. This finishes all the proofs of local class field theory.

We first restate Lemma 11.6.7.

Lemma. *Suppose that K is a nonarchimedean local field, K^{ur} is the maximal abelian unramified extension of K , and L is an abelian extension containing K^{ur} . Let $f : K^\times \rightarrow G(L/K)$ be a homomorphism satisfying (1) and either (2) or (2)':*

1. *The composition $K^\times \xrightarrow{f} G(L/K) \rightarrow G(K^{\text{ur}}/K)$ takes α to $\text{Frob}_{K^{\text{ur}}/K}(\pi)^{v(\alpha)}$.*

2. For any uniformizer $\pi \in K$, $f(\pi)|_{K_\pi} = 1$, where

$$K_\pi := L^{\phi_K(\pi)}.$$

2'. For any finite subextension K'/K of K_π , we have

$$f(\text{Nm}_{K'/K}(K'^{\times}))|_{K'} = \{1\}.$$

Then f equals the reciprocity map ϕ_K .

Proof of Lemma 11.6.7. It suffices to prove this for $L = K^{\text{ab}}$. We have the split exact sequence

$$1 \rightarrow U_K^{\times} \rightarrow K^{\times} \xrightarrow{v} \mathbb{Z} \rightarrow 1, \quad (14.7)$$

where the splitting is determined by the map $\mathbb{Z} \rightarrow K^{\times}$ sending $1 \mapsto \pi$, and the map $K^{\times} \rightarrow U_K$ sending $a \mapsto \frac{a}{\pi^{v(a)}}$. Under the Artin map, (14.7) gets sent to the split exact sequence of topological groups

$$1 \rightarrow I(K^{\text{ab}}/K) = G(K/K^{\text{ur}}) \rightarrow W(K^{\text{ab}}/K) \rightarrow W(K^{\text{ur}}/K) \cong \mathbb{Z} \rightarrow 1$$

by Theorem 11.2.4. This gives the exact sequence

$$1 \rightarrow G(K^{\text{ab}}/K^{\text{ur}}) \rightarrow G(K^{\text{ab}}/K) \rightarrow G(K^{\text{ur}}/K) \rightarrow 1,$$

where the splitting is by the map $\mathbb{Z} \cong G(K^{\text{ur}}/K) \rightarrow G(K^{\text{ab}}/K)$ sending $1 \mapsto \phi_K(\pi)$. This identifies $G(K^{\text{ab}}/K^{\text{ur}})$ with the quotient group $G(K_\pi/K)$ where

$$K_\pi = L^{\overline{\langle \phi_K(\pi) \rangle}} = L^{\phi_K(\pi)}.$$

If (2)' holds, then for any uniformizer π , we have that $\pi \in \text{Nm}_{K'/K}(K'^{\times})$ for every finite subextension K' of K_π . Then (2)' gives that $f(\pi)|_{K_\pi} = 1$. Then (2) holds.

We now show if (1) and (2) hold, then $f = \phi$. Indeed, (1) and (2) imply that $\phi(\pi)|_{K^{\text{ur}}K_\pi} = f(\pi)|_{K^{\text{ur}}K_\pi}$ for any uniformizer π . But $K^{\text{ur}}K_\pi = K^{\text{ab}}$ and the set of uniformizers generate K^{\times} (any unit is the quotient of two uniformizers). Hence $\phi = f$. \square

Proof of uniqueness in Theorem 11.2.1. Suppose ϕ' is another map satisfying the conditions of Theorem 11.2.1. It suffices to show ϕ' satisfies the conditions of Lemma 11.6.7 with $L = K^{\text{ab}}$. By assumption it satisfies (1). For condition (2)', we have $\phi_K(\pi)|_{K_\pi} = 1$ by definition of K_π . Hence π is a norm from every finite subextension of K_π . By condition 2 of Theorem 11.2.1, this shows $\phi'_{K'/K}(\text{Nm}_{K'/K}(K'^{\times})) = \{1\}$ for every subextension K'/K of L , as needed. Hence $\phi' = \phi$. \square

Problems

1. Using ϕ_K , construct a natural bijection between the following two sets.
 - continuous characters $W(\overline{K}/K) \rightarrow \mathbb{C}^\times$ (i.e. continuous representations $W(\overline{K}/K) \rightarrow GL_1(\mathbb{C})$).
 - continuous character $K^\times \rightarrow \mathbb{C}$ (i.e. continuous homomorphisms $GL_1(K) \rightarrow GL(\mathbb{C})$).

This is the “local Langlands correspondence for GL_1 over K .” Local class field theory generalizes more naturally in this form.

Chapter 15

Global class field theory

To prove the global reciprocity law we need to do two things, namely construct a map

$$\phi_K : \mathbb{I}_K / K^\times \text{Nm}_{L/K} \mathbb{I}_L \xrightarrow{\cong} G(L/K),$$

and show that it is an isomorphism. To show it is an isomorphism, we need to show that the two sides have the same cardinality:¹

$$|\mathbb{I}_K / K^\times \text{Nm}_{L/K} \mathbb{I}_L| = [L : K].$$

The first inequality “ \geq ” will be shown using cohomology, with lots of Herbrand quotient calculations. The second inequality “ \leq ” is most easily shown with L -functions, but can also be shown with a more complicated cohomological argument.

To construct a map, there are two approaches. We can define ϕ_K to be the map whose components are the local Artin map, and use the properties of the local Artin map given by local class field theory. Alternatively, we can construct it directly in the global case, without using local theory, and get local class field theory as a corollary. We will take the first approach. For an account of the second, see Lang [Lan94].

1 Basic definitions

First, some basic definitions.

Definition 15.1.1: Define the action of $G(L/K)$ on \mathbb{I}_L by permuting the places: For an idele $\mathbf{a} = (a_v)_{v \in V_L}$, define $\sigma \mathbf{a}$ by

$$(\sigma \mathbf{a})_{\sigma(v)} = \sigma(a_v).$$

Definition 15.1.2: Define the inclusion map $\mathbb{I}_K \hookrightarrow \mathbb{I}_L$ by

$$(a_v)_{v \in V_K} \mapsto ((a_v)_{w|v})_{v \in V_K},$$

¹More precisely, we use this to show the invariant map is an isomorphism, then get the Artin map from the machinery of class formations.

i.e. it is induced by componentwise inclusions $K_v \hookrightarrow L^w$. Let the inclusion map $\mathbf{C}_K \hookrightarrow \mathbf{C}_L$ be induced by the above inclusion.

For an infinite extension M/K , define

$$\mathbb{I}_M = \varinjlim_{K \subseteq L \subseteq M} \mathbb{I}_L, \quad \mathbf{C}_M = \varinjlim_{K \subseteq L \subseteq M} \mathbf{C}_L$$

where the limit is taken over finite Galois extensions L/K .

For short, let $H^n(L/K, A)$ denote $H^n(G(L/K), A)$ and $H^2(K, A) := H^2(\overline{K}/K, A)$. As in the local case, $H^n(L/K)$ denotes $H^n(G(L/K), K^\times)$.

Proposition 15.1.3: Let L/K be a Galois extension and $G = G(L/K)$. The inclusion map $\mathbb{I}_K \hookrightarrow \mathbb{I}_L$ sends $\mathbb{I}_K \xrightarrow{\cong} \mathbb{I}_L^G$ and the inclusion map $\mathbf{C}_K \hookrightarrow \mathbf{C}_L$ sends $\mathbf{C}_K \xrightarrow{\cong} \mathbf{C}_L^G$.

Proof. The first part holds because G acts transitively on all the places in L dividing a single $v \in V_K$, so any element of \mathbb{I}_L^G has to be constant on all $w \mid v$, i.e. in the image of \mathbb{I}_K .

For the second part², take the long exact sequence in cohomology associated to

$$1 \rightarrow L^\times \rightarrow \mathbb{I}_L \rightarrow \mathbf{C}_L \rightarrow 1$$

to get

$$\begin{array}{ccccccc} 1 & \longrightarrow & H^0(G, L^\times) & \longrightarrow & H^0(G, \mathbb{I}_L) & \longrightarrow & H^0(G, \mathbf{C}_L) \longrightarrow H^1(G, L^\times) \\ & & \parallel & & \parallel & & \parallel & \parallel \\ & & K^\times & & \mathbb{I}_L^G = \mathbb{I}_K & & \mathbf{C}_L^G & 1 \end{array}$$

where the equality on the right is Hilbert's Theorem 90 (Theorem 13.1.1) and the map $\mathbb{I}_K \rightarrow \mathbf{C}_L^G$ is induced by inclusion. Thus $\mathbf{C}_L^G = \mathbb{I}_K/K^\times = \mathbf{C}_K$. \square

2 The first inequality

In this section we will prove the following.

Theorem 15.2.1 (First inequality of global class field theory). *If L/K is cyclic, then*

$$|\mathbb{I}_K/K^\times \text{Nm}_{L/K} \mathbb{I}_L| \geq [L : K].$$

To prove the inequality, we first express the left-hand side in terms of cohomology. Letting $G = G(L/K)$, we know that

$$H_T^0(G, \mathbf{C}_L) = \mathbf{C}_K / \text{Nm}_{L/K} \mathbf{C}_L = \mathbb{I}_K / K^\times \text{Nm}_{L/K} \mathbb{I}_L.$$

²which isn't obvious, because we're taking quotients here

Then noting that the Herbrand quotient (with respect to G) of \mathbf{C}_L is $h(\mathbf{C}_L) = \frac{|H_T^0(G, \mathbf{C}_L)|}{|H_T^{-1}(G, \mathbf{C}_L)|}$, we have that

$$|\mathbb{I}_K/K^\times \text{Nm}_{L/K} \mathbb{I}_L| = |H_T^0(G, \mathbf{C}_L)| \geq h(\mathbf{C}_L). \quad (15.1)$$

To calculate $h(\mathbf{C}_L)$ our plan is as follows.

1. First express \mathbf{C}_L in terms of something involving a finite set of places; we find T so that

$$\mathbb{I}_L = L^\times \mathbb{I}_L^T.$$

(Proposition 15.2.2). Then calculation shows that $h(\mathbf{C}_L) = \frac{h(\mathbb{I}_L^T)}{h(U_L^T)}$, where U_L^T denotes the T -units in L .

2. Compute $h(\mathbb{I}_L^S) = \prod_{v \in S} n_v$. Note \mathbb{I}_L^S is a direct product, not a restricted direct product, so we can just take the product of the Herbrand quotient of the factors. Breaking up the places into $G(L/K)$ -orbits, we can calculate $h(\mathbb{I}_L^S)$ using the corollary to Shapiro's Lemma 12.8.7.

3. Compute $h(U_L^S) = \frac{1}{n} \prod_{v \in S} n_v$ by relating it to a lattice of codimension 1 in \mathbb{R}^s by the log map, where $s = |S|$. (See ANT, Chapter 5.) We use the fact that the Herbrand quotient of a full lattice depends only on the vector space it resides in (Theorem 15.2.5) to change to a more convenient lattice whose basis consists of vectors representing the s places in U_L^S , i.e. the lattice $\Lambda = \prod_{w \in S} \mathbb{Z}e_w$.

The set S breaks up into $G(L/K)$ -orbits, so the lattice breaks up into induced S -modules, and we can calculate $h(U_L^S)$ using again using Shapiro's Lemma 12.8.7.

4. Putting all the steps together gives

$$h(\mathbf{C}_L) = n,$$

as needed.

2.1 Reduce to finite number of places

Proposition 15.2.2: Let L be a number field. There exists a finite set of places T of L such that

$$\mathbb{I}_L = L^\times \mathbb{I}_L^T.$$

Proof. This basically follows from the finiteness of the class group.

For the first part, consider the map $p : \mathbb{I}_L \rightarrow \text{Cl}_L$, defined by sending

$$(a_v)_{v \in V_L} \mapsto \prod_{v=v_{\mathfrak{p}} \in V_L^0} \mathfrak{p}^{v(a_{\mathfrak{p}})}.$$

(Map a to the prime ideal whose valuation at each prime equals the valuations of the corresponding coordinates of a .) The kernel—the set sent to the principal ideals—is $L^\times \mathbb{I}_L^{V_\infty}$,

where V^∞ is the set of infinite places. Thus we have an isomorphism $\mathbb{I}_L/L^\times \mathbb{I}_L^{V^\infty} \rightarrow \text{Cl}_L^3$. The latter is finite; take the inverse image of a set of generators A . We can choose finite T containing V^∞ so that the coordinates of elements of A are units outside of T . Then \mathbb{I}_L^T generates \mathbb{I}_L/L^\times , as needed. \square

2.2 Cohomology of \mathbb{I}_L^S and \mathbb{I}_L

Proposition 15.2.3: Let L/K be a Galois extension of number fields. Let S be a set of places in K and let $\mathbb{I}_L^S := \mathbb{I}_L^T$ where $T = \{w \in V_L : w \mid v \text{ for some } v \in S\}$. Then for any $i > 0$ we have

$$H^i(G, \mathbb{I}_L^S) = \prod_{v \in S} H^i(G(L^v/K_v), L^{v^\times}) \times \prod_{v \notin S} H^i(G(L^v/K_v), U^v).$$

This is also true for Tate groups if G is finite.

In particular, if L/K is cyclic, and S contains all ramified places, then

$$\begin{aligned} H^1(G, \mathbb{I}_L^S) &= 1 \\ H^2(G, \mathbb{I}_L^S) &= \prod_{v \in S} \frac{1}{n_v} \mathbb{Z}/\mathbb{Z} \\ h(\mathbb{I}_L^S) &= \prod_{v \in S} n_v \end{aligned}$$

where n_v is the local degree $[L_w : K_v]$, for any $w \mid v$.

Proof. We have

$$\mathbb{I}_L^S = \prod_{w \in T} L_w^\times \times \prod_{w \notin T} U_w$$

where $U_w := U_{K_w}$. We calculate the cohomology groups of each factor.

$$\begin{aligned} H^i \left(G, \prod_{w \in T} L_w^\times \right) &= H^i \left(G, \prod_{v \in S} \prod_{w \mid v} L_w^\times \right) \\ &= \prod_{v \in S} H^i \left(G, \prod_{w \mid v} L_w^\times \right) && \text{cohomology respects products, Proposition 12.6.7} \\ &= \prod_{v \in S} H^i(G^v, L^{v^\times}) && \text{by Corollary 12.8.8 to Shapiro's Lemma} \\ &= \prod_{v \in S} H^i(G(L^v/K_v), L^{v^\times}) && (15.2) \\ &= \begin{cases} 1, & i = 1, \\ \prod_{v \in S} \frac{1}{n_v} \mathbb{Z}/\mathbb{Z}, & i = 2. \end{cases} && (15.3) \end{aligned}$$

³cf. Example 11.5.10; there $\mathbb{I}_L^{V^\infty}$ is written as \mathbb{U}_L .

For $i = 1$, the last result follows from Hilbert's Theorem 90, and for $i = 2$, it follows from the fact that $\text{inv}_{K_v} : H^2(G(L^v/K_v), L^{v\times}) \xrightarrow{\cong} \frac{1}{n_v}\mathbb{Z}/\mathbb{Z}$ is an isomorphism (a consequence of the class formation for LCFT, Theorem 14.4.14, or actually just Theorem 14.3.1 and Proposition 14.2.1).

For the units, we have,

$$H^i\left(G, \prod_{w \notin T} U_w\right) = \prod_{v \notin S} H^i(G(L^v/K_v), U_w) \quad \text{Proposition 12.6.7} \quad (15.4)$$

$$= 1 \quad \text{if } T \text{ unramified, by Theorem 14.1.1.} \quad (15.5)$$

For the general case, take the product of (15.2) and (15.4). For the special case, take the product of (15.3) and (15.5). The Herbrand quotient calculation follows directly. \square

If we consider the full group \mathbb{I}_L , we get the following result. (We won't need this until Section 5.)

Proposition 15.2.4: For any Galois extension L/K with Galois group G and any $n \geq 0$, we have

$$H^n(G, \mathbb{I}_L) \cong \bigoplus_{v \in V_K} H^n(L^v/K_v).$$

This is also true for Tate groups when G is finite.

In particular, we have

1. $H^1(G, \mathbb{I}_L) = 0$.
2. $H^2(G, \mathbb{I}_L) = \bigoplus_{v \in V_K} \frac{1}{n_v}\mathbb{Z}/\mathbb{Z}$.

Proof. We have

$$\mathbb{I}_L = \varinjlim_{S \text{ finite}} \mathbb{I}_L^S.$$

Hence using Proposition 12.14.3,

$$\begin{aligned} H^n(G, \mathbb{I}_L) &= H^n(G, \varinjlim_S \mathbb{I}_L^S) \\ &= \varinjlim_S H^n(G, \mathbb{I}_L^S) \\ &= \begin{cases} \varinjlim_S \prod_{v \in S} H^n(G^v, L^{v\times}) \times \prod_{v \notin S} H^n(G^v, U^v) = \bigoplus_{v \in V_K} H^n(G^v, L^{v\times}), & \text{general case} \\ 1, & n = 1 \\ \bigoplus_{v \in V_K} \frac{1}{n_v}\mathbb{Z}/\mathbb{Z}, & n = 2 \end{cases} \end{aligned}$$

where the last statement follows from Proposition 15.2.3. \square

2.3 Cohomology of lattices and U_L^T

Proposition 15.2.5: Suppose G is finite cyclic, V is a finite real vector space and $\mathbb{R}[G]$ -module, and M, N are two lattices in V , stable under the action of G . Then

$$h(M) = h(N).$$

(If one is defined, so is the other.)

Proof. We proceed in 2 steps.

Step 1: We show that $M \otimes_{\mathbb{Z}} \mathbb{Q} \cong N \otimes_{\mathbb{Z}} \mathbb{Q}$ as G -modules. We know $M \otimes_{\mathbb{Z}} \mathbb{R} = V = N \otimes_{\mathbb{Z}} \mathbb{R}$. Suppose $V = \mathbb{R}^n$. Choose bases $\{\beta_i\}$ for M and $\{\gamma_i\}$ for N . Let $B(\sigma)$ and $C(\sigma)$ be matrices representing the action of a generator $\sigma \in G$ on these bases.⁴ A linear map $M \otimes_{\mathbb{Z}} \mathbb{R} \rightarrow N \otimes_{\mathbb{Z}} \mathbb{R}$ represented by a matrix A with respect to $\{\beta_i\}$ and $\{\gamma_i\}$ is a isomorphism of G -modules if

$$A \cdot B(\sigma) = C(\sigma) \cdot A.$$

These determine a system of homogeneous linear equations in the entries of A , with coefficients in \mathbb{Z} , since $B(\sigma)$ and $C(\sigma)$ have entries in \mathbb{Z} .

Letting the solution space be $W \subseteq \mathcal{M}_{n \times n}(\mathbb{R})$, we have

$$\dim_{\mathbb{R}} W = \dim_{\mathbb{Q}}(W \cap \mathcal{M}_{n \times n}(\mathbb{Q})),$$

because Gaussian elimination never needs to leave the world of \mathbb{Q} . Hence we can find a basis for W contained in $\mathcal{M}_{n \times n}(\mathbb{Z})$, say $\{A_1, \dots, A_k\}$. By the existence of an isomorphism between $M \otimes_{\mathbb{Z}} \mathbb{R}$ and $N \otimes_{\mathbb{Z}} \mathbb{R}$, there exist $a_1, \dots, a_k \in \mathbb{R}$ such that $a_1 A_1 + \dots + a_k A_k$ is nonsingular, i.e.

$$\det(a_1 A_1 + \dots + a_k A_k) \neq 0.$$

The left hand side is hence a nonzero polynomial in the a_k ; since it has coefficients in the infinite field \mathbb{Q} it has a solution over \mathbb{Q} . Taking A to be the corresponding linear combination, we get the desired G -isomorphism $M \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow N \otimes_{\mathbb{Z}} \mathbb{Q}$.

Step 2: We have an isomorphism $f : M \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow N \otimes_{\mathbb{Z}} \mathbb{Q}$; by scaling f (since M, N are finite-dimensional lattices) we may assume f restricts to $f : M \rightarrow N$. Now $N/f(M)$ is finite; hence by Proposition 12.12.4(1) and (2),

$$h(N) = h(M)h(N/f(M)) = h(M).$$

□

Proposition 15.2.6: Let L/K be a finite cyclic extension of number fields of degree n . Let S be a set of places in K containing the infinite places and $T = \{w \in V_L : w \mid v \text{ for some } v \in S\}$. We have

$$h(U_L^T) = \frac{1}{n} \prod_{w \in T} n_w$$

⁴ G cyclic is not important here; we could work with all elements of G .

where U_L^T denotes the T -units in L and n_w is the local degree $[L_w : K_v]$, where $w \mid v$.

Proof. Consider the map $L : U_L^T \rightarrow \mathbb{R}^T$ defined by letting

$$L(a) = (\ln |a|_w)_{w \in T}$$

where $|\cdot|_w$ is the normalized valuation. Then $L(a)$ is a lattice of dimension $|T| - 1$ by Dirichlet's S -unit theorem 5.3.2; it is in the hyperplane where the sum of coordinates is 0 (take the log of the product formula 19.1). The kernel of L consists the roots of unity in L , $\mu \cap L$, which is a finite group. By Proposition 12.12.4(1)–(2) applied to $1 \rightarrow \mu \cap L \rightarrow U_L^T \rightarrow L(U_L^T) \rightarrow 0$,

$$h(U_L^T) = h(\mu \cap L)h(L(U_L^T)) = h(L(U_L^T)) \quad (15.6)$$

Let $G(L/K)$ act on \mathbb{R}^T by permuting the coordinates corresponding to the places. Note that L is a G -module homomorphism with respect to this action. Let \mathbf{x} be the vector $(1, 1, \dots, 1)$; note it is fixed by $G(L/K)$. Note that

$$\Lambda := L(U_L^T) \oplus (1, 1, \dots, 1)\mathbb{Z}$$

is a full lattice in \mathbb{R}^T . By Proposition 12.12.4(2)–(3), we have

$$h(\Lambda) = h(L(U_L^T))h(\mathbb{Z}) = n \cdot h(L(U_L^T)). \quad (15.7)$$

Consider the lattice $\Lambda' = \mathbb{Z}^T$ in \mathbb{R}^T , where e_v is the vector with 1 in the v position and 0's elsewhere. By Proposition 15.2.5, $h(\Lambda) = h(\Lambda')$. Since G permutes the places above $v \in S$ transitively, we have

$$\begin{aligned} h(\Lambda) &= h(\Lambda') = h\left(\bigoplus_{w \in T} e_w \mathbb{Z}\right) \\ &= h\left(\bigoplus_{v \in S} \bigoplus_{w \mid v} e_w \mathbb{Z}\right) \\ &= \prod_{v \in S} h\left(\bigoplus_{w \mid v} e_w \mathbb{Z}\right) \quad \text{cohomology respects products, Proposition 12.6.7} \\ &= \prod_{v \in S} h(G^v, \mathbb{Z}) \quad \text{by Corollary 12.8.8 to Shapiro's Lemma} \\ &= \prod_{v \in S} |G^v| \quad \text{Proposition 12.12.4(3)} \\ &= \prod_{v \in S} n_v. \end{aligned}$$

Together with (15.6) and (15.7), we get

$$h(U_L^T) = \frac{1}{n} h(\Lambda') = \frac{1}{n} \prod_{v \in S} n_v. \quad \square$$

2.4 Herbrand quotient of \mathbf{C}_L

Lemma 15.2.7. *If L/K is a cyclic extension of number fields of degree n ,*

$$h(\mathbf{C}_L) = n.$$

Proof. Choose a set of places T for L containing the ramified places and satisfying the conditions of Proposition 15.2.2. Enlarge T so it is stable under $G(L/K)$. Using Propositions 15.2.3 and 15.2.6, we have that

$$h(\mathbf{C}_L) = h(L^\times \mathbb{I}_L^T / L^\times) = h(\mathbb{I}_L^T / \mathbb{I}_L^T \cap L^\times) = \frac{h(\mathbb{I}_L^T)}{h(U_L^T)} = \frac{\prod_{v \in S} n_v}{\frac{1}{n} \prod_{v \in S} n_v} = n$$

□

Proof of Theorem 15.2.1. We have

$$|\mathbb{I}_K / K^\times \text{Nm}_{L/K}(\mathbb{I}_L)| = |H_T^0(G, \mathbf{C}_L)| = h(\mathbf{C}_L) |H_T^{-1}(G, \mathbf{C}_L)| \geq n$$

by Lemma 15.2.7. □

2.5 The Frobenius map is surjective

Using the first inequality, we can already prove surjectivity of the Artin map, defined on ideals.

Proposition 15.2.8: Let L/K be a finite abelian extension, and S be a finite set of primes. Define the map

$$\psi_{L/K} : I^S \rightarrow G(L/K)$$

by setting $\psi_{L/K}(\mathfrak{p}) = \text{Frob}_{L/K}(\mathfrak{p})$ for primes $\mathfrak{p} \notin S$ and extending to a group homomorphism. Then $\psi_{L/K}$ is surjective.

Proof. Let $H = \text{im}(\psi_{L/K})$. By compatibility of the Frobenius map, $\text{Frob}_{K^H/K}(\mathfrak{p})$ is the image of $\text{Frob}_{L/K}(\mathfrak{p})$ under the projection $G(L/K) \rightarrow G(K^H/K)$. Hence the map $\psi_{K^H/K} : I^S \rightarrow G(K^H/K)$ is trivial, giving $(K^H)^v = K_v$ for every $v \notin S$, and

$$\mathbb{I}_K^S \subseteq \text{Nm}_{K^H/K} \mathbb{I}_{K^H}.$$

However, $K^\times \mathbb{I}_K^S$ is dense in \mathbb{I}_K by the weak approximation theorem 7.3.4, so $K^\times \mathbb{I}_K^S = K^\times \text{Nm}_{K^H/K} \mathbb{I}_{K^H} = \mathbb{I}_K$. But by the First Inequality 15.2.1,

$$[K^H : K] \leq [\mathbb{I}_K : K^\times \text{Nm}_{K^H/K} \mathbb{I}_{K^H}] = 1.$$

Hence $K^H = K$, i.e. $H = G$. □

3 The second inequality

We give two proofs of the second inequality, an analytic proof and an algebraic proof. The first has the advantage of being short and sweet, while the second has the advantage of staying completely within the algebraic realm, i.e. not requiring knowledge of L -functions.

Theorem 15.3.1 (Second inequality for global class field theory). *For any extension L/K of degree n , and $G = G(L/K)$, we have*

1. $|H_T^0(G, \mathbf{C}_L)|$ and $|H^2(G, \mathbf{C}_L)|$ divide n .
2. (HT90 for ideles) $|H^1(G, \mathbf{C}_L)| = 1$.

In particular,

$$|\mathbb{I}_K/K^\times \text{Nm}_{L/K} \mathbb{I}_L| \leq [L : K].$$

3.1 Analytic approach

We first show the inequality $|\mathbb{I}_K/K^\times \text{Nm}_{L/K} \mathbb{I}_L| \leq [L : K]$.

Proof of inequality. Let \mathfrak{c} be admissible for L/K , i.e. such that $\mathbf{U}_K(1, \mathfrak{c}) \subseteq \text{Nm}_{L/K}(\mathbb{I}_L)$. By Proposition 11.5.9 we know that $\mathbb{I}_K/K^\times \text{Nm}_{L/K} \mathbb{I}_L \cong I_L^\mathfrak{c}/P_K(1, \mathfrak{c}) \text{Nm}_{L/K}(I_L^\mathfrak{c})$. We show that

$$[I_K^\mathfrak{c} : P_K(1, \mathfrak{c}) \text{Nm}_{L/K}(I_L^\mathfrak{c})] \leq [L : K].$$

Let $H = P_K(1, \mathfrak{c}) \text{Nm}_{L/K} I_L^\mathfrak{c}$ and let χ be a nontrivial character of $I_K^\mathfrak{c}/H$, viewed as a character of $I_K^\mathfrak{c}/P_K(1, \mathfrak{c})$.

Define the Hecke L -series $L_\mathfrak{c}(s, \chi)$ by

$$L_\mathfrak{c}(s, \chi) := \prod_{\mathfrak{p} \nmid \mathfrak{c}} \frac{1}{1 - \frac{\chi(\mathfrak{p})}{\mathfrak{N}\mathfrak{p}^s}} = \sum_{\mathfrak{a} \perp \mathfrak{c}} \frac{\chi(\mathfrak{a})}{\mathfrak{N}\mathfrak{a}^s},$$

where equality follows from expanding the product. Define

$$m(\chi) := \text{ord}_{s=1} L_\mathfrak{c}(s, \chi).$$

Since $L_\mathfrak{c}(s, \chi) = (s - 1)^{m(\chi)} g(s, \chi)$ for some $g(s, \chi)$ nonzero at $s = 1$, taking logs gives

$$\ln L_\mathfrak{c}(s, \chi) \sim m(\chi) \ln(s - 1) = -m(\chi) \ln \frac{1}{s - 1}.$$

Taking the sum over all characters of $I_K^{S(m)}$ gives

$$\ln \zeta_K(s) + \sum_{\chi \neq 1} \ln L_m(s, \chi) \sim \left[1 - \sum_{\chi \neq 1} m(\chi) \right] \ln \frac{1}{s - 1} \tag{15.8}$$

where we use the fact that $\zeta_K(s) := L(s, 1)$ has a pole at $s = 1$.

On the other hand, by the Taylor series expansion for \ln ,

$$\ln L_c(s, \chi) = - \sum_{\mathfrak{p} \nmid c} \ln \left(1 - \frac{\chi(\mathfrak{p})}{\mathfrak{N}\mathfrak{p}^s} \right) = \sum_{n=1}^{\infty} \sum_{\mathfrak{p} \nmid c} \frac{\chi(\mathfrak{p})^n}{n \mathfrak{N}\mathfrak{p}^{ns}} \sim \sum_{\mathfrak{p}} \frac{\chi(\mathfrak{p})}{\mathfrak{N}\mathfrak{p}^s} = \sum_{\mathfrak{K} \in I^c/H} \chi(\mathfrak{K}) \sum_{\mathfrak{p} \in \mathfrak{K}, \mathfrak{p} \nmid c} \frac{1}{\mathfrak{N}\mathfrak{p}^s}$$

where in the last step we grouped together the primes based on what they are modulo H . This is greater than the sum if we only include primes with $f(\mathfrak{P}/\mathfrak{p}) = 1$ (\mathfrak{P} in L). Again we are off by at most a constant if we only include primes splitting completely in L , because the ramified primes are at most a finite subset. We can then “unrestrict” to all the primes of L , and be off by at most a constant in a neighborhood of 1, because the other terms are in the form $\frac{1}{p^f s}$ for $f > 1$.

Let $h = [I^{S(m)} : H]$. We get, for $s \rightarrow 1^+$,

$$\begin{aligned} \ln \zeta_K(s) + \sum_{\chi \neq 1} \ln L_m(s, \chi) &\sim \sum_{\chi} \sum_{\mathfrak{K} \in I^c/H} \chi(\mathfrak{K}) \sum_{\mathfrak{p} \in \mathfrak{K}, \mathfrak{p} \nmid m} \frac{1}{\mathfrak{N}\mathfrak{p}^s} \\ &\asymp O(1) + h \sum_{\mathfrak{p} \in \text{Spl}(L/K)} \frac{1}{\mathfrak{N}\mathfrak{p}^s} & \sum_{\chi} \chi(\mathfrak{K}) = \begin{cases} 0, & \mathfrak{K} \neq H \\ h, & \mathfrak{K} = H. \end{cases} \\ &\sim O(1) + \frac{h}{N} \sum_{f(\mathfrak{P})=1} \frac{1}{\mathfrak{N}\mathfrak{P}^s} & N \text{ primes above each } \mathfrak{p} \\ &\sim O(1) + \frac{h}{N} \ln \zeta_L(s) \\ &\sim O(1) + \frac{h}{N} \ln \frac{1}{s-1}. \end{aligned}$$

Combining this with (15.8) gives $m(\chi) = 0$ (since $\frac{h}{N} > 0$) for all $\chi \neq 1$, and $h \leq N$, as needed. \square

3.2 Algebraic approach

The steps are as follows.

1. Carry out some preliminary local computations.
2. Consider the case where L/K is an extension such that $G(L/K) \cong (\mathbb{Z}/n\mathbb{Z})^r$, and K contains the n th roots of unity. Note this is a Kummer extension, so we can characterize it in terms of $L^{\times n} \cap K$. This will make computations easy for us.

We construct an explicit set E with

$$E \subseteq \text{Nm}_{L/K} \mathbb{I}_L \subseteq \mathbb{I}_K.$$

We have $[\mathbb{I}_K : K^{\times} \text{Nm}_{L/K} \mathbb{I}_L] \mid [I_K : K^{\times} E]$, so it suffices to show the latter equals n^r .

3. Show this.
4. This implies the cyclic prime case, and that the cyclic prime case implies the general case.

This section is incomplete; see Cassels-Frohlich [CF69], pg. 180-185.

3.2.1 Local computations

Proposition 15.3.2: Let K be a local field with $|\mu_n \cap K| = m$, i.e. K contains m n th roots of unity. Then

$$[K^\times : K^{\times n}] = \frac{nm}{|n|_v}$$

and

$$[U_K : U_K^n] = \frac{m}{|n|_v}.$$

Proof. There are two methods: appeal to the structure of K^\times or calculate a Herbrand quotient. □

3.2.2 Constructing E

Since L is a Kummer extension we can write it in the form $K(\sqrt[n]{a_1}, \dots, \sqrt[n]{a_r})$. Let S be a set of primes satisfying the following conditions.

1. S contains all infinite places.
2. S contains all divisors of n .
3. $\mathbb{I}_K = K^\times \mathbb{I}_K^S$. (This is possible by Proposition 15.2.2.)
4. S contains all prime factors in the numerator and denominator of all a_i , i.e. the a_i are all S -units.

Define

$$E = \prod_{v \in S} K_V^{\times n} \times \prod_{v \in T} K_v^\times \times \prod_{v \notin S \cup T} U_v.$$

Lemma 15.3.3. $E \subseteq \text{Nm}_{L/K} \mathbb{I}_L$.

We want to calculate $[\mathbb{I}_K : K^\times E]$ but $K^\times E$ is hard to deal with. E however, is not, because to calculate the index of E we can appeal to Proposition 15.3.2. Thus we use the following group theoretic fact.

Proposition 15.3.4: Let $B \subseteq A$ and C be subgroups of a group G . Then

$$[CA : CB][C \cap A : C \cap B] = [A : B].$$

Then

$$[\mathbb{I}_K : K^\times E] = [K^\times \mathbb{I}_K^{\text{SUT}} : K^\times E] = \frac{[\mathbb{I}_K^{\text{SUT}} : E]}{[K^\times \cap \mathbb{I}_K^{\text{SUT}} : K^\times \cap E]}.$$

See Cassels-Frohlich.

3.3 Finishing the proof

Now we prove Theorem 15.3.1.

Proof of Theorem 15.3.1. Step 1: We show the theorem when $[L : K]$ is prime. In this case, both the first and second inequality hold, so

$$|H_T^2(G, \mathbf{C}_L)| = [\mathbb{I}_K : K^\times \text{Nm}_{L/K}(\mathbb{I}_L)] = [L : K].$$

Since $h(\mathbf{C}_L) = n$ by Lemma 15.2.7, we get $|H_T^1(G, \mathbf{C}_L)| = 1$. Finally, note $H_T^2(G, \mathbf{C}_L) = H_T^0(G, \mathbf{C}_L)$ because $G(L/K)$ is cyclic.

Step 2: We show the theorem when $[L : K]$ is a prime power, by induction on the exponent. Suppose $|G| = p^n$. Every p -group has a normal subgroup of index p . Let $H \triangleleft G$ be such a group; it corresponds to $H = G(L/K')$ for some extension K'/K of degree p . The inflation-restriction exact sequence 12.11.10 gives

$$0 \rightarrow \underbrace{H^1(G/H, \mathbf{C}_{K'})}_{=0 \text{ by prime case}} \xrightarrow{\text{Inf}} H^1(G, \mathbf{C}_L) \xrightarrow{\text{Res}} \underbrace{H^1(H, \mathbf{C}_L)}_{=0 \text{ by induction hypothesis}}.$$

Thus $H^1(G, \mathbf{C}_L) = 0$. This shows part 2. Using $H^1(G, \mathbf{C}_L) = 0$, the inflation-restriction exact sequence gives

$$0 \rightarrow \underbrace{H^2(G/H, \mathbf{C}_{K'})}_{\text{order } p} \xrightarrow{\text{Inf}} H^2(G, \mathbf{C}_L) \xrightarrow{\text{Res}} \underbrace{H^2(H, \mathbf{C}_L)}_{\text{order } |p^{n-1}|}$$

by the case for cyclic extensions and the induction hypothesis. This shows $|H^2(G, \mathbf{C}_L)| \mid p^n$. Finally,

$$|H_T^0(G, \mathbf{C}_L)| = [\mathbf{C}_K : \text{Nm}_{L/K} \mathbf{C}_L] = [\mathbf{C}_K : \text{Nm}_{K'/K}(\mathbf{C}_{K'})][\text{Nm}_{K'/K}(\mathbf{C}_{K'}) : \text{Nm}_{L/K}(\mathbf{C}_L)].$$

Now $[\mathbf{C}_K : \text{Nm}_{K'/K}(\mathbf{C}_{K'})] = p$ by the cyclic case, and the surjection $\text{Nm}_{K'/K} : \mathbf{C}_{K'} / \text{Nm}_{L/K'}(\mathbf{C}_L) \twoheadrightarrow \text{Nm}_{K'/K}(\mathbf{C}_{K'}) / \text{Nm}_{L/K}(\mathbf{C}_L)$ and the induction hypothesis gives that the second factor divides p^{n-1} . This finishes the induction step.

Step 3: We show the theorem holds in general, using Corollary 12.11.7: the map

$$\text{Res}^n : H^n(G, M) \rightarrow H^n(G_p, M)$$

is injective on the p -primary component. Using step 2, for $n = 1$, this gives us that $p \nmid |H_T^1(G, \mathbf{C}_L)|$ for any p , i.e. $H_T^1(G, \mathbf{C}_L) = 0$. For $n = 0, 2$, this gives that $v_p(|H^n(G, M)|) \leq v_p(|H^n(G_p, M)|) \leq v_p(G)$, giving part 1. \square

3.4 Local-to-global principle for algebras

The fact that $H^1(G, \mathbf{C}_L) = 0$ also gives the following corollary.

Theorem 15.3.5 (Brauer-Hasse-Noether Theorem). *Let L/K be any Galois extension with Galois group G . Then the map*

$$H^2(G, L^\times) \rightarrow \bigoplus_{v \in V_K} H^2(G^v, L^{v \times})$$

is injective. A central simple algebra over a number field K is split over K iff it is split locally everywhere.

Proof. Taking cohomology of $0 \rightarrow L^\times \rightarrow \mathbb{I}_L \rightarrow \mathbf{C}_L \rightarrow 0$ gives

$$\begin{array}{ccccccc} H^1(G, \mathbf{C}_L) & \longrightarrow & H^2(G, L^\times) & \longrightarrow & H^2(G, \mathbb{I}_L) & \longrightarrow & \cdots \\ \parallel & & \parallel & & \parallel & & \\ 0 & \longrightarrow & \text{Br}_K & \longrightarrow & \bigoplus_{v \in V_K} \text{Br}_{K_v} & & \end{array} \quad (15.9)$$

Here $H^1(G, \mathbf{C}_L) = 0$ directly from HT90 for ideles (Theorem 15.3.1), and equality on the right comes from

$$H^2(G, \mathbb{I}_L) = \bigoplus_{v \in V_K} H^2(L^v/K_v)$$

(Proposition 15.2.4). Brauer group is H^2 by Theorem 13.5.2. Injectivity of the bottom map gives the result.

(We do need to check that in the above diagram, the map $\text{Br}_K \rightarrow \bigoplus_{v \in V_K} \text{Br}_{K_v}$ is exactly the map sending an algebra to its reduction over every local field. This is a matter of tracing the long windy road between Br and H^2 and left to the reader.) \square

4 Proof of the reciprocity law

To construct the Artin map in the local case, we constructed the invariant map $\text{inv}_K : H^2(K^{\text{ur}}/K) \rightarrow \mathbb{Q}/\mathbb{Z}$. Then we used the fact that $H^2(K^{\text{ur}}) = 0$, i.e. every $a \in H^2(K)$ splits in an unramified extension, to conclude that $H^2(K) \cong H^2(K^{\text{ur}}/K)$.

In the global case we will construct the invariant map $\text{inv}_K : H^2(K_c/K, \mathbb{I}_{K_c}) \rightarrow \mathbb{Q}/\mathbb{Z}$, for a certain infinite cyclotomic extension K_c . Then we show $H^2(K_c, \mathbb{I}_{\overline{K}}) = 0$, i.e. every $a \in H^2(K, \mathbb{I}_{\overline{K}})$ splits in this cyclotomic extension, to conclude $H^2(K, \mathbb{I}_{\overline{K}}) \cong H^2(K_c/K, \mathbb{I}_{\overline{K}_c})$.

We construct the global Artin map by taking the product of the local Artin maps:

$$\begin{aligned} \phi_{L/K} &: \mathbb{I}_K \rightarrow G(L/K) \\ \phi_{L/K}(\mathbf{a}) &= \prod_{v \in V_K} \phi_v(a_v). \end{aligned} \quad (15.10)$$

(Only a finite number of the factors—those where L^v/K_v is ramified or $a_v \notin U_v$ —are not equal to the identity.)

We need to show that $K^\times \subseteq \ker \phi_{L/K}$, so that it factors through $\mathbb{I}_K/K^\times \cdot \text{Nm}_{L/K} \mathbb{I}_L$. Consider the following two properties.

- (A) Define the map $\phi_{L/K}$ as in (15.10). The map $\phi_{L/K}$ takes the value 1 on the principal ideles $K^\times \subseteq \mathbb{I}_K$.
- (B) For all $\alpha \in H^2(G(L/K), L^\times) = \text{Br}_{L/K}$,

$$\text{inv}(\alpha) := \sum_{v \in V_K} \text{inv}_v(i(\alpha)) = 0.$$

Note in (B), inv_v is defined as follows.

Definition 15.4.1: Define inv_v as the following composition:

$$\text{inv}_v : H^2(G, \mathbb{I}_L) \xrightarrow{\text{Res}_{G/G_v}} H^2(G_v, \mathbb{I}_L) \xrightarrow{H^2(G_v, p_v)} H^2(G_v, (L^v)^\times) \xrightarrow{\text{inv}} \mathbb{Q}/\mathbb{Z}$$

where $p_v : \mathbb{I}_L \rightarrow (L^v)^\times$ is the projection map. (This looks complicated, but it is just what you think it is.)

We prove (A) for all finite abelian extensions of number fields and (B) for all finite Galois extensions of number fields.

We first show that (A) holds for a special class of extensions, and then use an “unscrewing” argument to show (A) and (B) hold for more general extensions. The plan of attack is as follows.

1. Show (A) holds for $\mathbb{Q}(\zeta_m)/\mathbb{Q}$.
2. Show (A) holds for all cyclotomic extensions.
3. Show that (B) holds for α split by a cyclotomic extension.
4. Every α is split by a cyclic cyclotomic extension, so (B) holds for all $\alpha \in H^2(K, \overline{K}^\times)$.
5. Show that (A) holds for all abelian extensions.

Note that (A) is a statement about $H_T^{-2} \rightarrow H_T^0$ while (B) is a statement about H^2 . We “transfer” the problem from (A) to (B) so that we can apply our characterization of ϕ_v in terms of the local invariant map (Theorem 14.4.9). First, we need an analogue of Theorem 14.4.9 in the global case.

Lemma 15.4.2. *Let $G = G(L/K)$. For all $v \in V_K$ and all $\chi \in H^1(G, \mathbb{Q}/\mathbb{Z}) = \text{Hom}(G, \mathbb{Q}/\mathbb{Z})$, we have $\text{inv}_v(\bar{\mathbf{a}} \cup \delta\chi) = \chi_v(\phi_v(a_v))$. (χ_v is the restriction of χ to G_v and $\bar{\mathbf{a}}$ is the image of \mathbf{a} in $H_T^0(G(L/K), \mathbb{I}_L)$). Hence*

$$\text{inv}(\bar{\mathbf{a}} \cup \delta\chi) = \sum_v \text{inv}_v(\bar{\mathbf{a}} \cup \delta\chi) = \chi(\phi(a)).$$

Proof. Since restriction commutes with cup products (Proposition 12.11.9) and with δ , we have

$$\begin{aligned} \text{inv}_v(\bar{\mathbf{a}} \cup \delta\chi) &= \text{inv}(p_v \text{Res}_{G/G_v}(\bar{\mathbf{a}} \cup \delta\chi)) \\ &= \text{inv}(p_v(\bar{\mathbf{a}}) \cup \delta\chi_v) && \text{Res}_{G/G_v}(\chi) = \chi_v \\ &= \text{inv}(\bar{a}_v \cup \delta\chi_v) = \chi_v(\phi_v(a_v)). \end{aligned}$$

We invoked Theorem 14.4.9 in the last step.

Taking the product gives the second statement:

$$\chi(\phi(\mathbf{a})) = \chi\left(\prod_v \phi_v(a_v)\right) = \sum_v \chi_v(\phi_v(\mathbf{a})) = \sum_v \text{inv}_v(\bar{\mathbf{a}} \cup \delta\chi).$$

□

4.1 (A) holds for $\mathbb{Q}(\zeta_n)/\mathbb{Q}$

Proposition 15.4.3: For any $m \in \mathbb{N}$,

$$\phi_{\mathbb{Q}(\zeta_m)/\mathbb{Q}}(\mathbb{Q}^\times) = 1.$$

First reduce to the case where $m = p$ is prime. We give two approaches.

Proof 1. By Example 11.4.3, we know the ideal version of global class field theory holds for all cyclotomic extensions of \mathbb{Q} . Note the maximal unramified extension of \mathbb{Q}_p is included in $\mathbb{Q}_p(\zeta_\infty)$ for all p (Theorem 8.2.6). Hence by Theorem 11.6.6(2), there is a map ϕ' satisfying the conditions of the idele version of GCFT, except that $\phi'_\mathbb{R}$ may not equal $\phi_\mathbb{R}$. Letting ϕ'_v be the restriction of ϕ' to K_v , we have (on $G(\mathbb{Q}(\zeta_\infty)/\mathbb{Q})$)

$$\phi'(\mathbf{a}) = \phi'_\mathbb{R}(a_\mathbb{R}) \prod_{v \in V_\mathbb{Q}^0} \phi'_v(a_v) \stackrel{?}{=} \prod_{v \in V_\mathbb{Q}} \phi_v(a_v) = \phi'(\mathbf{a}) \tag{15.11}$$

where the middle inequality is pending a proof that $\phi'_\mathbb{R} = \phi_\mathbb{R}$. We check this is true.

Since $\phi'_\mathbb{R}$ is a map $\mathbb{R}/\mathbb{R}_{>0} \rightarrow G(\mathbb{C}/\mathbb{R})$, it suffices to show complex conjugation is in the image of ϕ' . We have $G(\mathbb{C}/\mathbb{R}) \cong G(\mathbb{R}(i)/\mathbb{R})$, so consider ϕ' on $G(\mathbb{Q}(i)/\mathbb{Q})$. As $\phi'(\mathbb{Q}^\times) = 1$, we have by (15.11) and the fact that $\mathbb{Q}(i)/\mathbb{Q}$ is only ramified at 2 that on $G(\mathbb{Q}(i)/\mathbb{Q})$,

$$1 = \phi'(-7) = \phi'_2(-7)\phi'_7(-7)\phi'_\mathbb{R}(-7)$$

Now $-7 \equiv 1 \pmod{8}$ so $-7 \in \text{Nm}_{\mathbb{Q}_2(i)/\mathbb{Q}_2}(\mathbb{Q}_2(i)^\times)$, and $\phi'_2(-7) = 1$. We have $v_7(-7) = 1$, so $\phi'_7(-7)$ equals the Frobenius element, complex conjugation. Hence $\phi'_\mathbb{R}(-7)$ is also complex conjugation.

Thus (15.11) holds, and we have $\phi_{\mathbb{Q}(\zeta_\infty)/\mathbb{Q}}(\mathbb{Q}^\times) = \phi'_{\mathbb{Q}(\zeta_\infty)/\mathbb{Q}}(\mathbb{Q}^\times) = 1$, as needed. □

Proof 2. Use explicit computations of local symbols, obtained from Lubin-Tate theory. See Cassels-Frohlich [CF69], p. 191. □

4.2 (B) holds for all cyclomic extensions

We prove the following more general proposition.

Proposition 15.4.4 (Devissage): ⁵ If (A) is true for L/K , then (A) holds for

1. any subextension M/K and
2. any extension LK'/K' .

For an extension $K'(\zeta_n)/K'$, apply the proposition with $L = \mathbb{Q}(\zeta_n)$ and $K = \mathbb{Q}$ to obtain the following.

Corollary 15.4.5. (A) holds for all cyclotomic extensions.

Proof of Proposition 15.4.4.

1. For any place v , ϕ_{M^v/K_v} is the composition of ϕ_{L^v/K_v} and the projection $G(L^v/K_v) \rightarrow G(M^v/K_v)$. Since the global map is the product of the local maps, $\phi_{M/K}$ is the composition of $\phi_{L/K}$ and $G(L/K) \rightarrow G(M/K)$. Hence $\phi_{M/K}(K^\times) = 1$.
2. Let $L' = L \cdot K'$. We have a natural inclusion $G(L'/K') \hookrightarrow G(L/K)$. The local Artin map is compatible with basefield extension with respect to the norm map. Since the norm on ideles is computed componentwise, it follows the map $\phi = \prod_{v \in V_K} \phi_v$ is also compatible with field extensions.

$$\begin{array}{ccc} \mathbb{I}_{K'} & \xrightarrow{\phi_{L'/K'}} & G(L'/K') \\ \downarrow \text{Nm}_{K'/K} & & \downarrow i \\ \mathbb{I}_K & \xrightarrow{\phi_{L/K}} & G(L/K). \end{array}$$

Suppose $a \in K'^\times$. By commutativity and (A) for the extension L/K , we have

$$i \circ \phi_{L'/K'}(a) = \phi_{L/K}[\underbrace{\text{Nm}_{K'/K}(a)}_{\in K^\times}] = 1.$$

Since i is injective, this implies $\phi_{K'/K'}(a) = 1$. □

4.3 (A) for cyclotomic implies (B) for α split by cyclic cyclotomic

This follows from the more general proposition:

Proposition 15.4.6: If L/K is cyclic, then (A) implies (B).

⁵Devissage means “unscrewing” in French.

Proof. Since L/K is cyclic, we can take $\chi \in H^1(G, \mathbb{Q}/\mathbb{Z})$ to be a generating character. We have the following commutative diagram.

$$\begin{array}{ccccc}
 K^\times & \hookrightarrow & \mathbb{I}_K & \xrightarrow{\phi_{L/K}} & G(L/K) \\
 \downarrow \bullet \cup \delta\chi & & \downarrow \bullet \cup \delta\chi & & \downarrow \chi \\
 H^2(G, L^\times) & \longrightarrow & H^2(G, \mathbb{I}_L) & \xrightarrow{\text{inv}} & \mathbb{Q}/\mathbb{Z}
 \end{array}$$

The left-hand square commutes by functoriality of cup products; the right-hand square commutes by Lemma 15.4.2. Recall $\bullet \cup \delta\chi$ is an isomorphism for G cyclic, by Proposition 12.12.1. Hence if $a \in H^2(G, L^\times)$, then it is equal to $b \cup \delta\chi$ for some $b \in K^\times$, and

$$\text{inv}(a) = \text{inv}(b \cup \delta\chi) = \chi(\phi_{L/K}(b)) = 0.$$

In the last step we use (A) to give $\phi_{L/K}(b) = 0$. □

4.4 (B) for cyclic cyclotomic implies (B) in general

It suffices to prove the following.

Theorem 15.4.7. *For any $\beta \in H^2(K)$ there exists a cyclic cyclotomic extension L/K such that β maps to 0 in $H^2(L)$.*

There exists a cyclotomic extension $K_c \subseteq K(\zeta_\infty)$ with $G(K_c/K) \cong \widehat{\mathbb{Z}}$ such that the inclusion map

$$H^2(K_c/K) \rightarrow H^2(K)$$

is an isomorphism.

We first give a criterion for β to map to 0 in $H^2(L)$, then find a cyclotomic L/K where this criterion holds.

Lemma 15.4.8. *Let $\alpha \in H^2(K)$. Then $\text{Res}_{K/L}(\alpha) = 0$ in $H^2(L)$ if and only if $[L^v : K_v] \text{inv}_v(\alpha) = 0$ for every $v \in V_K$.*

Proof. By the Brauer-Hasse-Noether Theorem 15.3.5, $\text{Res}_{K/L}(\alpha) = 0$ in $H^2(L/K, L^\times)$ iff $\text{Res}_{K/L}(\alpha) = 0$ in $H^2(L^v/K_v, L^{v\times}) = 0$ for all v . Since inv_{K_v} is an isomorphism, this is true iff $\text{inv}_{K_v} \text{Res}_{K_v/L^v}(\alpha) = 0$ for all v . But we know

$$\text{inv}_{K_v} \text{Res}_{K_v/L^v}(\alpha) = [L^v : K_v] \text{inv}_v(\alpha),$$

from the class formation for LCFT (Theorem 14.4.14). □

Lemma 15.4.9. *Suppose K/\mathbb{Q} is a finite extension and S be a finite set of places of K . There exists a cyclic cyclotomic extension L/K such that*

$$\begin{aligned}
 m & \mid [L^v : K_v] \text{ for every finite } v \in S \\
 2 & \mid [L^v : K_v] \text{ for every real } v \in S.
 \end{aligned}$$

(The second condition is just equivalent to L being complex.)

Proof. First consider the case $K = \mathbb{Q}$. Note that for an odd prime q ,

$$G(\mathbb{Q}(\zeta_{q^r})/\mathbb{Q}) \cong (\mathbb{Z}/q^r)^\times \cong \mathbb{Z}/(q-1)q^{r-1} \cong \mathbb{Z}/(q-1) \times \mathbb{Z}/q^{r-1}.$$

Let $L(q^r)$ be the subextension of $\mathbb{Q}(\zeta_{q^r})$ with Galois group \mathbb{Z}/q^{r-1} . Because \mathbb{Q}_p only has a finite number of roots of unity, $v_q([L(q^r) : \mathbb{Q}_p]) \rightarrow \infty$ as $r \rightarrow \infty$.

Similarly for $q = 2$,

$$G(\mathbb{Q}(\zeta_{2^r})/\mathbb{Q}) \cong (\mathbb{Z}/2^r)^\times \cong \mathbb{Z}/2 \times \mathbb{Z}/2^{r-2}.$$

The subextension $\mathbb{Q}(\zeta_{2^r} - \zeta_{2^r}^{-1})$ corresponds to the automorphisms $\zeta \mapsto \zeta^s$ with $s \equiv 1 \pmod{4}$, which form a group isomorphic to $\mathbb{Z}/2^{r-2}$. Let $L(2^r) = \mathbb{Q}(\zeta_{2^r} - \zeta_{2^r}^{-1})$ (note this is complex), then similarly $\lim_{r \rightarrow \infty} v_2([L(2^r) : \mathbb{Q}]) = \infty$. Now take

$$L := \prod_{q_i | 2m} L(q_i^{r_i})$$

for r_i large enough. As it is a compositum of cyclic cyclotomic extensions of relatively prime degrees, L is cyclic cyclotomic.

Now suppose we are given general K . First construct $\mathbb{Q}(\zeta_n)/\mathbb{Q}$ satisfying the conditions for \mathbb{Q} with $m[K : \mathbb{Q}]$. Then take $L = K(\zeta_n)$. We have $[K^v(\zeta_n) : K^v] \mid m$ for finite primes v of \mathbb{Q} since $[K^v : \mathbb{Q}_v] \mid [K : \mathbb{Q}]$.

We can take $K_c = \cup_{S, r_i} K \cdot \prod_{q_i \in S} L(q_i^{r_i})$. □

Proof of Theorem 15.4.7. We know $\text{inv}_v(\alpha) = 0$ except for a finite number of primes, say primes in S . Suppose $m \text{inv}_v(\alpha) = 0$. Use Lemma 15.4.9 to get $L = K(\zeta_N)$ such that works for m and S . Then by Lemma 15.4.8, $\text{Res}_{K/L}(\alpha) = 0$ in $H^2(K(\zeta_N))$. □

4.5 (B) implies (A) for all abelian extensions

This will follow from the following proposition.

Proposition 15.4.10: If L/K is abelian, then (B) for L/K implies (A) for L/K .

Proof. Let $a \in K^\times$. By Lemma 15.4.2, for any character χ ,

$$\chi(\phi_{L/K}(a)) = \text{inv}\left(\underbrace{\bar{a} \cup \delta\chi}_{\in H^2(L/K, L^\times)}\right) = 0.$$

Hence $\phi_{L/K}(a) = 0$. □

We have now proved the following.

Theorem 15.4.11. *The following hold.*

(A) For an abelian extension L/K , define the map $\phi_{L/K}$ as in (15.10). The map $\phi_{L/K}$ takes the value 1 on the principal ideles $K^\times \subseteq \mathbb{I}_K$.

(B) For any $\alpha \in H^2(\overline{K}/K)$,

$$\text{inv}(\alpha) := \sum_{v \in V_K} \text{inv}_v(\alpha) = 0.$$

5 The ideles are a class formation

We now complete the proof of global class field theory by showing that the ideles are a *class formation* and invoking the theorems in Section 4. In the local case, the G -modules in the class formations are the fields themselves, but in the global case, the G -modules are the ideles.

Theorem 15.5.1. *Let K be a global field. Then*

$$(G(\overline{K}/K), \{G(L/K) : L/K \text{ finite Galois}\}, \mathbf{C}_{\overline{K}})$$

is a class formation.

Note that $\mathbf{C}_{\overline{K}}^{G(\overline{K}/L)} = \mathbf{C}_L$ for each L by Proposition 15.1.3.

Proof. We check the axioms in Definition 14.4.5.

Step 1: First, $H^1(G(L/K), \mathbf{C}_L) = 0$ for every cyclic extension of prime degree (in fact every finite extension), by Theorem 15.3.1.

Second, we need maps $\text{inv}_{L/K} : H^2(L/K, \mathbf{C}_K) \xrightarrow{\cong} \mathbb{Q}/\mathbb{Z}$. Right now we just have a map

$$\text{inv}_{L/K} : H^2(G(L/K), \mathbb{I}_L) \rightarrow \mathbb{Q}/\mathbb{Z}.$$

We need to show $\text{inv}_{L/K}$ “factors through” $H^2(G(L/K), \mathbf{C}_L)$. We also need to show compatibility with inflation and restriction, and that

$$\text{inv}_{L/K} : H^2(G(L/K), \mathbf{C}_L) \xrightarrow{\cong} \frac{1}{[L : K]} \mathbb{Z}/\mathbb{Z}$$

for all L/K . It is hard to show this directly, except in the cyclic case, when we know the first inequality holds. As we will see, though, showing the cyclic case is enough, because by Theorem 15.4.7, every element of $H^2(G(\overline{K}/K), \mathbf{C}_{\overline{K}})$ is contained in $H^2(G(L/K), \mathbf{C}_L)$ for some cyclic (in fact, also cyclotomic) L/K .

Step 2: Consider the following commutative diagram, whose columns are inflation-restriction sequences.

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & H^2(L/K, L^\times) & \xrightarrow{i_1} & H^2(L/K, \mathbb{I}_L) & \xrightarrow{p_1} & H^2(L/K, \mathbf{C}_L) \\
 & & \downarrow \text{Inf} & & \downarrow \text{Inf} & & \downarrow \text{Inf} \\
 0 & \longrightarrow & H^2(M/K, M^\times) & \xrightarrow{i_2} & H^2(M/K, \mathbb{I}_M) & \xrightarrow{p_2} & H^2(M/K, \mathbf{C}_M) & \xrightarrow{\text{inv}} & \frac{1}{n}\mathbb{Z}/\mathbb{Z} \\
 & & \downarrow \text{Res} & & \downarrow \text{Res} & & \downarrow \text{Res} & & \downarrow \text{Res} \\
 0 & \longrightarrow & H^2(M/L, M^\times) & \xrightarrow{i_3} & H^2(M/L, \mathbb{I}_M) & \xrightarrow{p_3} & H^2(M/L, \mathbf{C}_M) & \xrightarrow{\text{inv}} & \mathbb{Q}/\mathbb{Z} \\
 & & & & & & & & \downarrow \text{inv} \\
 & & & & & & & & \mathbb{Q}/\mathbb{Z}.
 \end{array}$$

The columns are exact by the inflation-restriction exact sequence (Proposition 12.11.10) and the following:

1. $H^1(M/L, M^\times) = 0$ by Hilbert's Theorem 90 (Theorem 13.1.1).
2. $H^1(M/L, \mathbb{I}_M) = 0$ by Proposition 15.2.4.
3. $H^1(M/L, \mathbf{C}_M) = 0$ by Theorem 15.3.1.

The rows are exact because they come from the long exact sequences of $0 \rightarrow L^\times \rightarrow \mathbb{I}_L \rightarrow \mathbf{C}_L \rightarrow 0$ and $0 \rightarrow M^\times \rightarrow \mathbb{I}_M \rightarrow \mathbf{C}_M \rightarrow 0$, and the fact that H^1 of $\mathbf{C}_L, \mathbf{C}_M$ is trivial (again by Theorem 15.3.1).

Step 3: Next we show the maps inv are compatible with inflation. Indeed, since we have a class formation for local class field theory (Theorem 14.4.14), for every $w \mid v$ we have the diagram

$$\begin{array}{ccc}
 H^2(L_w/K_v) & \xrightarrow{\text{inv}_{K_v}} & \frac{1}{[L_w:K_v]}\mathbb{Z}/\mathbb{Z} \\
 \downarrow \text{Inf}_{L_w/L_v} & & \downarrow i \\
 H^2(K_v) & \xrightarrow{\text{inv}_{K_v}} & \mathbb{Q}/\mathbb{Z}.
 \end{array}$$

Now $H^2(G(L/K), \mathbb{I}_K) \cong \bigoplus_{v \in V_K} H^2(G^v, (L^v)^\times)$ by Proposition 15.2.4 and $\text{inv} = \sum_{v \in V_K} \text{inv}_v$, so inv is compatible with inflation.

Step 4: Thus, we can take the direct limit over M , noting direct limits preserve exactness,

to get (we will explain the dashed and dotted lines)

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 & & (15.12) \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & H^2(L/K, L^\times) & \xrightarrow{i_1} & H^2(L/K, \mathbb{I}_L) & \xrightarrow{p_1} & H^2(L/K, \mathbf{C}_L) & & \\
 & & \downarrow \text{Inf} & & \downarrow \text{Inf} & & \downarrow \text{Inf} & & \\
 0 & \longrightarrow & H^2(K, \overline{K}^\times) & \xrightarrow{i_2} & H^2(K, \mathbb{I}_{\overline{K}}) & \xrightarrow{p_2} & H^2(K, \mathbf{C}_{\overline{K}}) & \xrightarrow{\text{inv}'_1} & \frac{1}{n}\mathbb{Z}/\mathbb{Z} \\
 & & \downarrow \text{Res} & & \downarrow \text{Res} & & \downarrow \text{Res} & & \downarrow \text{inv}'_2 \\
 0 & \longrightarrow & H^2(L, \overline{L}^\times) & \xrightarrow{i_3} & H^2(L, \mathbb{I}_{\overline{K}}) & \xrightarrow{p_3} & H^2(L, \mathbf{C}_{\overline{K}}) & \xrightarrow{\text{inv}'_3} & \mathbb{Q}/\mathbb{Z} \\
 & & & & \searrow \text{inv}_3 & & & & \downarrow n \\
 & & & & & & & & \mathbb{Q}/\mathbb{Z}
 \end{array}$$

Step 5: Now we show the maps inv_j are compatible under restriction. Again, since we have a class formation for local class field theory (Theorem 14.4.14), we have the diagram

$$\begin{array}{ccc}
 H^2(K_v) & \xrightarrow{\text{inv}_{K_v}} & \mathbb{Q}/\mathbb{Z} \\
 \downarrow \text{Res}_{K_v/L_w} & & \downarrow [L_w:K_v] \\
 H^2(L_w) & \xrightarrow{\text{inv}_{L_w}} & \mathbb{Q}/\mathbb{Z}
 \end{array}$$

Using $H^2(G(L/K), \mathbb{I}_K) \cong \bigoplus_{v \in V_K} H^2(G^v, (L^v)^\times)$, we can write an element of $H^2(K, \mathbb{I}_K)$ as $\mathbf{x} = (x_v)_{v \in V_K}$, where $x_v \in H^2(G^v, (K_v)^\times)$. On degree 0, $\text{Res}_{K/L}$ is the diagonal imbedding $\mathbb{I}_K \xrightarrow{\cong} \mathbb{I}_L^G \hookrightarrow \mathbb{I}_L$ of Proposition 15.1.3, so on degree 2,

$$\text{Res}_{K/L} \mathbf{x} = \left((\text{Res}_{K_v/L_w} x_v)_{w|v} \right)_{v \in V_K} \in \bigoplus_{v \in V_K} \bigoplus_{w|v} H^2(G_w, \overline{K}_v^\times).$$

The invariant map then sends this to

$$\sum_{v \in V_K} \sum_{w|v} \text{inv}_{L_w} (\text{Res}_{K_v/L_w} x_v) = \sum_{v \in V_K} \sum_{w|v} n_{w/v} \text{inv}_{K_v} x_v = n \sum_{v \in V_K} \text{inv}_{K_v} x_v = n \text{inv}_K \mathbf{x},$$

using the fact that $[L : K] = \sum_{w|v} n_{w/v}$, where $n_{w/v}$ is the local degree.

Step 6: By Theorem 15.4.11, the bent maps are complexes, i.e. $\text{im}(i_j) \subseteq \ker(\text{inv}_j)$ for all three rows.

Thus the maps inv_j factor through the images $\text{im}(p_j)$, for $j = 1, 2, 3$ to give the maps inv'_j . Be careful: we have only so far defined $\text{inv}'_1 : \text{im}(p_1) \rightarrow \frac{1}{n}\mathbb{Z}/\mathbb{Z}$, and not $\text{inv}'_1 : H^2(L/K, \mathbf{C}_L) \rightarrow \frac{1}{n}\mathbb{Z}/\mathbb{Z}$. We want to show that for certain extensions L/K , the p_j are in fact surjective, so the map inv'_j is an isomorphism $H^2(L/K, \mathbf{C}_L) \rightarrow \frac{1}{n}\mathbb{Z}/\mathbb{Z}$.

To do this we orders of $\text{im}(\text{inv}_1)$ and $|H^2(L/K, \mathbf{C}_L)|$. Again, we use $H^2(L/K, \mathbb{I}_L) \cong \bigoplus_{v \in V_K} \frac{1}{n_v} \mathbb{Z}/\mathbb{Z}$ (Proposition 15.2.4). Making this identification using the invariant maps inv_v , the invariant map takes $(a_v)_v \in \bigoplus_{v \in V_K} \frac{1}{n_v} \mathbb{Z}/\mathbb{Z}$ to $\sum_{v \in V_K} a_v$. Thus $\text{im}(\text{inv}_1) = \frac{1}{\text{lcm}_v(n_v)} \mathbb{Z}/\mathbb{Z}$ and

$$|\text{im}(\text{inv}_1)| = \text{lcm}_v(n_v).$$

We have that

$$\text{lcm}_v(n_v) = |\text{im}(\text{inv}_1)| = |\text{im}(\text{inv}'_1)| \leq |\text{im}(p_1)| \leq |H^2(L/K, \mathbf{C}_L)| \leq n, \quad (15.13)$$

where the last step is the second inequality. We don't get any information out of this unless $\text{lcm}_v(n_v) = n$. For certain extensions L/K , we do know it is true, though.

Step 7: We show that if L/K is cyclic, then $\text{lcm}_v(n_v) = n$. Let S be the set of ramified primes and infinite places of K . By Proposition 15.2.8, $G(L/K)$ is generated by the elements $\text{Frob}_{L/K}(\mathfrak{p})$ for $\mathfrak{p} \notin S$. Now $\langle \text{Frob}_{L/K}(\mathfrak{p}) \rangle$ is sent to a subgroup of index n_v in $G(L/K)$. Since $G(L/K)$ to be generated by these elements, we must have $\text{lcm}_v(n_v) = n$.

Then equality holds everywhere in (15.13), we have the exact sequence

$$0 \rightarrow H^2(L/K, L^\times) \rightarrow H^2(L/K, \mathbb{I}_L) \rightarrow H^2(L/K, \mathbf{C}_L) \cong \frac{1}{n} \mathbb{Z}/\mathbb{Z} \rightarrow 0,$$

where the map $H^2(L/K, \mathbb{I}_L) \rightarrow \frac{1}{n} \mathbb{Z}/\mathbb{Z}$ is the invariant map.

Step 8: Taking the direct limit over all $L \subseteq K_c$ (as defined in Theorem 15.4.7) we get

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^2(K, K_c^\times) & \longrightarrow & H^2(K_c/K, \mathbb{I}_{K_c}) & \longrightarrow & H^2(K_c/K, \mathbf{C}_{K_c}) \cong \mathbb{Q}/\mathbb{Z} \longrightarrow 0 \\ & & \downarrow \text{Inf} & & \downarrow \text{Inf} & & \downarrow \text{Inf} \\ 0 & \longrightarrow & H^2(K) & \longrightarrow & H^2(K, \mathbb{I}_{\overline{K}}) & \xrightarrow{\text{inv}} & \mathbb{Q}/\mathbb{Z} \end{array}$$

where the top row is exact. By Theorem 15.4.7, the left vertical map is an isomorphism. The middle map is also an isomorphism because Theorem 15.2.4 gives that it is the map

$$\bigoplus_{v \in V_K} H^2(K_c^v/K_v) \rightarrow \bigoplus_{v \in V_K} H^2(K_v).$$

This is surjective because $H^2(K_c^v/K_v) \cong \mathbb{Q}/\mathbb{Z}$ via the invariant map, K_c^v being the directed union of L_w with $[L_w : K_v]$ arbitrarily divisible. Hence it is an isomorphism. Finally, the right vertical map is clearly an isomorphism. Thus the bottom row is short exact and inv_K gives an isomorphism $H^2(K, \mathbf{C}_{\overline{K}}) \rightarrow \mathbb{Q}/\mathbb{Z}$, i.e. the map inv'_2 in (15.12). Restricting to $H^2(L'/K, \mathbf{C}_{L'})$, it is an isomorphism to $\frac{1}{[L':K]} \mathbb{Z}/\mathbb{Z}$ for any L' , as needed. \square

We are now ready to reap the rewards of our hard work.

Theorem (Global reciprocity, Theorem 11.6.1). *Given a finite abelian extension L/K , there is a unique continuous homomorphism $\phi_{L/K}$ that is compatible with the local Artin maps, i.e. the following diagram commutes:*

$$\begin{array}{ccc} \mathbb{I}_K & \xrightarrow{\phi_{L/K}} & G(L/K) \\ i_v \uparrow & & \uparrow \\ K_v^\times & \xrightarrow{\phi_v} & G(L^v/K_v). \end{array}$$

Moreover, $\phi_{L/K}$ satisfies the following properties.

1. (Isomorphism) For every finite abelian extension L/K , ϕ_K defines an isomorphism

$$\phi_{L/K} : \mathbf{C}_K / \text{Nm}_{L/K}(\mathbf{C}_L) = \mathbb{I}_K / (K^\times \cdot \text{Nm}_{L/K}(\mathbb{I}_L)) \xrightarrow{\cong} G(L/K).$$

2. (Compatibility over all extensions) For $L \subseteq M$, L, M both finite abelian extensions of K , the following commutes:

$$\begin{array}{ccc} & & G(M/K) \\ & \nearrow \phi_{M/K} & \downarrow p_L \\ \mathbb{I}_K & \xrightarrow{\phi_{L/K}} & G(L/K) \end{array}$$

Thus we can define $\phi_K := \varprojlim_{L/K \text{ abelian}} \phi_{L/K}$ as a map $\mathbb{I}_K \rightarrow G(K^{ab}/K)$.

3. (Compatibility with norm map) ϕ_K is a continuous homomorphism $\mathbb{I}_K \rightarrow G(K^{ab}/K)$, and the following commutes.

$$\begin{array}{ccc} \mathbb{I}_L & \xrightarrow{\phi_L} & G(L^{ab}/L) \\ \downarrow \text{Nm}_{L/K} & & \downarrow \bullet|_{K^{ab}} \\ \mathbb{I}_K & \xrightarrow{\phi_K} & G(K^{ab}/K) \end{array}$$

Proof. By Theorem 15.5.1 and the abstract reciprocity law (Theorem 14.4.8) we get isomorphisms $\phi'_{L/K} : \mathbf{C}_K / \text{Nm}_{L/K} \mathbf{C}_L \rightarrow G(L/K)$ satisfying the required compatibility properties. We only have to check that $\phi'_{L/K} = \phi_{L/K}$ (recall we defined $\phi_{L/K}$ as the product of local maps). From Theorem 14.4.9, for every character χ , $\chi(\phi'_{L/K}(\mathbf{a})) = \text{inv}_K(\bar{\mathbf{a}} \cup \delta\chi)$. But this is also true for $\phi_{L/K}$ by Proposition 15.4.2. Hence $\phi_{L/K} = \phi'_{L/K}$, as needed.

Uniqueness is clear from the condition that ϕ restricts to the local Artin maps. □

6 Existence theorem

We now prove the existence theorem for global class field theory.

Proof of Theorem 11.6.3 and Theorem 11.6.4. This involves explicitly constructing norm groups and calculating norm indices, which overlaps with Section 3.2. The proof is omitted for now. See Cassels-Frohlich [CF69], pg. 201-202.

Theorem 11.6.4 now follows from the Existence Theorem and Theorem 14.4.13. \square

Finally, we prove that ϕ_K gives a topological isomorphism $\mathbb{I}_K/\overline{K^\times(K_\infty^\times)^0} \rightarrow G(K^{\text{ab}}/K)$. This finishes the proof of all theorems of global class field theory.

Proof of Theorem 11.6.5. First we prove that ϕ_K is surjective. We know that $\phi_{H_K/K} : \mathbb{I}_K \rightarrow G(H_K/K)$ is surjective, where H_K is the Hilbert class field (See Definition 5), since this is a finite extensions. Thus it suffices to show $\phi_{H_K/K} : \mathbb{I}_K \rightarrow G(K^{\text{ab}}/H_K)$ is surjective.

We know that for each place v of K , $\phi_K : K_v \twoheadrightarrow W(K_v^{\text{ab}}/K_v)$ is surjective (Theorem 11.2.4). Restricting to U_v , we get that $\phi_K|_{U_v} : U_v \twoheadrightarrow I(K_v^{\text{ab}}/K_v) \cong I_v(K^{\text{ab}}/K)$ is surjective. Since $K^\times (\prod_{v \in V_K} U_v) / K^\times \subseteq \mathbb{I}_K$, it suffices to show $\prod_{v \in V_K} I_v(K^{\text{ab}}/K) = G(K^{\text{ab}}/K)$. Let $K_v^{\text{ab,ur}}$ denote the maximal abelian extension of K unramified at v . We have by Theorem 2.7.2 that

$$\prod_{v \in V_K} I_v(K^{\text{ab}}/K) = \prod_{v \in V_K} G(K^{\text{ab}}/K_v^{\text{ab,ur}}) = G\left(K^{\text{ab}} / \bigcap_{v \in V_K} K_v^{\text{ab,ur}}\right) = G(K/H_K)$$

since H_K is the maximal abelian extension unramified at all places. This shows surjectivity.

To show the kernel is $\overline{K^\times(K_\infty^\times)^0}$, note that this is exactly the intersection of all open subsets of finite index in \mathbb{I}_K . \square

Chapter 16

Applications

In this chapter we give several important applications of class field theory to number theory, rewarding the reader for reading the difficult proofs in the last few chapters (or conversely, motivating the reader to read the proofs).

Why is class field theory useful? It relates a field K to its Galois group $G(K^{\text{ab}}/K)$, so transfers information about the extensions of a field into information *contained in the field itself*, or conversely, relates the behavior of elements in the field K , to their behavior in various extension fields. Moreover, because the global Artin map is constructed from the local Artin maps, questions in number theory involving global fields like \mathbb{Q} can be understood by patching together information from its completions (local fields). In the chapter, we will use the full power of class field theory to give solutions to the following problems.

Throughout, we will assume that K is a number field.

1. **Reciprocity laws:** We show, roughly, that whether a prime p is a perfect n th power modulo q , depends only on $q \bmod p$ (actually, some multiple of p). Reciprocity hence shows that the Legendre symbol $\left(\frac{*}{\bullet}\right)$, is like a group homomorphism in *both* the top and bottom. The Artin isomorphism will give us the homomorphism in the bottom.
2. **Local-to-global principle:** We show the **Hasse-Minkowski theorem**: a quadratic form has a solution in K iff it has a solution in every completion of K .
3. **Density of primes:** We prove the **Chebotarev density theorem** on the distribution of prime ideals in a number field.
4. **Splitting of primes:** We show how a prime \mathfrak{p} splits in an abelian extension L/K depends only on \mathfrak{p} modulo a *ray class group*, since splitting behavior can be expressed in terms of the Artin map (Proposition 11.1.3). We show this characterization is unique to abelian extensions, and give some examples for splitting in nonabelian extensions.
5. **Maximal unramified abelian extension:** We characterize the maximal unramified abelian extension H_K of a number field K , and show that all ideals of K become principal in H_K . H_K can be computed for quadratic extensions using the modular function j , which we show in Chapter 17.

6. **Primes represented by quadratic forms:** We relate quadratic forms to primes using the Gauss correspondence (Theorem 4.5.1), then use the Hilbert class field to characterize which primes are represented by a given quadratic form.
7. **Artin and Hecke L -functions:** We use class field theory to show that for abelian extensions, all Artin L -functions are Hecke L -functions. This is useful because it is relatively easy to show Hecke L -functions satisfy nice properties such as analytic continuation and functional equation. This was Emil Artin's original motivation for class field theory.

Finally, we describe how class field theory fits as the “1-dimensional case” of the Langlands program.

1 Reciprocity laws

First we interpret and generalize the Legendre symbol using class field theory. We derive a generalize reciprocity law using class field theory, and then specialize to quadratic, cubic, and biquadratic reciprocity.

Reciprocity laws take two forms. The first is as follows.

Theorem 16.1.1 (Weak reciprocity). *Let K be a number field containing all n th roots of unity. Let p be a fixed prime. Then there exists a modulus \mathfrak{m} and a finite subset $S \in I_K^{\mathfrak{m}}/P_{K,1}(\mathfrak{m})$, such that for all p relatively prime to \mathfrak{m} ,*

$$p \text{ is a perfect } n\text{th power mod } q \iff (q \bmod P_{K,1}(\mathfrak{m})) \in S.$$

In fact, S is the kernel of a certain homomorphism $I_K(\mathfrak{m})/P_{K,1}(\mathfrak{m}) \rightarrow \mu_n$.

This tells us that whether p is a perfect n th power modulo q , depends only on the modular properties of q , and is moreover characterized by a group homomorphism. However, it does not give an efficient method to actually determine whether p is a perfect n th power modulo q . To get this we turn to strong reciprocity.

We know that the Legendre symbol $\left(\frac{\bullet}{p}\right)$ (and its generalizations to n th powers, $\left(\frac{\bullet}{p}\right)_n$), is a homomorphism in the upper component as well, so it is natural to relate these two homomorphisms: what is their ratio $\left(\frac{p}{q}\right)_n \left(\frac{q}{p}\right)_n^{-1}$? This will give us a natural algorithm to compute the Legendre symbol $\left(\frac{a}{p}\right)_n$. We will prove strong reciprocity at the end of this section, after we discuss the Hilbert symbol.

1.1 Weak reciprocity and the Legendre symbol

The key observations linking reciprocity to the Artin map are that a is a perfect n th power modulo \mathfrak{p} iff $a^{\frac{\mathfrak{N}\mathfrak{p}-1}{n}} \equiv 1 \pmod{\mathfrak{p}}$ (just like $\left(\frac{a}{p}\right) = a^{\frac{p-1}{2}}$ in the quadratic case), and the homomorphism $a \mapsto a^{\frac{\mathfrak{N}\mathfrak{p}-1}{n}}$ can be linked to the Frobenius map.

Definition 16.1.2: Let K be a number field containing an n th root of unity, and let \mathfrak{p} be a prime ideal with $an \perp \mathfrak{N}\mathfrak{p}$. Define the **Legendre symbol** $\left(\frac{a}{\mathfrak{p}}\right)_n$ to be the unique n th root of unity ζ such that

$$\zeta \equiv a^{\frac{\mathfrak{N}\mathfrak{p}-1}{n}} \pmod{\mathfrak{p}}.$$

To see this is well-defined, note the following two points.

1. The n th roots of unity are distinct modulo \mathfrak{p} because $n \perp \mathfrak{N}\mathfrak{p}$. Hence $\frac{\mathfrak{N}\mathfrak{p}-1}{n}$ is an integer.
2. $(a^{\frac{\mathfrak{N}\mathfrak{p}-1}{n}})^n = 1 \equiv 1 \pmod{\mathfrak{p}}$ by Fermat's little theorem so $a^{\frac{\mathfrak{N}\mathfrak{p}-1}{n}}$ is equivalent to a unique n th root of unity.

Proposition 16.1.3: Let K be a number field containing an n th root of unity, let \mathfrak{p} be a prime ideal with $an \perp \mathfrak{N}\mathfrak{p}$. Then a is a perfect n th power modulo \mathfrak{p} iff $\left(\frac{a}{\mathfrak{p}}\right)_n = 1$.

Proof. Let the residue field of \mathfrak{p} be k . As k^\times has order $\mathfrak{N}\mathfrak{p} - 1$ and is generated by 1 element, a is a perfect n th power modulo \mathfrak{p} iff $a^{\frac{\mathfrak{N}\mathfrak{p}-1}{n}} = \left(\frac{a}{\mathfrak{p}}\right)_n = 1$. □

Proposition 16.1.4: $\left(\frac{a}{\mathfrak{p}}\right)_n$ is a group homomorphism factoring through $\mathcal{O}_K/\mathfrak{p}$.

Proof. Clear. □

How can class field theory give us an expression like this? Well, the Frobenius element corresponding to \mathfrak{p} acts like taking the $\mathfrak{N}\mathfrak{p}$ power modulo p . How do we get to $a^{\frac{\mathfrak{N}\mathfrak{p}-1}{n}}$? By acting by the Frobenius on $\sqrt[n]{a}$ instead.

Proposition 16.1.5: The following holds:

$$\left(\frac{a}{\mathfrak{p}}\right)_n = \frac{[\psi_{L/K}(\mathfrak{p})](\sqrt[n]{a})}{\sqrt[n]{a}},$$

where $L = K(\sqrt[n]{a})$.

Proof. First note $\mathfrak{p} \nmid an$ implies that $K(\sqrt[n]{a})/K$ is unramified at \mathfrak{p} , by Theorem 8.2.5.

By definition $\psi_{L/K}(\mathfrak{p})$ is the homomorphism that sends b to $b^{\mathfrak{N}\mathfrak{p}}$ modulo \mathfrak{p} . Thus

$$[\psi_{L/K}(\mathfrak{p})](\sqrt[n]{a}) \equiv \sqrt[n]{a}^{\mathfrak{N}\mathfrak{p}} \equiv a^{\frac{\mathfrak{N}\mathfrak{p}-1}{n}} \sqrt[n]{a} \pmod{\mathfrak{p}}.$$

But $\sqrt[n]{a}$ satisfies $X^n - a = 0$, so $(\mathfrak{p}, L/K)$ must send $\sqrt[n]{a}$ to $\zeta \sqrt[n]{a}$ where ζ is some root of unity. The above equation shows that we must have $\zeta = \left(\frac{a}{\mathfrak{p}}\right)_n$, as needed. □

We define $\left(\frac{a}{\mathfrak{b}}\right)_n$ for any $\mathfrak{b} \in I_K^{(na)}$ by extending multiplicatively the map $\left(\frac{a}{\mathfrak{p}}\right)_n$, originally defined for primes \mathfrak{p} . Equivalently (by Proposition 16.1.5), define $\left(\frac{a}{\mathfrak{b}}\right)_n = \frac{[\psi_{L/K}(\mathfrak{b})](\sqrt[n]{a})}{\sqrt[n]{a}}$.¹

We can now prove weak reciprocity.

¹We can extend the definition to all prime elements p by defining $\left(\frac{a}{p}\right)_n = \frac{\phi_{L/K}(i_v(p))(\sqrt[n]{a})}{\sqrt[n]{a}}$, then extend

Proof of Theorem 16.1.1. By Proposition 16.1.5,

$$\left(\frac{a}{\mathfrak{p}}\right)_n = \frac{[\psi_{K(\sqrt[n]{a})/K}(\mathfrak{p})](\sqrt[n]{a})}{\sqrt[n]{a}}. \quad (16.1)$$

Taking $a = p$ and $\mathfrak{p} = (q)$, we get

$$\left(\frac{p}{q}\right)_n = \frac{[\psi_{K(\sqrt[n]{p})/K}(q)](\sqrt[n]{p})}{\sqrt[n]{p}}.$$

Let \mathfrak{m} be the conductor of $K(\sqrt[n]{p})/K$. Since $\psi_{K(\sqrt[n]{p})/K}$ is an homomorphism on $I_K^{\mathfrak{m}}/i(P_{K,1}(\mathfrak{m}))$ (Theorem 11.4.1), its kernel contains $i(P_{K,1}(\mathfrak{m}))$. In other words, when $q \in i(P_{K,1}(\mathfrak{m}))$, then $\left(\frac{p}{q}\right) = \frac{[\psi_{K(\sqrt[n]{p})/K}(q)](\sqrt[n]{p})}{\sqrt[n]{p}} = \frac{\text{id}(\sqrt[n]{p})}{\sqrt[n]{p}} = 1$ and p is a perfect n th power modulo q . \square

1.2 Strong reciprocity and the Hilbert symbol

To prove strong reciprocity we need to actually compute (16.1). Supposing \mathfrak{p} is a principal ideal (b) , our statement about reciprocity seems to suggest that b and a play similar roles in the equation:²

$$\left(\frac{a}{b}\right)_n = \frac{[\psi_{L/K}(b)](\sqrt[n]{a})}{\sqrt[n]{a}}. \quad (16.2)$$

However, (16.2) is not symmetric. We seek to symmetrize it.

But look at Proposition 13.2.2. Equation (16.1) is the character corresponding to the element $a \in K^\times$. Using the map in Kummer Theory, we can get the equation symmetric in a and b . In fact, we did this already when we defined the Hilbert symbol.

If motivation was lacking when we defined the Hilbert symbol, hopefully this clears things up: it explains and clarify the duality in a and b observed above by making it symmetric in a and b .

Proposition 16.1.6: Let $b \nmid n$ be prime in K and K_b the completion at b . Let $(\cdot, \cdot)_b : K_b^\times/K_b^{\times n} \times K_b^\times/K_b^{\times n} \rightarrow \mu_n$ denote the Hilbert symbol. Then for $a \perp b$,

$$(a, b)_{b,n} = \left(\frac{a}{b}\right)_n.$$

In general, if $K_\pi(\sqrt[n]{a})/K_\pi(a)$ is unramified,

$$(a, b)_{\pi,n} = \left(\frac{(-1)^{v(a)v(b)} a^{v(b)} b^{-v(a)}}{\pi}\right)_n.$$

where $(a, b)_{v,n}$ denotes $(a, b)_n$ when a, b are considered in K_v .

the definition of $\left(\frac{a}{b}\right)_n$ to encompass any $b \in K^\times$ by multiplicativity. For instance, in the case $n = 2$, this gives the Jacobi symbol. For $b = 2$, $\left(\frac{a}{2}\right)$ tells us whether a is a perfect square modulo *any power of 2*.

²Caution: we're using the Artin map on ideals; we write $\psi_{L/K}(b)$ to mean $\psi_{L/K}((b))$. In contrast, $\phi_{L/K}(b) = 1$ since $b \in K$.

Proof. Proposition 16.1.5 and Proposition 14.6.3 give

$$\left(\frac{a}{b}\right)_n = \frac{[\psi_{L/K}(b)](\sqrt[n]{a})}{\sqrt[n]{a}} = \frac{[\psi_{L_b/K_b}(b)](\sqrt[n]{a})}{\sqrt[n]{a}} = (a, b)_{b,n}.$$

For the second part write $a = \pi^j u$ and $b = \pi^k u'$ where u, u' are units, and use bilinearity 14.6.4 to compute

$$\begin{aligned} (\pi^j u, \pi^k u') &= (\pi, \pi^k u')^j (u, \pi)^k && (u, u') = 1 \text{ since } K(\sqrt[n]{a}) \text{ unramified, 14.6.5} \\ &= (\pi, -\pi)^{jk} (\pi, (-1)^k u')^j \left(\frac{u}{\pi}\right)_n^k && \text{by the first part} \\ &= ((-1)^k u', \pi)^{-j} \left(\frac{u}{\pi}\right)_n^k && (\pi, -\pi) = 1, \text{ Theorem 14.6.4(2)} \\ &= \left(\frac{(-1)^k u'}{\pi}\right)_n^{-j} \left(\frac{u}{\pi}\right)_n^k \\ &= \left(\frac{(-1)^{jk} u^k u'^{-j}}{\pi}\right)_n \\ &= \left(\frac{(-1)^{v(a)v(b)} a^{v(b)} b^{-v(a)}}{\pi}\right)_n. \end{aligned}$$

□

The last main ingredient is the product formula for Hilbert symbols.

Theorem 16.1.7 (Product formula for Hilbert symbols). *Let K be a number field containing the n th roots of unity. Then*

$$\prod_{v \in V_K} (a, b)_v = 1.$$

Proof. Using the fact that the global Artin map can be written as the product of local Artin maps,

$$\prod_{v \in V_K} \phi_{K_v(\sqrt[n]{a})/K_v}(b) = \phi_K(b) = 1,$$

because ϕ_K is the identity on K . Now operate on this by the character $\chi(\sigma) = \frac{\sigma(\sqrt[n]{a})}{\sqrt[n]{a}} \in K$ and use Proposition 14.6.3 to get

$$\prod_{v \in V_K} (a, b)_v = \prod_{v \in V_K} \chi(\phi_{K(\sqrt[n]{a})/K}(b)) = 1.$$

□

Combining Proposition 16.1.6 and 16.1.7 gives the strong reciprocity law.

Theorem 16.1.8 (Strong reciprocity). *Let K be a number field containing a primitive n th root of unity and suppose a, b, n are pairwise relatively prime. Then*

$$\left(\frac{a}{b}\right)_n \left(\frac{b}{a}\right)_n^{-1} = \prod_{v|n\infty} (b, a)_{v,n}.$$

Suppose b, n are relatively prime and a is a prime dividing n . Then

$$\left(\frac{a}{b}\right)_n = \prod_{v|n\infty} (a, b)_{v,n}.$$

Proof. Suppose a, b, n are pairwise relatively prime. For a number c let $S(c)$ denote the finite places v where $v(c) \neq 0$. We calculate $\left(\frac{a}{b}\right)_n$ and $\left(\frac{b}{a}\right)_n$ using multiplicativity. We have

$$\begin{aligned} \left(\frac{a}{b}\right)_n \left(\frac{b}{a}\right)_n^{-1} &= \left(\frac{a}{\prod_{\pi \in S(b)} \pi^{v_\pi(b)}}\right) \left(\frac{b}{\prod_{\pi \in S(a)} \pi^{v_\pi(a)}}\right)^{-1} & (b) &= \left(\prod_{\pi \in S(b)} \pi^{v_\pi(b)}\right), \quad (a) = \left(\prod_{\pi \in S(a)} \pi^{v_\pi(a)}\right) \\ &= \prod_{v_\pi \in S(b)} \left(\frac{a}{\pi}\right)_n^{v_\pi(b)} \prod_{v_\pi \in S(a)} \left(\frac{b}{\pi}\right)_n^{-v_\pi(a)} \\ &= \prod_{v_\pi \in S(b)} \left(\frac{a}{\pi}\right)_n^{v_\pi(b)} \prod_{v_\pi \in S(a)} \left(\frac{b^{-v_\pi(a)}}{\pi}\right)_n \\ &= \prod_{v \in S(b)} (a, b)_v \prod_{v \in S(a)} (a, b)_v && \text{by Proposition 16.1.6} \\ &= \prod_{v \nmid n\infty} (a, b)_v && (a, b)_\mathbb{C} = 1, (a, b)_v = 1 \text{ when } a, b \in U_v, \text{ 14.6.5} \\ &= \prod_{v|n\infty} (b, a)_v \end{aligned}$$

where in the last step we used the product formula 16.1.7, which tells us $\prod_{v \in V_K} (a, b)_v = 1$.

Now suppose a is a prime dividing n . Then again using multiplicativity, Proposition 16.1.6, and the fact that $(a, b)_v = 1$ for $v | n\infty$, $n \nmid a$ (Corollary 14.6.5),

$$\left(\frac{a}{b}\right)_n = \prod_{v_\pi \in S(b)} \left(\frac{a}{\pi}\right)_n^{v_\pi(b)} = \prod_{v \in S(b)} (a, b)_v = \prod_{v|n\infty} (a, b)_v.$$

□

In practice, we can compute the action of the Hilbert symbol for each $v | n\infty$, since $K_v^\times / K_v^{\times n}$ is a finite set. We will carry out these computations in the cases $n = 2, 4$, for $K = \mathbb{Q}$ and $\mathbb{Q}(i)$.

1.3 Quadratic and biquadratic reciprocity

We derive quadratic and biquadratic reciprocity using Theorem 16.1.8.

Theorem 16.1.9 (Quadratic reciprocity). *Let p, q be odd primes. Then*

$$\left(\frac{-1}{p}\right) = (-1)^{\frac{p-1}{2}}, \quad \left(\frac{2}{p}\right) = (-1)^{\frac{p^2-1}{8}}, \quad \left(\frac{p}{q}\right) \left(\frac{q}{p}\right) = (-1)^{\frac{p-1}{2} \cdot \frac{q-1}{2}}.$$

Proof. The first follows from definition of the Legendre symbol. By strong reciprocity 16.1.8,

$$\begin{aligned} \left(\frac{2}{p}\right) &= (2, p)_2 \\ \left(\frac{p}{q}\right) \left(\frac{q}{p}\right) &= (p, q)_2. \end{aligned}$$

Let $U^{(i)}$ denote $1 + (2)^i$ in \mathbb{Q}_2 .

1. We have $(2, p)_2 = 1$ iff p is a norm from $\mathbb{Q}_2(\sqrt{2})$ (Theorem 14.6.4), iff p is in the form $x^2 - 2y^2$ in \mathbb{Q}_2 . Looking at this modulo 8, we must have $p \in \{1, 5\}2^{\mathbb{Z}}$. This is sufficient as we know $[\mathbb{Q}_2^\times : \text{Nm}_{\mathbb{Q}_2(\sqrt{2})/\mathbb{Q}_2}(\mathbb{Q}_2(\sqrt{2})^\times)] = [\mathbb{Q}_2(\sqrt{2}) : \mathbb{Q}_2] = 2$, so we must have $\text{Nm}_{\mathbb{Q}_2(\sqrt{2})/\mathbb{Q}_2}(\mathbb{Q}_2(\sqrt{2})^\times) = \{1, 5\}2^{\mathbb{Z}}$. Hence $(2, p)_2 = 1$ iff $p \equiv 1, 5 \pmod{8}$, iff $\frac{p^2-1}{8}$ is even. This gives

$$\left(\frac{2}{p}\right) = (-1)^{\frac{p^2-1}{8}}.$$

2. We have $(p, q)_2 = 1$ iff $q \in N := \text{Nm}_{\mathbb{Q}_2(\sqrt{p})/\mathbb{Q}_2}(\mathbb{Q}_2(\sqrt{p})^\times)$, iff q is in the form $x^2 - py^2$.
 - (a) If $p \equiv 1 \pmod{4}$, then $x^2 - py^2$ can attain any odd residue modulo 8. Since $[Q : N] = [\mathbb{Q}_2(\sqrt{p}) : \mathbb{Q}_2] \leq 2$, we have $U^{(3)}2^{2\mathbb{Z}} = \mathbb{Q}_2^{\times 2} \subseteq N$. Since N contains all residues modulo 8, $U2^{2\mathbb{Z}} \subseteq N$. Hence $q \in N$, and $(p, q)_2 = 1$.
 - (b) If $p \equiv 3 \pmod{4}$, then $x^2 - py^2$ cannot be $3 \pmod{4}$. Hence $N = U^{(2)}2^{\mathbb{Z}}$, and $q \in N$ iff $q \equiv 1 \pmod{4}$. Hence $(p, q)_2 = 1$ iff $q \equiv 1 \pmod{4}$.

It remains to note $(-1)^{\frac{p-1}{2} \cdot \frac{q-1}{2}} = 1$ iff either $p \equiv 1 \pmod{4}$ or $q \equiv 1 \pmod{4}$. \square

Theorem 16.1.10 (Biquadratic reciprocity). *Suppose p, q are primes in $\mathbb{Z}[i]$ with $p, q \equiv 1 \pmod{(1+i)^3}$. Then*

$$\left(\frac{p}{q}\right)_4 = (-1)^{\frac{\text{nr}_p-1}{4} \cdot \frac{\text{nr}_q-1}{4}} \left(\frac{q}{p}\right)_4.$$

Note every prime contains an associate that is equivalent to $1 \pmod{4}$.

Proof. Note $p \equiv 1 \pmod{(1+i)^3}$ means $p \equiv 1$ or $1 + 2i \pmod{(1+i)^3}$.

By strong reciprocity 16.1.8,

$$\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4^{-1} = (q, p)_{2,4} = (p, q)_{2,4}^{-1}.$$

We have $(p, q)_{2,4} = 1$ iff $q \in \text{Nm}_{\mathbb{Q}_2(\sqrt{p})/\mathbb{Q}_2}(\mathbb{Q}_2(\sqrt{p})^\times)$. Consider 2 cases.

1. $\mathfrak{N}p \equiv 1 \pmod{8}$. Equivalently (writing out $p = a + bi$ and calculating the norm), $p \equiv 1 \pmod{8}$. We can calculate that $(1 + i)^{3\mathbb{Z}}U^{(3)} \subseteq N := \text{Nm}_{\mathbb{Q}_2(\sqrt{p}, i)/\mathbb{Q}_2(i)}(\mathbb{Q}_2(\sqrt{p}, i)^\times)$, so $q \in N$. (The calculations are lengthy, but here's the idea: by examining the structure of $\mathbb{Q}_2(i)$, or using Proposition 15.3.2, we find that $\mathbb{Q}_2(i)^{\times 4} = U^{(7)}(1 + i)^{4\mathbb{Z}}$. Hence the norm group N satisfies

$$U^{(7)}(1 + i)^{4\mathbb{Z}} \subseteq N \subseteq \mathbb{Q}_2(i)^\times$$

and has index at most 4. Now calculate the norm of enough numbers in $\mathbb{Q}_2(\sqrt{p}, i)$ until we can determine $(1 + i)^{3\mathbb{Z}}U^{(3)} \subseteq N$. Using a computer algebra system is advised.)

2. $\mathfrak{N}p \equiv 5 \pmod{8}$. Equivalently, $p \equiv 5 \pmod{8}$. We can calculate that $(1 + i)^{4\mathbb{Z}}U^{(3)} \subseteq \text{Nm}_{\mathbb{Q}_2(\sqrt{p})/\mathbb{Q}_2}(\mathbb{Q}_2(\sqrt{p})^\times)$ but $(1 + 2i)(1 + i)^{4\mathbb{Z}}U^{(3)} \not\subseteq \text{Nm}_{\mathbb{Q}_2(\sqrt{p})/\mathbb{Q}_2}(\mathbb{Q}_2(\sqrt{p})^\times)$. Hence $(p, q)_4 = 1$ iff $q \equiv 1 \pmod{4}$, i.e. iff $\mathfrak{N}q \equiv 1 \pmod{8}$.

In the case where $\mathfrak{N}p, \mathfrak{N}q \equiv 5 \pmod{8}$, we have $(p, q)_4^2 = (p, q^2)_4 = 1$ but $(p, q)_4 \neq 1$ so $(p, q)_4 = -1$. □

1.4 Reciprocity for odd primes

We give an algorithm for finding reciprocity laws for $\mathbb{Q}(\zeta_p)/\mathbb{Q}$ for p prime, and then specialize to $p = 3$.

Theorem 16.1.11. *Let p be an odd prime, let $K = \mathbb{Q}(\zeta_p)$, and let v be the valuation corresponding to $1 - \zeta_p$. Let $\pi = 1 - \zeta_p$. Then the elements*

$$\begin{aligned} & \pi \\ \eta_1 &= 1 - \pi = \zeta_p \\ \eta_2 &= 1 - \pi^2 \\ & \vdots \\ \eta_p &= 1 - \pi^p \end{aligned}$$

generate $K_v^\times / K_v^{\times p}$, and $(a, b)_v$ is the unique skew-symmetric pairing $K_v^\times \times K_v^\times \rightarrow \mu_p$ satisfying the following.

1. $(\eta_i, \eta_j)_v = (\eta_i, \eta_{i+j})_v (\eta_{i+j}, \eta_j)_v (\eta_{i+j}, \pi)_v^{-j}$.

2. $(\eta_i, \pi)_v = \begin{cases} 1, & 1 \leq i \leq p - 1 \\ \zeta, & i = p. \end{cases}$

Moreover, if $i + j \geq p + 1$, then $(a, b)_v = 1$ for all $a \in U^{(i)}$ and $b \in U^{(j)}$.

We start with the following lemma.

Lemma 16.1.12. *Let K be a number field containing p th roots of unity. Let ζ be a primitive p th root of unity, $\pi = 1 - \zeta$, and \mathfrak{p} a prime dividing π . Suppose $a = 1 + \pi^p c$ with $\pi = 1 - \zeta$ and $c \in \mathcal{O}_v$. Then for all b ,*

$$(a, b)_{\mathfrak{p}} = \zeta^{-\mathrm{tr}_{k/\mathbb{F}_p}(\bar{c})v_{\mathfrak{p}}(b)}.$$

We will just need the case where $K = \mathbb{Q}(\zeta_p)$, in which case $k = \mathbb{F}_p$.

Proof. Because $a \notin \mathfrak{p}$, $K(\sqrt[p]{a})/K$ is unramified by Lemma 8.2.5. We have (cf. Proposition 6.2.2)

$$\begin{aligned} \frac{\zeta^p - 1}{\zeta - 1} &= 0 \\ \implies \frac{(1 - \pi)^p - 1}{(1 - \pi) - 1} &= 0 \\ \implies \pi^{p-1} - p\pi^{p-2} + \cdots + p &= 0 \\ \implies \pi^{p-1} &\equiv -p \pmod{p\pi} \end{aligned}$$

and we get

$$\frac{\pi^{p-1}}{p} \equiv -1 \pmod{\mathfrak{p}}. \quad (16.3)$$

Let $\alpha = \sqrt[p]{a}$ be a p th root of a , and write $\alpha = 1 + \pi x$, where $x \in L$. Now $\alpha^m - a = 0$ becomes $(1 + \pi x)^p - (1 + \pi^p c) = 0$. Hence x is a zero of the polynomial $f(X) = \frac{1}{\pi^p} ((1 + \pi x)^p - (1 + \pi^p c))$. Using (16.4), we find that $f(X)$ is integral, so $x \in \mathcal{O}_L$, and that modulo π ,

$$f(X) = \frac{1}{\pi^p} (\pi^p x^p + p\pi x - \pi^p c) \equiv x^p - x - c \pmod{\pi}$$

Let $\mathfrak{N}\mathfrak{p} = p^f$. Letting σ be the Frobenius, we find that $\sigma(x) \equiv x^{p^f} \pmod{\mathfrak{p}}$. Note that

$$x^{p^j} \equiv (x + c)^{p^{j-1}} \equiv x^{p^{j-1}} + c^{p^{j-1}} \pmod{\mathfrak{p}}.$$

Hence by induction

$$\sigma(x) = x^{p^f} = x + \bar{c} + \bar{c}^p + \cdots + \bar{c}^{p^{f-1}} = x + \mathrm{tr}_{k/\mathbb{F}_p}(\bar{c}) \quad (16.4)$$

in k . Now by Proposition 14.6.3,

$$(a, b)_{\mathfrak{p}} = \frac{[\phi_{K_{\pi}(\alpha)/K_{\pi}}(b)](\alpha)}{\alpha} = \frac{\sigma^{v(b)}(\alpha)}{\alpha}$$

To get the second equality, note that by construction, $\phi_{K_{\pi}(\alpha)/K_{\pi}}(\pi)$ is the Frobenius element; as $K_{\pi}(\alpha)/K_{\pi}$ is unramified, $U_{K_{\pi}} \subseteq \ker \phi_{K_{\pi}(\alpha)/K_{\pi}}$ (Example 14.5.1), and the Artin map depends only on $v(b)$. We have

$$(a, b)_v = \zeta^n \text{ where } \zeta^n \alpha = \sigma^{v(b)}(\alpha);$$

to find n we reduce both sides modulo $\mathfrak{p}\pi$. We calculate

$$\zeta\alpha \equiv (1 - \pi)(1 + \pi x) \equiv 1 + (x - 1)\pi \pmod{\mathfrak{p}\pi} \quad (16.5)$$

$$\implies \zeta^n\alpha \equiv 1 + (x - n)\pi \pmod{\mathfrak{p}\pi} \quad (16.6)$$

$$\sigma^{v(b)}(x) \equiv x + v(b) \operatorname{tr}_{k/\mathbb{F}_p}(\bar{c}) \pmod{\pi} \quad \text{by (16.4)} \quad (16.7)$$

$$\implies \sigma^{v(b)}(\alpha) \equiv 1 + (x + v(b) \operatorname{tr}_{k/\mathbb{F}_p}(\bar{c}))\pi \pmod{\mathfrak{p}\pi}. \quad (16.8)$$

Matching (16.6) and (16.8) gives $n = -v(b) \operatorname{tr}_{k/\mathbb{F}_p}(\bar{c})$ and

$$(a, b)_v = \zeta^{-v(b) \operatorname{tr}_{k/\mathbb{F}_p}(\bar{c})} \pmod{\pi}.$$

□

In particular, note that $(a, b)_v = 1$ if $a \equiv 1 \pmod{\pi^{p+1}}$. By nondegeneracy of the pairing (Theorem 14.6.4), we get that $a \in (K_v^\times)^p$. Hence $U^{(p+1)} \subseteq (K_v^\times)^p$.

Proof of Theorem 16.1.11. Note that η_i generates $U^{(i)}/U^{(i+1)}$, and π generates $K_\pi^\times/(K_\pi^\times)^p U^{(1)}$. As mentioned above, $U^{(p+1)} \subseteq (K_\pi^\times)^p$ so $\pi, \eta_1, \dots, \eta_p$ generate $K_\pi^\times/(K_\pi^\times)^p$. Since the group has order $\frac{p^2}{|p|_{v_\pi}} = p^{p+1}$ (Proposition 15.3.2), these generators are independent.

We use a relation between the η_i, η_j to derive the first relation. Namely, we have $\frac{\eta_j}{\eta_{i+j}} + \pi^j \frac{\eta_i}{\eta_{i+j}} = 1$, so

$$\left(\frac{\eta_j}{\eta_{i+j}}, \pi^j \frac{\eta_i}{\eta_{i+j}} \right)_p = 1$$

by Theorem 14.6.4. Note $(a, -1) = 1$ for any a because -1 is a p th power. Expanding the above bilinearity gives

$$\begin{aligned} 1 &= (\eta_j, \pi^j \eta_i)(\eta_{i+j}, \pi^j \eta_i)^{-1} \underbrace{(\eta_{i+j}, -\eta_{i+j})}_1 \underbrace{(\eta_{i+j}, -1)}_1 (\eta_j, \eta_{i+j})^{-1} \\ &= (\eta_j, \eta_i) \underbrace{(\eta_j, \pi^j)}_{=1, \eta_j + \pi^j = 1} (\eta_{i+j}, \pi)^{-j} (\eta_{i+j}, \eta_i)^{-1} (\eta_j, \eta_{i+j})^{-1} \\ &= (\eta_i, \eta_j)^{-1} (\eta_{i+j}, \pi)^{-j} (\eta_{i+j}, \eta_i)^{-1} (\eta_{i+j}, \eta_j) \\ \implies (\eta_i, \eta_j) &= (\eta_i, \eta_{i+j})(\eta_{i+j}, \eta_j)(\eta_{i+j}, \pi)^{-j}. \end{aligned}$$

This shows item 1. For item 2, note for $1 \leq i \leq p-1$ that since $\eta_i + \pi^i = 1$,

$$1 = (\eta_i, \pi^i) = (\eta_i, \pi)^i \implies 1 = (\eta_i, \pi).$$

For $i = p$, we use the lemma to find

$$(\eta_p, \pi)_v = \zeta^{-\operatorname{tr}_{k/\mathbb{F}_p}(-1)} = \zeta$$

because $k = \mathbb{F}_p$.

Note that if $i + j \geq p + 1$, then $\eta_{i+j} \in U^{(p+1)} \subseteq (K_v^\times)^p$ so item 1 gives that $(\eta_i, \eta_j) = 1$. Now as a skew-symmetric bilinear pairing (η_i, η_j) is determined by items 1 and 2, because we can expand (η_i, η_j) using item 1, then repeatedly expand factors (the indices increase each time) until we only have factors in the form (\bullet, π) , and use item 2 to get a value out. □

We now use this to derive cubic reciprocity.

Theorem 16.1.13 (Cubic reciprocity). *Let $K = \mathbb{Q}(\omega)$, where $\omega = \zeta_3 = \frac{-1+\sqrt{-3}}{2}$. For $a \equiv \pm 1 \pmod{3\mathcal{O}_K}$, write*

$$a = \pm(1 + 3(m + n\omega)).$$

Then

$$\begin{aligned} \left(\frac{b}{a}\right)_3 &= \left(\frac{a}{b}\right)_3 && \text{if } b \perp a, b \equiv \pm 1 \pmod{3\mathcal{O}_K} \\ \left(\frac{\omega}{a}\right)_3 &= \omega^{-m-n} \\ \left(\frac{1-\omega}{a}\right)_3 &= \omega^m. \end{aligned}$$

Note that if $q \not\equiv 1 \pmod{3}$ is prime, then $3 \nmid |\mathbb{F}_q^\times|$ so any element of \mathbb{F}_q^\times is a cubic residue. Note any element of K relatively prime to 3 can be written in the form $\omega^i(1-\omega)^j a$ where $a \equiv \pm 1 \pmod{3\mathcal{O}_K}$.

Proof. First suppose $a, b \equiv 1 \pmod{3}$. By Strong Reciprocity 16.1.8,

$$\left(\frac{a}{b}\right)_3 \left(\frac{b}{a}\right)_3^{-1} = (b, a)_3.$$

Note $a, b \in U^{(2)}$ so by Theorem 16.1.11, $(b, a)_3 = 1$. This shows the first equation.

For the second, letting $\pi = 1 - \omega$, note that

$$(1 - \pi^2)^\alpha (1 - \pi^3)^\beta = (1 + 3\omega)^\alpha (1 + 3(2\omega + 1))^\beta \in [1 + 3(\beta + (2\beta + \alpha)\omega)]U^{(4)}$$

Setting $\alpha = n - 2m$ and $\beta = m$, we get

$$\begin{aligned} a &\in (1 - \pi^2)^{n-2m} (1 - \pi^3)^m U^{(4)} \\ (1 - \pi^2)^{2m-n} (1 - \pi^3)^{-m} &\in aU^{(4)} \end{aligned}$$

Now Theorem 16.1.11 tells us

$$\begin{aligned} (\omega, 1 - \pi^2) &= (\eta_1, \eta_2) = (\eta_3, \pi)^{-2} = \omega \\ (\omega, 1 - \pi^3) &= (\eta_1, \eta_3) = 1 \\ (\pi, 1 - \pi^2) &= (\eta_2, \pi)^{-1} = 1 \\ (\pi, 1 - \pi^3) &= (\eta_3, \pi)^{-1} = \omega^{-1}. \end{aligned}$$

Thus

$$\begin{aligned} \left(\frac{\omega}{a}\right) &= (\omega, (1 - \pi^2)^{2m-n} (1 - \pi^3)^{-m}) = \omega^{-m-n} \\ \left(\frac{\pi}{a}\right) &= (\pi, (1 - \pi^2)^{2m-n} (1 - \pi^3)^{-m}) = \omega^m. \end{aligned}$$

□

As an application, we show the following.

Theorem 16.1.14. *If $q \equiv 1 \pmod{3}$ is a prime, then 2 is a cubic residue modulo q iff q is in the form*

$$q = x^2 + 27y^2, \quad \text{for some } x, y \in \mathbb{Z}.$$

Proof. Since $q \equiv 1 \pmod{3}$, q splits in \mathcal{O}_K as $\alpha\bar{\alpha}$. By multiplying by a root of unity, we may assume $\alpha \equiv 1 \pmod{3\mathcal{O}_K}$, i.e. α is in the form $\alpha = 3(x + y\omega) \pm 1$. In order for 2 to be a cubic residue, it must be a cubic residue modulo α . If $a^3 \equiv 2 \pmod{\alpha}$, then $\bar{a}^3 \equiv 2 \pmod{\bar{\alpha}}$, so it would also be a cubic residue modulo $\bar{\alpha}$ and hence modulo q .

Now $\left(\frac{2}{\alpha}\right) = 1$ iff $\left(\frac{\alpha}{2}\right) = 1$, by Cubic Reciprocity 16.1.13. Since 2 remains inert in \mathcal{O}_K , and the only cube in \mathbb{F}_4^\times is 1, we get that α must actually be in the form

$$\alpha = 6(x + y\omega) \pm 1.$$

Taking the norm gives

$$p = (6x + 3y \pm 1)^2 + 27y^2.$$

This is in the form $x'^2 + 27y'^2$; conversely, any prime in the form $x'^2 + 27y'^2$ must have $x' \equiv \pm 1 \pmod{3}$, and hence is in the above form. \square

2 Hasse-Minkowski Theorem

The global Artin map can be expressed as the product of local Artin maps. From class field theory, we get various “local-to-global” results such as the Hasse-Brauer-Noether Theorem 15.3.5 and the Hasse Norm Theorem 16.2.2. The most famous is the local-to-global principle for quadratic forms, the Hasse-Minkowski Theorem.

Definition 16.2.1: A quadratic form is said to **represent** a if there is a solution to $q(X_1, \dots, X_n) = a$ with $(x_1, \dots, x_n) \neq (0, \dots, 0)$. A quadratic form representing 0 is said to be **isotropic**.

(For a review of quadratic forms, see Chapter 4.)

Where class field theory comes in is that a quadratic form in 2 variables representing a number a can be interpreted as a norm equation, $a = x^2 + by^2$. We can write this as $a = (x + y\sqrt{b})(x - y\sqrt{b}) = \text{Nm}_{K(\sqrt{b})/K}(x + y\sqrt{b})$ when $\sqrt{b} \notin K$. Class field theory gives us a local-to-global theorem for norms, the Hasse Norm Theorem. This will prove the $n = 2$ case of Hasse-Minkowski. Then a series of elaborate reductions will prove the local-to-global principal for any number of variables.

2.1 Hasse norm theorem

Theorem 16.2.2 (Hasse norm theorem). *Suppose L/K is cyclic. Then a is a global norm iff it is a local norm everywhere: $a \in \text{Nm}_{L/K} L^\times$ iff $a \in \text{Nm}_{L^v/K_v} L^{v\times}$ for all $v \in V_K$.*

Compare this to the proof of Theorem 15.3.5.

Proof. The forward direction is clear.

Let $G = G(L/K)$. Take the long exact sequence in Tate cohomology of

$$0 \rightarrow L^\times \rightarrow \mathbb{I}_L \rightarrow \mathbf{C}_L \rightarrow 0$$

to get the top row of the following.

$$\begin{array}{ccccccc} H_T^{-1}(G, \mathbf{C}_L) & \longrightarrow & H_T^0(G, L^\times) & \longrightarrow & H_T^0(G, \mathbb{I}_L) & \longrightarrow & \cdots \\ \parallel & & \parallel & & \parallel & & \\ 0 & \longrightarrow & K^\times / \text{Nm}_{L/K} L^\times & \hookrightarrow & \bigoplus_{v \in V_K} K_v^\times / \text{Nm}_{K_v}(L^{v^\times}) & & \end{array} \quad (16.9)$$

We explain the bottom row. First note the equalities of H_T^0 are by definition of H_T^0 , plus Proposition 15.2.4. Next note cohomology is 2-periodic because G is cyclic (Proposition 12.12.1), and $H_T^1(G, \mathbf{C}_L) = 0$ by Theorem 15.3.1 (HT90 for ideles), so

$$H_T^{-1}(G, \mathbf{C}_L) = H_T^1(G, \mathbf{C}_L) = 0.$$

Then (16.9) gives that the map $K^\times / \text{Nm}_{L/K} L^\times \hookrightarrow \bigoplus_{v \in V_K} K_v^\times / \text{Nm}_{K_v}(L^{v^\times})$ is injective. If $a \in K^\times$ is a norm in every completion, then it is 0 in $\bigoplus_{v \in V_K} K_v^\times / \text{Nm}_{K_v}(L^{v^\times})$, hence 0 in $K^\times / \text{Nm}_{L/K} L^\times$, hence a global norm. \square

2.2 Quadratic forms

We prove the following.

Theorem 16.2.3 (Hasse-Minkowski). *Let K be a number field. The following hold.*

1. *A quadratic form f defined over K represents a iff f represents a in every completion K_v .*
2. *Two quadratic forms over K are equivalent iff they are equivalent over every completion K_v .*

First we note that item 1 implies item 2.

Proof that 1 implies 2. The forward direction is clear. For the reverse direction, induct on the rank n , $n = 0$ being the base case. Suppose f, g are equivalent over every completion K_v . Suppose f represents a . Then f represents a over every K_v . Since $g \sim f$ over every K_v , g represents a over every K_v . By item 1, g represents a .

Thus we can write $f \sim aX^2 + f'$, $g \sim aX^2 + g'$. Now $aX^2 + f' \sim aX^2 + g'$ over every K_v implies (see Serre [Ser73, IV.1.7, Prop. 4]) $f' \sim g'$ over every K_v . By the induction hypothesis, $f' \sim g'$ over K . Thus $f \sim g$. \square

Next we show that we can reduce item 1 to a statement about quadratic forms representing 0.

Lemma 16.2.4. *Suppose $\text{char}(K) \neq 2$. An nondegenerate isotropic quadratic form over K represents all of K .*

Proof. Let B be the bilinear form associated to q . Suppose $\mathbf{x} \neq 0$ is such that $q(\mathbf{x}) = 0$. Since q is nondegenerate, there exists \mathbf{y} such that $B(\mathbf{x}, \mathbf{y}) \neq 0$. Then $q(\mathbf{x} + a\mathbf{y}) = a^2q(\mathbf{y}) + 2aB(\mathbf{x}, \mathbf{y})$ attains every value as a ranges over K . \square

Lemma 16.2.5. *A quadratic form $q(X_1, \dots, X_{n-1})$ represents a iff $q(X_1, \dots, X_{n-1}) - aX_n^2$ represents 0.*

Proof. For the forward direction, suppose $q(x_1, \dots, x_{n-1}) = a$. Then $q(x_1, \dots, x_{n-1}) - a \cdot 1^2 = 0$.

For the reverse direction, let (x_1, \dots, x_n) be a solution. If $x_n = 0$ then $q(x_1, \dots, x_n) = 0$ so q represents 0. Thus q is isotropic and represents a . If $x_n \neq 0$ then $q\left(\frac{x_1}{x_n}, \dots, \frac{x_{n-1}}{x_n}\right) = a$. \square

Thus it suffices to prove item 1 of Hasse-Minkowski for $a = 0$. Specifically, item 1 for forms with n variables is a consequence of item 1 for $a = 0$ for forms with $n + 1$ variables. We now prove Hasse-Minkowski. Every quadratic form over a field not of characteristic 2 can be put in diagonal form, so it suffices to consider diagonal forms. By scaling, we may assume one of the coefficients is 1.

2.2.1 Proof for $n \leq 2$

For $n = 1$ the theorem is trivial. For $n = 2$, we need the following.

Lemma 16.2.6. *An element $a \in K$ is a square iff it is a square in every completion K_v .*

Proof. (cf. the proof of Proposition 15.2.8) The forward direction is clear.

So suppose a is a square in every completion. Then $K_v(\sqrt{a}) = K$ so $\text{Nm}_{K_v(\sqrt{a})/K_v} K_v(\sqrt{a})^\times = K_v^\times$. This shows $\text{Nm}_{K(\sqrt{a})/K}(\mathbb{I}_{K(\sqrt{a})}) = \mathbb{I}_K$. By the first inequality 15.2.1,

$$[K(\sqrt{a}) : K] \leq [\mathbb{I}_K : \text{Nm}_{K(\sqrt{a})/K}(\mathbb{I}_{K(\sqrt{a})})] = 1$$

so $K(\sqrt{a}) = K$, i.e. a is a square in K . \square

Now a quadratic form

$$q(X, Y) = X^2 - aY^2$$

represents 0 iff a is a square (it rearranges to $\left(\frac{X}{Y}\right)^2 = a$), so q represents 0 over K if it represents 0 over every K_v .

2.2.2 Proof for $n = 3$

As promised, we re-express the condition for $p(x)$ to represent 0 as a condition on norms.

Lemma 16.2.7. *Let K be any field. A quadratic form*

$$q(X, Y, Z) = X^2 - bY^2 - cZ^2$$

represents 0 iff $c \in \text{Nm}_{K(\sqrt{b})/K}(K(\sqrt{b})^\times)$.

Proof. Note if $q(x, y, z) = 0$ with $z = 0$, then b must be a perfect square. If b is a perfect square then $K(\sqrt{b})/K$ is trivial and c is trivially a norm.

So it suffices to consider solutions with $z \neq 0$ and b not a perfect square. In this case,

$$x^2 - by^2 - cz^2 = 0$$

iff

$$c = \left(\frac{x}{z}\right)^2 - b\left(\frac{y}{z}\right)^2 = \left(\frac{x}{z} - \sqrt{b} \cdot \frac{y}{z}\right) \left(\frac{x}{z} + \sqrt{b} \cdot \frac{y}{z}\right) = \text{Nm}_{K(\sqrt{b})/K} \left(\frac{x}{z} - \sqrt{b} \frac{y}{z}\right).$$

□

By the Hasse Norm Theorem 16.2.2, $c \in \text{Nm}_{K(\sqrt{b})/K}(K(\sqrt{b})^\times)$ if this is true for every completion K_v . Combined with the lemma above, this gives Hasse-Minkowski for $n = 3$.

We will need the following in the proof for $n \geq 5$.

Lemma 16.2.8. *The form $f = X^2 - bY^2 - cZ^2$ represents 0 in a local field K_v iff $(b, c)_v = 1$. Moreover, f represents 0 in K_v for all but a finite number of places v .*

Proof. Note f represents 0 iff $c \in \text{Nm}_{K(\sqrt{b})/K}(K(\sqrt{b})^\times)$, which is equivalent to $(b, c)_v = 1$ by Theorem 14.6.4. Only finitely many of these are not equal to 1 by Corollary 14.6.5. □

2.2.3 Proof for $n = 4$

We reduce the $n = 4$ case to the $n = 3$ case (but for a different field extension) by the following string of equivalences. The brilliant idea here is to turn the quadratic form equation into a quotient of norms.

Theorem 16.2.9. *For any field K , the following are equivalent, for $a, b, c \in K^\times$.*

1. *The form $f(X, Y, Z, T) = X^2 - bY^2 - cZ^2 + acT^2$ represents 0 in K .*
2. *c is a product of norms from $K(\sqrt{a})$ and $K(\sqrt{b})$:*

$$c \in \text{Nm}_{K(\sqrt{a})/K}(K(\sqrt{a})^\times) \text{Nm}_{K(\sqrt{b})/K}(K(\sqrt{b})^\times).$$

3. *$c \in \text{Nm}_{K(\sqrt{a}, \sqrt{b})/K(\sqrt{ab})}(K(\sqrt{a}, \sqrt{b})^\times)$.*

4. *The form $g(X, Y, Z) = X^2 - bY^2 - cZ^2$ represents 0 in $K(\sqrt{ab})$.*


Proof. (1) \iff (2): If (x, y, z, t) is a solution with $z^2 - at^2 = 0$, then $x^2 - by^2 = 0$ as well. Then a, b are squares in K and (2) is clear. So it suffices to consider solutions with $z^2 - at^2 \neq 0$. In that case,

$$x^2 - by^2 - cz^2 + act^2 = 0 \iff c = \frac{x^2 - by^2}{z^2 - at^2} = (x - \sqrt{b}y)(x + \sqrt{b}y)(z - \sqrt{a}t)^{-1}(z + \sqrt{a}t)^{-1},$$

and this has a solution iff (2) holds.

(4) \iff (3): Applying Lemma 16.2.7, we see (4) is equivalent to c being a norm from $K(\sqrt{b}, \sqrt{ab})/K(\sqrt{ab})$. But $K(\sqrt{b}, \sqrt{ab}) = K(\sqrt{a}, \sqrt{b})$.

(2) \iff (3): This is the hard part. We consider the field extensions



If any of a, b, ab is in $K^{\times 2}$ then the result is clear: If $a \in K^{\times 2}$ then both (2) and (3) are true for any c , since $K(\sqrt{a}) = K$ and $K(\sqrt{a}, \sqrt{b}) = K(\sqrt{ab})$. If $ab \in K^{\times 2}$ then $K(\sqrt{a}) = K(\sqrt{b})$ so both (2) and (3) are equivalent to $c \in \text{Nm}_{K(\sqrt{a})/K}(K(\sqrt{a})^\times)$.

Now assume $a, b, ab \notin K^{\times 2}$. In this case $G(K(\sqrt{ab})/K) \cong \mathbb{Z}/2 \times \mathbb{Z}/2$ with the 3 subextensions corresponding to 3 subgroups. Let σ be the non-identity element fixing $K(\sqrt{a})$, τ fix $K(\sqrt{b})$ and $\rho = \sigma\tau$ fix $K(\sqrt{ab})$. (I.e. σ switches $\pm\sqrt{b}$ and τ switches $\pm\sqrt{a}$.) We convert the statements in (2) and (3) into the language of Galois theory, using the fixed field theorem.

Note (2) is equivalent to the following:

$$(2)' : \quad \text{There exist } x, y \in L, \quad \sigma(x) = x, \tau(y) = y, x\rho(x)y\rho(y) = c.$$

To go between these statements take

$$x' = z - \sqrt{a}t, y = x - \sqrt{b}y$$

and note ρ conjugates both \sqrt{a} and \sqrt{b} . Similarly, (3) is equivalent to the following.

$$(3)' : \quad \text{There exists } z \in L, \quad z\rho(z) = c;$$

just take $z' = (x - \sqrt{b}y)(z - \sqrt{a}z)$. To go from (2)' to (3)' just take $z = xy$. To go back from (3)' to (2)' requires more work. Given z , let $u = \frac{z \cdot \sigma(z)}{c}$. Now $\sigma(u) = u$ and $u\rho(u) = \frac{z\rho(z)\sigma(z)\sigma(\rho(z))}{c^2} = 1$. Since $\sigma(u) = u$. i.e. $u \in K(\sqrt{a})$, and $G(K(\sqrt{a})/K) = \{1, \tau|_{K(\sqrt{a})}\}$, by

Hilbert's Theorem 90 (13.1.1) there exists $x \in K(\sqrt{a})$ (i.e. x satisfying $\sigma(x) = x$) such that $\frac{\tau(x)}{x} = u$. Set $y = \frac{\rho(z)}{x}$. We've chosen x satisfying the conditions. For y , note

$$\begin{aligned} \tau(y) &= \frac{\sigma(z)}{\tau(x)} & \tau\rho &= \sigma \\ &= \frac{\sigma(z)}{xu} & \frac{\tau(x)}{x} &= u \\ &= \frac{c}{xz} & u &= \frac{z\sigma(z)}{c} \\ &= \frac{\rho(z)}{x} = y & z\rho(z) &= c. \end{aligned}$$

Finally, $xy\rho(xy) = \rho(z)\rho(\rho(z)) = c$. This shows (2)' \implies (3)' and finishes the proof. \square

Now we show Hasse-Minkowski holds for $n = 4$. By (1) \iff (4) in Theorem 16.2.9, Hasse-Minkowski for $f = X^2 - bY^2 - cZ^2 + acT^2$ over K is equivalent to Hasse-Minkowski for $g = X^2 - bY^2 - cZ^2$ over $K(\sqrt{ab})$, and we have already proved Hasse-Minkowski for $n = 3$.

2.2.4 Proof for $n \geq 5$

We now prove Hasse-Minkowski for $n \geq 5$. We proceed by induction. The idea is to “replace” $aX_1^2 + bX_2^2$ by just cX^2 .

Suppose it proved for $n - 1$, and write

$$f(X_1, \dots, X_n) = aX_1^2 + bX_2^2 - g(X_3, \dots, X_n).$$

Suppose f represents 0 in each K_v . Then there exists c_v such that

$$aX_1^2 + bX_2^2 = c_v = g(X_3, \dots, X_n)$$

has a nontrivial solution in K_v . By Lemma 16.2.8, there exists a finite set S such that g represents all elements of K_v when $v \notin S$. We only need to focus on $v \in S$.

Note $K_v^{\times 2}$ is open in K_v^\times by Theorem 8.1.5. By the Weak Approximation Theorem 7.3.4, there exists c such that $c \in c_v K_v^{\times 2}$ for all $v \in S$. Since c_v is in the form $ax_1^2 + bx_2^2$, so is c . Then $c = g(X_3, \dots, X_n)$ has a solution for all v .

Thus

$$h(X, X_3, \dots, X_n) := cX^2 - g(X_3, \dots, X_n)$$

represents 0 in all K_v . By the induction hypothesis, it represents 0 in K as well. Then f represents 0: if $c = ax_1^2 + bx_2^2$ then replace the solution (x, x_3, \dots, x_n) with $(xx_1, xx_2, x_3, \dots, x_n)$. This finishes the proof.

We now use Hasse-Minkowski show that most quadratic forms in $n \geq 5$ variables represent 0.

Lemma 16.2.10. *A form $f = X^2 - bX^2 - cZ^2 + acT^2$ represents every nonzero element over a local field K unless $K = \mathbb{R}$ and f is positive definite.*

A form f in $n \geq 5$ variables over K represents 0 unless K is real and f is definite.

Proof. First we show that if f does not represent 0 in K , then $a, b \notin K^{\times 2}$, $ab \in K^{\times 2}$, and $c \notin \text{Nm}_{K(\sqrt{a})} K(\sqrt{a})^\times = \text{Nm}_{K(\sqrt{b})} K(\sqrt{b})^\times$. If a or b is in $K^{\times 2}$ then f clearly represents 0, so $a, b \notin K^{\times 2}$. By $\sim (1) \implies \sim (2)$ of Theorem 16.2.9, $c \notin \text{Nm}_{K(\sqrt{a})/K}(K(\sqrt{a})^\times) \text{Nm}_{K(\sqrt{b})/K}(K(\sqrt{b})^\times)$. If $K(\sqrt{a}) \neq K(\sqrt{b})$, then the norm groups are distinct groups of index 2 in K^\times , by the correspondence between norm groups and extensions. Then their product must be all of K^\times , a contradiction. Hence, $K(\sqrt{a}) = K(\sqrt{b})$ and $ab \in K^{\times 2}$. Then $\sim (2)$ becomes simply $c \notin \text{Nm}_{K(\sqrt{a})/K}(K(\sqrt{a})^\times)$.

Conversely, suppose $a, b \notin K^{\times 2}$, $ab \in K^{\times 2}$, and $c \notin \text{Nm}_{K(\sqrt{a})} K(\sqrt{a})^\times = \text{Nm}_{K(\sqrt{b})} K(\sqrt{b})^\times$. Let $N := \text{Nm}_{K(\sqrt{a})/K}(K(\sqrt{a})^\times)$; as noted it has index 2 in K^\times . Then

$$\{x^2 - by^2 - cz^2 + act^2 : x, y, z, t \in K \text{ not all zero}\} = \{x^2 - by^2\} - c\{z^2 - at^2\} = N - cN$$

where $A \pm B$ denotes $\{a \pm b : a \in A, b \in B\}$. Since $c \notin N$, $0 \notin N - cN$. Since $N - cN$ is invariant under multiplication by elements of N , it is a union of cosets of N . Suppose that $N - cN \neq K^\times$. Then $N - cN$ is either N or cN , and

$$\{N - cN, cN - c^2N\} = \{N, cN\}$$

so

$$N - cN + cN - c^2N = N + cN.$$

If $-1 \in N$, then $N + cN = N - cN$ is cN or N , which is a contradiction because 0 is in the LHS above. Hence $-1 \notin N$. Then

$$(N - cN) - (cN - c^2N) \in \{N - cN, cN - N\} = \{N, cN\}$$

Now $c, -1 \notin N$ imply $-c \in N$, so $(N - cN) - (cN - c^2N) = N + N + N + N \in \{N, cN\}$. We have $1^2 + 1^2 + 1^2 + 1^2 = 2^2 \in N$ and $3^2 + 4^2 = 5^2$, so $N + N + N + N = N$ and $N + N = N$ (as it is a union of cosets). This implies that there exists a choice of sign in K : K^\times is the disjoint union of the closed semigroups N of “positive” elements and $-N$ of “negative” elements. If K is p -adic then this cannot happen as we must have $N \supseteq \overline{\mathbb{Z}} = K$ where $\overline{\bullet}$ denotes closure in the v -adic topology. The only possibility is $K = \mathbb{R}$. Because N consists just of positive numbers, f is positive definite. This proves the first part.

For the second part, write $f(X_1, \dots, X_n) = g(X_1, \dots, X_{n-1}) - a_n X_n^2$. By the first part, $g(X_1, \dots, X_{n-1})$ represents every element of K^\times unless $K \cong \mathbb{R}$ and g is positive definite. (Just consider $g(X_1, \dots, X_4, 0, \dots, 0)$.) In the first case, g represents a_n so f represents 0. In the second case, g represents all positive reals, and f fails to represent all reals iff a_n is negative, i.e. f is positive definite. \square

Corollary 16.2.11. *A form f in $n \geq 5$ variables represents 0 in K unless there is a real place v with f positive definite in K_v .*

Proof. This follows directly from Lemma 16.2.10 and the Hasse-Minkowski Theorem 16.2.3. \square

3 Chebotarev density theorem

Definition 16.3.1: The **density** of a set of primes S in K is d if

$$d = \lim_{N \rightarrow \infty} \frac{|\{\mathfrak{p} \in S \mid \mathfrak{N}\mathfrak{p} \leq N\}|}{|\{\mathfrak{p} \mid \mathfrak{N}\mathfrak{p} \leq N\}|}.$$

The **Dirichlet density** of a set of primes S in K is δ if

$$\delta = \lim_{s \rightarrow 1^+} \frac{\sum_{\mathfrak{p} \in S} \frac{1}{\mathfrak{N}\mathfrak{p}^s}}{\ln \frac{1}{s-1}}.$$

(Note that $\sum_{\mathfrak{p}} \frac{1}{\mathfrak{N}\mathfrak{p}^s} \sim \ln \frac{1}{s-1}$ as $s \rightarrow 1^+$ by a weak version of the prime number theorem for number fields.)

Note if a set of primes has density d , then it has Dirichlet density d (an exercise in partial summation), but a set of primes having a Dirichlet density may not have a well-defined density.

Theorem 16.3.2 (Chebotarev density theorem). *Let L/K be a finite Galois extension of number fields, and let C be a conjugacy class G . The set of prime ideals \mathfrak{p} of K such that $(\mathfrak{p}, L/K) = C$ has density $\frac{|C|}{|G|}$.*

In the special case that G is abelian, the conjugacy classes are just elements and they occur with density $\frac{1}{|G|}$. An especially notable case is the following.

Example 16.3.3 (Dirichlet): Let $n \in \mathbb{N}$ and k be relatively prime to n . Then the set

$$\{q \text{ prime} \mid q \equiv k \pmod{n}\}$$

has density $\frac{1}{\varphi(n)}$.

Indeed, Chebotarev gives that the density of q where $(q, L/K)$ is a specific element is $\frac{1}{\varphi(n)}$. By Example 11.1.6, this gives that the density of q being a specific (relatively prime) residue modulo n is $\frac{1}{\varphi(n)}$.

Example 16.3.4: If L/K is a Galois extension, then the density of primes of K splitting in L is $\frac{1}{[L:K]}$.

Indeed, a prime splits completely iff $(\mathfrak{p}, L/K) = 1$, by Proposition 11.1.3.

3.1 Proof


We prove a weaker form of the Chebotarev Density Theorem, with Dirichlet density. We will need the following.

Theorem 16.3.5 (Dirichlet’s theorem for number fields). *Let K be a number field, let H be a congruence subgroup modulo \mathfrak{m} , and let \mathfrak{K} be a class in $I_K^{\mathfrak{m}}/H$. The set of prime ideals \mathfrak{p} of K such that $\mathfrak{p} \in \mathfrak{K}$ has density $\frac{1}{[I_K^{\mathfrak{m}}:H]}$.*

Proof. See Lang [Lan94, VIII. §4] for the proof with Dirichlet density. □

In the proof below, we use “density” to mean “Dirichlet density.”

Proof of Chebotarev Density Theorem 16.3.2. We can’t deal with nonabelian extensions directly, so the idea is to reduce to the abelian case as follows. Consider L/L^σ ; this is cyclic. A prime \mathfrak{P} in L with $(\mathfrak{P}, L/K) = \sigma$ descends to a prime \mathfrak{P}' such that $(\mathfrak{P}', L/L^\sigma) = \sigma|_{L^\sigma}$. Since L/L^σ is abelian, these primes \mathfrak{P}' are characterized by a modular condition, and we can find their density using Theorem 16.3.5. Then we will relate the density of primes with $(\mathfrak{p}, L/K) = C$ to the density of primes with $(\mathfrak{P}, L/K) = \sigma$.



Let

$$S = \{ \mathfrak{p} : (\mathfrak{p}, L/K) = C \}.$$

Note that fixing $\sigma \in C$, $\mathfrak{p} \in S$ iff there exists $\mathfrak{P} \mid \mathfrak{p}$ in L such that $(\mathfrak{P}, L/K) = \sigma$.

Suppose $\sigma \in C$ has order f . Then L/L^σ is a cyclic extension of degree f . Let \mathfrak{c} be the conductor of this extension. The Artin map gives an isomorphism

$$I_{L^\sigma}^{\mathfrak{c}}/H \xrightarrow{\cong} G(L/K^\sigma)$$

for some congruence subgroup H .

Let $S_{L,\sigma}$ be those primes in L whose Frobenius element is σ :

$$S_{L,\sigma} = \{ \mathfrak{P} : (\mathfrak{P}, L/K) = \sigma \}.$$

(Note that $\bigcup_{\sigma \in C} S_{L,\sigma}$ gives all primes above those in S .) Let $S_{L'}$ be those primes in $L' := L^\sigma$ below a prime in L' :

$$S_{L'} = \{ \mathfrak{P} \cap L^\sigma : \mathfrak{P} \in S_{L,\sigma} \}.$$

We have a bijection $S_{L,\sigma} \cong S_{L'}$ by $\mathfrak{P} \mapsto \mathfrak{P} \cap L^\sigma$, because σ generates the decomposition group $D_{L/K}(\mathfrak{P})$, and $L/L^{D_{L/K}(\mathfrak{P})}$ has no splitting.

Now the density depends only on primes of degree 1 over \mathbb{Q} . Since H is a subgroup of index f in $I_{L^\sigma}^{\mathfrak{c}}$, by Theorem 16.3.5, $S_{L'}$ has density $\frac{1}{f}$.

Given \mathfrak{p} such that $(\mathfrak{p}, L/K) = C$, how many primes \mathfrak{P} above \mathfrak{p} satisfy $(\mathfrak{P}, L/K) = \sigma$? Choose \mathfrak{P}_0 above \mathfrak{p} . The primes above \mathfrak{p} are $\tau\mathfrak{P}_0$ for $\tau \in G(L/K)$. Each prime is hit $|D_{L/K}(\mathfrak{P})| = f$ times. Now we have $(\tau\mathfrak{P}_0, L/K) = \sigma$ iff

$$\tau(\mathfrak{P}_0, L/K)\tau^{-1} = \sigma.$$

The number of such τ is equal to the order of the stabilizer of the conjugation action (i.e. the number of elements commuting with τ) which is N divided by the number of elements in an orbit, i.e. $\frac{N}{|C|}$. Hence the number of \mathfrak{P} lying above \mathfrak{p} with $(\mathfrak{P}, L/K) = \sigma$ is

$$\frac{N/|C|}{f} = \frac{N}{|C|f}.$$

The density of $S_{L,\sigma}$ is $\frac{1}{f}$. Now every $\frac{N}{|C|f}$ good primes in L correspond to 1 good prime down below, so we get the desired density to be

$$\frac{1/f}{N/(|C|f)} = \frac{|C|}{N}.$$

□

3.2 Applications

Often, we will need Chebotarev just for the existence of infinitely many primes with $(\mathfrak{p}, L/K) = C$, or just for the existence of a prime after we exclude a set of zero density. Here is a typical application.

Corollary 16.3.6. *Let K be a number field. There exist infinitely many primes p of \mathbb{Q} such that there is a prime $\mathfrak{p} \mid p$ of K with $(\mathfrak{p}, L/K) = C$ and $\mathfrak{N}\mathfrak{p} = p$.*

Proof. Chebotarev's Theorem 16.3.2 says there is a positive Dirichlet density of primes \mathfrak{p} with $(\mathfrak{p}, L/K) = C$. The Dirichlet density of primes \mathfrak{p} with residue degree greater than 1 is 0, because a sum of terms of the form $\frac{1}{p^{fs}}$ with $f \geq 2$ converges. Hence infinitely many primes must remain. □

Definition 16.3.7: For two sets S, T , we write $S \lesssim T$ to mean $S \subseteq T \cup A$ for some finite set A , i.e. we have inclusion except for finitely many elements. We write $S \approx T$ if $S \lesssim T$ and $T \lesssim S$.

Definition 16.3.8: Define

$$\text{Spl}(M/K) = \{\mathfrak{p} \text{ prime of } K \text{ splitting completely in } M\}.$$

$$\widetilde{\text{Spl}}(M/K) = \{\mathfrak{p} \text{ prime of } K \text{ unramified in } M, f(\mathfrak{P}/\mathfrak{p}) = 1 \text{ for some } \mathfrak{P} \text{ in } M\}.$$

If \mathfrak{p} is unramified in K and $f(\mathfrak{P}/\mathfrak{p}) = 1$, we say that \mathfrak{P} is a **split factor** of \mathfrak{p} .

Note $\widetilde{\text{Spl}}(M/K) = \text{Spl}(M/K)$ if M/K is Galois.

The following says that the primes that split in a Galois extension characterize the extension uniquely, as well as giving inclusions between extensions.

Theorem 16.3.9. *Let L/K and M/K be finite field extensions.*

1. *If L/K is Galois, then $L \subseteq M$ iff $\widetilde{\text{Spl}}(M/K) \lesssim \text{Spl}(L/K)$.*
2. *If M/K is Galois, then $L \subseteq M$ iff $\text{Spl}(M/K) \gtrsim \text{Spl}(L/K)$.*
3. *If L/K and M/K are Galois, then $L = M$ if and only if $\text{Spl}(M/K) \approx \text{Spl}(L/K)$.*

In (1) and (2), inclusions actually hold.

Proof.

1. Suppose $L \subseteq M$, and $\mathfrak{p} \in \widetilde{\text{Spl}}(M/K)$. Say that $\mathfrak{P} \mid \mathfrak{p}$ and $f(\mathfrak{P}/\mathfrak{p}) = 1$. Let $\mathfrak{P}' = \mathfrak{P} \cap \mathcal{O}_K$. Then $f(\mathfrak{P}'/\mathfrak{p}) = 1$. Additionally, $e(\mathfrak{P}/\mathfrak{p}) = 1$ implies $e(\mathfrak{P}'/\mathfrak{p}) = 1$. Since L/K is Galois, the ramification indices and residue field degrees are equal for all primes above \mathfrak{p} . Hence $\widetilde{\text{Spl}}(M/K) \subseteq \text{Spl}(L/K)$.

Conversely suppose $\widetilde{\text{Spl}}(M/K) \lesssim \text{Spl}(L/K)$. Let N/K be a Galois extension containing L and M . It suffices to show $G(N/M) \subseteq G(N/L)$; then Galois theory gives $M \supseteq L$.

Take any $\sigma \in G(N/M)$. By Chebotarev Density 16.3.2, there exist infinitely many primes \mathfrak{p} in K such that $(\mathfrak{p}, N/K) = [\sigma]$. For such a prime \mathfrak{p} , let \mathfrak{P} be a prime lying above \mathfrak{p} in N such that $(\mathfrak{P}, N/K) = \sigma$ and let $\mathfrak{P}' = \mathfrak{P} \cap \mathcal{O}_M$. For such a prime we have

$$\alpha \equiv \sigma(\alpha) \equiv \alpha^{n_{\mathfrak{P}}} \pmod{\mathfrak{P}'}, \quad \alpha \in \mathcal{O}_M.$$

The left equality holds because σ fixes M and the right equality holds by definition of $(\mathfrak{P}, N/K)$. Hence $\mathcal{O}_M/\mathfrak{P}' \subseteq \mathbb{F}_{n_{\mathfrak{P}}} = \mathcal{O}_K/\mathfrak{p}$, and equality must hold. In other words, $f(\mathfrak{P}'/\mathfrak{p}) = 1$. Hence $\mathfrak{p} \in \widetilde{\text{Spl}}(M/K)$. Since $\widetilde{\text{Spl}}(M/K) \lesssim \text{Spl}(L/K)$, we can take \mathfrak{p} such that $\mathfrak{p} \in \text{Spl}(L/K)$ as well. Then $\sigma|_L = 1$ hence $G(N/M) \subseteq G(N/L)$ and $M \supseteq L$.

2. Suppose $L \subseteq M$. Then any prime splitting completely in M splits completely in L , so $\text{Spl}(M/K) \subseteq \text{Spl}(L/K)$.

Conversely suppose $\text{Spl}(M/K) \gtrsim \text{Spl}(L/K)$. Let L^{gal} be the Galois closure of L . Since M/K is Galois, $\widetilde{\text{Spl}}(M/K) = \text{Spl}(M/K)$; we also have $\text{Spl}(L/K) = \text{Spl}(L^{\text{gal}}/K)$ (Any prime splitting completely in L splits completely in the Galois closure, by exercise 2 in 8). Thus

$$\widetilde{\text{Spl}}(M/K) \subseteq \text{Spl}(L^{\text{gal}}/K)$$

and we can apply part 1 to get $L^{\text{gal}} \subseteq M$; *a fortiori* $L \subseteq M$.

3. Apply part 2 twice. □

4 Splitting of primes

4.1 Splitting of primes

Theorem 16.4.1. *Let L/K be an extension of number fields.*

1. *If L^{gal}/K is abelian, then there is a modulus \mathfrak{m} and a congruence subgroup modulo \mathfrak{m} such that*

$$\text{Spl}(L/K) = \{\text{prime } \mathfrak{p} \in H\}.$$

2. *If there exists $\mathfrak{K} \in C_K(\mathfrak{m}) = I_K^{\mathfrak{m}}/P_K(1, \mathfrak{m})$ such that*

$$\{\text{prime } \mathfrak{p} : \mathfrak{p} \pmod{P_K(1, \mathfrak{m})} = \mathfrak{K}\} \simeq \text{Spl}(L/K),$$

(i.e. all but finitely many primes satisfying a certain modular condition split) then L^{gal}/K is abelian.

In other words the law of decomposition of primes in an extension L/K is determined by modular conditions iff L/K is an abelian extension.

Proof. ³ As $\text{Spl}(L/K) = \text{Spl}(L^{\text{gal}}/K)$, it suffices to consider L/K Galois.

Part 1: By global class field theory, the kernel of the Artin map $I_K^{\mathfrak{m}} \rightarrow G(L/K)$ is a congruence subgroup H . But we have by Proposition 11.1.3 that \mathfrak{p} splits completely iff $\psi_{L/K}(\mathfrak{p}) = (\mathfrak{p}, L/K) = 1$. Hence

$$H = \ker(\psi_{L/K}) = \text{Spl}(L/K).$$

Part 2: Let $K_{\mathfrak{m}}$ be the ray class field of K modulo \mathfrak{m} and $M = LK_{\mathfrak{m}}$. There is a natural map

$$p = p_1 \times p_2 : G(M/K) \hookrightarrow G(K_{\mathfrak{m}}/K) \times G(L/K) \xrightarrow{\cong} C_K(\mathfrak{m}) \times G(L/K)$$

where the second map is given by $\psi_{L/K}^{-1}$ in the first component.

For all but finitely many primes, we have the following string of facts.

1. $\mathfrak{p} \in \mathfrak{K}$.
2. $\mathfrak{p} \in \text{Spl}(L/K)$.
3. $(\mathfrak{p}, L/K) = 1$.
4. For any prime $\mathfrak{P} \mid \mathfrak{p}$ in M , $p((\mathfrak{P}, M/K)) = (\mathfrak{K}, 1)$.

³This proof is from <http://mathoverflow.net/questions/11688>.

(1) \implies (2) is by assumption, (2) \iff (3) is Proposition 11.1.3, and (3) \iff (4) is by compatibility of the Frobenius elements (the map $G(M/K) \rightarrow G(L_{\mathfrak{m}}/K) \times G(L/K)$ is compatible with the map on residue fields $G(m/k) \rightarrow G(k_{\mathfrak{m}}/k) \times G(l/k)$).

Suppose $\sigma \in G(M/K)$ and $p(\sigma) = (\mathfrak{K}, g)$. By Chebotarev's Theorem there exist primes $\mathfrak{P} \mid \mathfrak{p}$ in M and K , respectively, such that $(\mathfrak{P}, M/K) = \sigma$. But (1) \implies (4) shows that $g = 1$. Hence

$$p(G(M/K)) \cap (\mathfrak{K}, G(L/K)) = \{(\mathfrak{K}, 1)\}.$$

Since p is a group homomorphism that is surjective in the first component, $p(G(M/K)) \cap (\mathfrak{K}', G(L/K))$ must consist of 1 element for every \mathfrak{K}' , in particular for $\mathfrak{K}' = 1$. Thus if $p(\sigma) = (P_K(1, \mathfrak{m}), g)$, then $g = 1$. Given a prime \mathfrak{p} splitting completely in $K_{\mathfrak{m}}$, i.e. \mathfrak{p} such that $\mathfrak{p} \in P_K(1, \mathfrak{m})$, take any $\mathfrak{P} \mid \mathfrak{p}$ in M . Then $p(\mathfrak{P}, M/K) = (P_K(1, \mathfrak{m}), g)$ for some g , so $g = 1$ and

$$(\mathfrak{p}, L/K) = (\mathfrak{P}, M/K)|_L = p_2(\mathfrak{P}, M/K) = g = 1,$$

i.e. \mathfrak{p} splits completely in L . Thus $\text{Spl}(L_{\mathfrak{m}}/K) \stackrel{\sim}{\subset} \text{Spl}(L/K)$, showing by Theorem 16.3.9 that $L \subseteq L_{\mathfrak{m}}$. \square

For nonabelian extensions, the set of primes that split has to be specified by more than just a modulo condition.

Example 16.4.2: We show that a prime splits completely in $\mathbb{Q}(\zeta_3, \sqrt[3]{2})$ iff $p \equiv 1 \pmod{3}$ and p is in the form $x^2 + 27y^2$.

Note that $\mathbb{Q}(\zeta_3, \sqrt[3]{2})$ is the splitting field of $x^3 - 2 = 0$. For an unramified prime, p splits completely iff the residue field extension has degree 1, i.e. $x^3 - 2$ splits completely in \mathbb{F}_p . This is true iff 2 is a cubic residue modulo p . As we saw in Theorem 16.1.14, this is true iff p is of the form $x^2 + 27y^2$.

4.2 Roots of polynomials over finite fields

We can recast the problem of splitting behavior in terms of finding roots of univariate polynomials over finite fields. Let L/K be a finite extension, and $f \in \mathcal{O}_K[X]$ be the minimal polynomial of a primitive element in L/K . Then Theorem 2.6.3 tells us that for a prime \mathfrak{p} relatively prime to the conductor of L/K , the factorization of f in $\mathcal{O}_K/\mathfrak{p}$ corresponds to the factorization of \mathfrak{p} . In particular, \mathfrak{p} splits completely iff f splits completely, and \mathfrak{p} has a split factor iff f has a root in $\mathcal{O}_K/\mathfrak{p}$.

Definition 16.4.3: Let $N_{\mathfrak{p}}(f)$ denote the number of zeros of f in $\mathcal{O}_K/\mathfrak{p}$.

Thus we can rephrase Theorem 16.4.1 as follows.

Theorem 16.4.4. *Let f be an irreducible polynomial over K . Let α be a root of f and L be the Galois closure of $K(\alpha)$.*

1. *For all except a finite number of primes, $N_{\mathfrak{p}}(f) = m$ iff $\psi_{L/K}(\mathfrak{p}) = [\sigma]$ for some $\sigma \in G(L/K)$ fixes m of the roots of L .*

2. The sets $\{\mathfrak{p} : N_{\mathfrak{p}}(f) = m\}$ are given by modular conditions iff L/K is abelian.

3. The density of primes \mathfrak{p} such that $N_{\mathfrak{p}}(f) = m$ is $\frac{\{\sigma \in G(L/K) : \sigma \text{ fixes } m \text{ roots}\}}{[L:K]}$.

Proof. The first item follows from Theorem 2.6.3. The second item follows from this and Theorem 16.4.1. The third item follows from the Chebotarev Density Theorem 16.3.2. \square

Even the reciprocity laws (at least, weak reciprocity) can be put in the same framework: in a field K containing n th roots of unity, a is a perfect n th power modulo \mathfrak{p} iff $x^n - a$ splits completely modulo \mathfrak{p} (the polynomial viewpoint), i.e. the prime \mathfrak{p} splits completely in $K(\sqrt[n]{a})/K$ (the splitting viewpoint).

5 Hilbert class field

Definition 16.5.1: The **Hilbert class field** of K is the largest abelian field extension of K unramified over K at all places. (For infinite places this means that no real embedding becomes complex.) It is denoted H_K .

The **large Hilbert class field** of K is the largest abelian field extension of K unramified over K at all finite places, with no restrictions for infinite places (i.e. they are allowed to ramify). It is denoted H_K^+ .

Note if K is already totally complex then $H_K = H_K^+$.

Proposition 16.5.2: The Hilbert class field and large Hilbert class field exist, and the global reciprocity map gives isomorphisms

$$\begin{aligned} G(H_K/K) &\cong \text{Cl}_K \\ G(H_K^+/K) &\cong \text{Cl}_K^+. \end{aligned}$$

Proof. The Hilbert class field is exactly the ray class field corresponding to the modulus 1, and the narrow Hilbert class field is exactly the ray class field corresponding to the modulus $\mathfrak{m} = \prod_{v \text{ real}} v$. Indeed, by global class field theory the fields corresponding to congruence subgroups of $C_K(1)$ are just the fields unramified over K , and the fields corresponding to congruence subgroups of $C_K(\mathfrak{m})$ are just the fields unramified at every infinite place.

The global reciprocity map gives the desired isomorphisms. \square

The most interesting property of the Hilbert class field is the following.

Theorem 16.5.3. *Let K be a global field. Every fractional ideal of K becomes principal in the Hilbert class field L of K .*

Proof. Let M be the Hilbert class field of L . By Proposition 16.5.2, the global reciprocity map gives $C_K \xrightarrow{\cong} G(L/K)$ and $C_L \xrightarrow{\cong} G(M/L)$. We will transfer the map $C_K \rightarrow C_L$ to the Galois groups. By definition, L is the maximal unramified *abelian* extension of K ; since M is also unramified over K , L is the maximal *abelian* subextension of M/K . But by Galois

theory, intermediate Galois extensions correspond to quotient groups of $G(M/K)$. This means that

$$G(L/K) = G(M/K)/G(M/L)$$

is the largest abelian quotient of $G(L/K)$. From group theory this means that $G(M/L)$ is the *derived subgroup* $(G(L/K))'$.

The following diagram commutes by compatibility of the Artin map (the last diagram in Theorem 14.4.10 together with Theorem 15.5.1)

$$\begin{array}{ccc} C_K & \xrightarrow[\cong]{\phi_{L/K}} & G(L/K)^{\text{ab}} \\ \downarrow & & \downarrow V \\ C_L & \xrightarrow[\cong]{\phi_{M/L}} & G(M/L)^{\text{ab}} \end{array}$$

where V is the transfer.

However, the transfer map is 0 by Theorem 12.11.13 and the fact that $G(M/L) = G(L/K)'$. Hence the map $C_K \rightarrow C_L$ is trivial, i.e. every fractional ideal of K becomes trivial in L . \square

6 Primes represented by quadratic forms

We now give a complete characterization of which primes can be represented by which binary (positive definite integral) quadratic forms. First consider the form $x^2 + ny^2$.

A prime is in the form $p = x^2 + ny^2$ iff p splits as $\mathfrak{p}\bar{\mathfrak{p}} = (x + y\sqrt{-n})(x - y\sqrt{-n})$ in $\mathbb{Z}[\sqrt{-n}]$, with its factors being principal ideals. We can think of this as saying that \mathfrak{p} goes to 0 in the ideal class group of $\mathbb{Z}[\sqrt{-n}]$. Unfortunately, this is not the same class group as C_K . However, this class group is essentially a quotient of a ray class group (Theorem 4.6.2). But by class field theory, we can find a field extension L such that the Artin map to $G(L/K)$ is an isomorphism. The primes in the kernel of the Artin map are exactly those that split completely in L , so this relates the equation $x^2 + ny^2$ to the splitting of primes in the Hilbert class field.

Definition 16.6.1: Let \mathcal{O} be an integral quadratic order and $f := \text{disc}(\mathcal{O})$.

1. Suppose $f < 0$. The field L corresponding to the congruence subgroup

$$P_K(\mathbb{Z}, f) := \{(a) \in I_K(f) : a \pmod{f} \in \mathbb{Z} \pmod{f}\} \subseteq I_K(f)$$

is called the **ring class field** of \mathcal{O} .

2. Suppose $f > 0$. The field L corresponding to the congruence subgroup

$$P_K(\mathbb{Z}, \infty f) := \{(a) \in I_K(f) : a \pmod{f} \in \mathbb{Z} \pmod{f}\} \subseteq I_K(f)$$

is called the **ring class field** of \mathcal{O} .

The reason for this definition is that $I_K(f)/P_K(\mathbb{Z}, \infty f) \cong I(\mathcal{O})/P^+(\mathcal{O}) = C^+(\mathcal{O})$ via the map $\mathfrak{a} \mapsto \mathfrak{a} \cap \mathcal{O}$, by Theorem 4.6.2. (Ignore the ∞ when K is imaginary; in this case $C^+(\mathcal{O}) = C(\mathcal{O})$.)

Example 16.6.2: When $\mathcal{O} = \mathcal{O}_K$, with K/\mathbb{Q} a quadratic extension, then the ring class field is just the large Hilbert class field of K , because $I(\mathcal{O})/P^+(\mathcal{O}) = C_K^+$.

Theorem 16.6.3. *Let $n \geq 1$. Let Q be a quadratic form that corresponds to $\mathfrak{a} \subseteq R$ under the Gauss correspondence 4.5.1, let $K = \text{Frac}(R)$, and let p be an odd prime not dividing $f := \text{disc}(R)$. Let \mathfrak{b} be the ideal corresponding to \mathfrak{a} under the map $I_K(f)/P_K(\mathbb{Z}, \infty f) \rightarrow I(\mathcal{O})/P^+(\mathcal{O}) = C^+(\mathcal{O})$. Let L be the ring class field of R and suppose $(L/K, \mathfrak{b}) = \sigma$. Then*

$$f \text{ represents } \mathfrak{p} \iff (L/\mathbb{Q}, p) = [\sigma]$$

where $[\sigma]$ denotes the conjugacy class of σ in $G(L/\mathbb{Q})$.

Proof. Let $K = \mathbb{Q}(\sqrt{-n})$. We have the following string of equivalences.

1. Q represents p .
2. $pR = \mathfrak{p}\bar{\mathfrak{p}}$ in R for some ideal \mathfrak{p} in the same ideal class as \mathfrak{a} .
3. $p\mathcal{O}_K = \mathfrak{p}\bar{\mathfrak{p}}$ for $\mathfrak{p} \sim \mathfrak{b}$ where the ideals are considered in $I_K(f)/P_K(\mathbb{Z}, \infty f)$.
4. $p\mathcal{O}_K = \mathfrak{p}\bar{\mathfrak{p}}$ for $(L/K, \mathfrak{p}) = \sigma$.
5. $(L/\mathbb{Q}, p) = [\sigma]$.

The equivalence (1) \iff (2) follows from Proposition 4.5.4. We have (2) \iff (3) by Theorem 4.6.2, which gives an isomorphism $I_K(f)/P_K(\mathbb{Z}, \infty f) \rightarrow I(\mathcal{O})/P^+(\mathcal{O}) = C^+(\mathcal{O})$ by sending \mathfrak{a} to $\mathfrak{a} \cap \mathcal{O}$. By definition of ring class field, the Artin map is an isomorphism $I_K(f)/P_K(\mathbb{Z}, \infty f) \rightarrow G(L/K)$, so (3) \iff (4).

For (4) \iff (5), note by definition of the Artin symbol that (4) is equivalent to

$$p \text{ splits in } \mathcal{O}_K \text{ and } \sigma(\alpha) \equiv \alpha^{|k|} \pmod{\mathfrak{P}} \text{ for all } \alpha \in L$$

where \mathfrak{P} is any prime dividing \mathfrak{p} in L . Since p is unramified, p splits in \mathcal{O}_K iff $[k : \mathbb{F}_p] = 1$, iff $|k| = p$. Hence the above is equivalent to

$$\sigma(\alpha) \equiv \alpha^p \pmod{\mathfrak{P}}$$

This says exactly that $(L/\mathbb{Q}, p) = [\sigma]$. □

Corollary 16.6.4. *Suppose $n \neq 0$ is an integer.*

1. *Let L be the ring class field of $\mathbb{Z}[\sqrt{-n}]$. Then p can be represented as*

$$p = x^2 + ny^2, \quad x, y \in \mathbb{Z}$$

if and only if p splits completely in L .

2. For $-n \equiv 1 \pmod{4}$, let L' be the ring class field of $\mathbb{Z}\left[\frac{1+\sqrt{-n}}{2}\right]$. Then p can be represented as

$$p = x^2 + xy + \frac{1-n}{2}y^2$$

iff p splits completely in L' .

Remark 16.6.5: It is not hard to show that we can replace the conditions by the following uniform statement: $4p$ can be represented as $4p = x^2 + dy^2$ iff p splits completely in the order of discriminant $-d$.

Proof. These quadratic forms correspond to the principal ideals in $\mathbb{Z}[\sqrt{-n}]$ and $\mathbb{Z}\left[\frac{1+\sqrt{-n}}{2}\right]$, respectively (Example 4.5.3), so the theorem says p can be represented by the quadratic forms iff

$$(L/K, p) = 1.$$

This is true iff \mathfrak{p} splits completely in L (Proposition 11.1.3). \square

How is this useful? Algorithmically, there are fast ways to find solutions to $p = x^2 + ny^2$ (Cornacchia's algorithm), so we can obtain primes splitting completely in the Hilbert class field H_K . This means that the minimal polynomial of H_K/K factors completely modulo p . As we will in Chapter 17, the roots are the j -invariants of CM elliptic curves; the fact that they are in \mathbb{F}_p gives us an easy way to calculate the action of the class group on elliptic curves.

Additionally, this description of solutions to $p = x^2 + ny^2$ gives a way to find the density of primes represented by a quadratic form.

Theorem 16.6.6. Let Q be a primitive positive definite quadratic form of discriminant $D < 0$, and let S be the set of primes represented by Q . Then the density of primes $d(S)$ represented by S is

$$d(S) = \begin{cases} \frac{1}{2h(D)}, & Q \text{ properly equivalent to its opposite,} \\ \frac{1}{h(D)}, & \text{else,} \end{cases}$$

where $h(D)$ is the class number of the quadratic ring with discriminant D . In particular, Q represents infinitely many prime numbers.

Note “ Q properly equivalent to its opposite” is equivalent to saying that the ideal class corresponding to Q has order dividing 2.

Example 16.6.7: $h(-27) = 3$ so $\frac{1}{6}$ of all primes can be represented by the form $x^2 + 27y^2$.

In fact, the ring class field of $\mathbb{Z}[\sqrt{-27}]$ is $\mathbb{Q}(\zeta_3, \sqrt[3]{2})$, so $p = x^2 + 27y^2$ iff p splits completely in $\mathbb{Q}(\zeta_3, \sqrt[3]{2})$. This shows Example 16.4.2 in a different way.

Proof of Theorem 16.6.6. Let K be the quadratic field of discriminant D .

By Theorem 16.6.3, p is represented by Q iff $(L/\mathbb{Q}, p) = [\sigma]$ where L is the ring class field of the order corresponding to Q and Q corresponds to σ under the Gauss correspondence. We need to find $[\sigma]$, so we first need to understand $G(L/\mathbb{Q})$.

Since $C(\mathcal{O}) \cong I_K(f)/P_K(\mathbb{Z}, f) \cong G(L/K)$ via the Artin map,

$$[L : K] = |C(\mathcal{O})| = h(D) \implies [L : \mathbb{Q}] = 2h(D).$$

Next we show $G(L/\mathbb{Q}) = G(L/K) \rtimes G(K/\mathbb{Q})$ where, denoting complex conjugation by $\sigma \in G(K/\mathbb{Q})$, we have $\sigma\tau\sigma^{-1} = \tau^{-1}$ for all $\tau \in G(L/K)$. Let \mathfrak{m} be the modulus corresponding to $f\mathcal{O}_K$, where f is the conductor. By construction of L , it is the unique field such that $\ker(\psi_{L/K}) = P_K(\mathbb{Z}, f)$. However, because the Artin map commutes with Galois action (see the third diagram in Theorem 14.4.10),

$$\ker(\psi_{\sigma(L)/K}) = \sigma \ker(\psi_{L/K}) = \sigma P_K(\mathbb{Z}, f) = P_K(\mathbb{Z}, f).$$

Uniqueness hence gives $\sigma(L) = L$, i.e. $\sigma \in L$. Hence $|G(L/\mathbb{Q})| = 2|G(L/K)| = [L : \mathbb{Q}]$, giving that L/\mathbb{Q} is Galois. Given $\tau \in G(L/K)$, by surjectivity of the Frobenius map 15.2.8, $\tau = (L/K, \mathfrak{p})$ for some \mathfrak{p} . Then by Lemma 11.1.2,

$$\sigma\tau\sigma^{-1} = \sigma(L/K, \mathfrak{p})\sigma^{-1} = (L/K, \sigma\mathfrak{p}) = (L/K, \bar{\mathfrak{p}}) = (L/K, \mathfrak{p})^{-1} = \tau^{-1},$$

as needed.

From the structure of $G(L/\mathbb{Q})$, we see that the conjugacy class of any element σ is $\{\sigma, \sigma^{-1}\}$. By the Chebotarev density theorem 16.3.2, the density of primes such that $(L/\mathbb{Q}, p) = [\sigma] = \{\sigma, \sigma^{-1}\}$ is hence

$$\frac{|[\sigma]|}{[L : \mathbb{Q}]} = \begin{cases} \frac{1}{2h(D)}, & \sigma = \sigma^{-1}, \\ \frac{1}{h(D)}, & \text{else,} \end{cases}$$

as needed. □

7 Introduction to the Langlands program

In this section, we'll give the big picture, and be content with morally, rather than mathematically correct, statements.

Much of modern number theory is occupied with the relationship between the following three objects.

1. Algebraic varieties, i.e. polynomial equations.
2. Galois representations, i.e. continuous functions from $G(\bar{K}/K)$ to algebraic groups such as $GL_n(\mathbb{C})$.
3. Automorphic forms, i.e. continuous functions defined on an algebraic group on the ideles, such as $GL_n(\mathbb{A}_K)$, and satisfying certain conditions.

The relationship between Galois representations and automorphic forms is known as the Langlands correspondence. More precisely, there is a conjectural correspondence

$$\left\{ \begin{array}{l} \text{cuspidal automorphic} \\ \text{representations of } \mathrm{GL}_n(\mathbb{A}_K) \\ \text{algebraic at } \infty \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} \text{irreducible continuous} \\ G(\overline{K}/K) \rightarrow \mathrm{GL}_n(\mathbb{C}) \\ \text{algebraic at } \ell \end{array} \right\}$$

We can define L -series from both Galois representations and automorphic forms. L -series from Galois representations arise more naturally in number theory (because it is relatively easy to go from algebraic varieties to Galois representations), but as automorphic forms are analytic objects, L -series of automorphic forms are known to satisfy more properties. The Langlands correspondence allows us to show that L -series of Galois representations arise from automorphic forms, hence have nice analytic properties as well. This allows us to prove various results about algebraic varieties, such as density theorems on the number of solutions over finite fields, for example the Sato-Tate conjecture.

We first give some more precise definitions, then describe this relationship in the 1-dimensional abelian case (which we have in fact proved!), and then give an overview of how it generalizes.

7.1 Definitions

Definition 16.7.1: Let k be a topological field (for instance, \mathbb{C} or \mathbb{Q}_ℓ), and let $V \cong k^n$ be a n -dimensional vector space over k . A n -dimensional **Galois representation** of K over k is a continuous homomorphism

$$\rho : G(K^s/K) \rightarrow \mathrm{GL}(V) = \mathrm{GL}_n(k).$$

Let \mathfrak{p} be a prime of K . We say ρ is **unramified** at \mathfrak{p} if $I_{\mathfrak{p}}(K^s/K) \subseteq \ker(\rho)$.

Let K be a number field. Let $\mathrm{Frob}(\mathfrak{p})$ be a Frobenius element of \mathfrak{p} in $K_{\mathfrak{p}}$ (defined in $G(\overline{K}_{\mathfrak{p}}/K_{\mathfrak{p}})$ up to $I(\overline{K}_{\mathfrak{p}}/K_{\mathfrak{p}})$). Define the (modified) **characteristic polynomial** of ρ at \mathfrak{p} to be

$$P_{\rho}(X) := \det(1 - X \cdot \rho(\mathrm{Frob}(\mathfrak{p}))|V^{I(\overline{K}_{\mathfrak{p}}/K_{\mathfrak{p}})}).$$

(Here, $V^{I(\overline{K}_{\mathfrak{p}}/K_{\mathfrak{p}})}$ denotes the subspace of V fixed by the inertia group. $P_{\rho}(X)$ is well-defined because $\mathrm{Frob}(\mathfrak{p})$ is defined up to $I(\overline{K}_{\mathfrak{p}}/K_{\mathfrak{p}})$, and $I(\overline{K}_{\mathfrak{p}}/K_{\mathfrak{p}}) \subseteq \ker(\rho|_{V^{I(\overline{K}_{\mathfrak{p}}/K_{\mathfrak{p}})}})$. In particular, if ρ is unramified at \mathfrak{p} , then $V = V^{I(\overline{K}_{\mathfrak{p}}/K_{\mathfrak{p}})}$.)

We can now define the L -function associated to a Galois representation.

Definition 16.7.2: In the above, suppose V is a complex vector space and K is a number field. The **local L -factor** at a prime \mathfrak{p} is

$$L_{\mathfrak{p}}(\rho, s) = P_{\rho}(\mathfrak{N}\mathfrak{p}^{-s})^{-1}.$$

The **Artin L -function** of ρ is⁴

$$L(\rho, s) = \prod_{\mathfrak{p}} L_{\mathfrak{p}}(\rho, s).$$

We have the following conjecture.

Conjecture 16.7.3 (Artin’s conjecture). *Every Artin L -function has analytic continuation to \mathbb{C} and satisfies a functional equation.*

7.2 Class field theory is 1-dimensional Langlands

For a different take on some of these ideas, with concrete examples, see Dalawat [Dal11].

7.2.1 Galois representations are automorphic representations

We rephrase global class field theory in the form that generalizes under the Langlands program.

Theorem 16.7.4 (Rephrase of GCFT). *There is a bijection between continuous homomorphisms $\chi : \mathbb{A}_K^\times / \overline{K^\times (K_\infty^\times)^0} \rightarrow \mathbb{C}^\times$ and continuous homomorphisms $\rho : G(\overline{K}/K) \rightarrow \mathrm{GL}_1(\mathbb{C})$, given by the following.*

$$\begin{aligned} \{\chi : \mathbb{A}_K^\times / \overline{K^\times (K_\infty^\times)^0} \rightarrow \mathbb{C}^\times\} &\leftrightarrow \{\rho : G(\overline{K}/K) \rightarrow \mathrm{GL}_1(\mathbb{C})\} \\ \chi &\mapsto \chi \circ \phi_K^{-1} \end{aligned}$$

Proof. From Theorem 11.6.5, the Artin map gives a topological isomorphism $\mathbb{A}_K^\times / \overline{K^\times (K_\infty^\times)^0} \rightarrow G(K^{\mathrm{ab}}/K)$. It remains to note that any function $G(\overline{K}/K) \rightarrow \mathrm{GL}_1(\mathbb{C})$ factors through $G(\overline{K}/K)^{\mathrm{ab}} = G(K^{\mathrm{ab}}/K)$, since $\mathrm{GL}_1(\mathbb{C})$ is abelian. \square

The functions on the left side have a special name.

Definition 16.7.5: A **Hecke character** is a continuous homomorphism $\mathbb{A}_K^\times / \overline{K^\times (K_\infty^\times)^0} \rightarrow \mathbb{C}^\times$, or equivalently, a homomorphism

$$\chi : \mathbf{C}_K \rightarrow S^1 := \{x \in \mathbb{C} : |x| = 1\}$$

with finite image. The **conductor** of χ is the smallest modulus \mathfrak{m} such that χ factors through $\mathbb{A}_K^\times / K^\times \mathbf{U}_K(1, \mathfrak{m}) \cong C_K(\mathfrak{m})$.

The homomorphisms $\chi : \mathbb{A}_K^\times / \overline{K^\times (K_\infty^\times)^0} \rightarrow \mathbb{C}$ are “automorphic functions” on $\mathrm{GL}_1(\mathbb{A}_K)$, a.k.a. Hecke characters, and the homomorphisms $\rho : G(\overline{K}/K) \rightarrow \mathrm{GL}_1(\mathbb{C})$ are 1-dimensional “Galois representations.” Our correspondence is unsatisfactory, however, because we would like to get all continuous homomorphisms $\mathbb{A}_K^\times / K^\times \rightarrow \mathbb{C}^\times$, not just those factoring through

⁴Sometimes infinite places are included. The factors at infinite places take more thought to define so we exclude them here.

$\mathbb{A}_K^\times / K^\times (K_\infty^\times)^0$. Since $G(K^{\text{ab}}/K)$ has the profinite topology, any continuous homomorphism $G(\overline{K}/K) \rightarrow \text{GL}_1(\mathbb{C})$ must have finite image, while functions $\mathbb{A}_K^\times / K^\times \rightarrow \mathbb{C}^\times$ can have infinite image. To remedy this, we introduce functions $G(\overline{K}/K) \rightarrow \text{GL}_1(\mathbb{C})$ with infinite image (no longer continuous under the complex topology).

For simplicity, we just consider the case of \mathbb{Q} .

Example 16.7.6: We say a function $\pi : \mathbb{A}_\mathbb{Q}^\times / \mathbb{Q}^\times \rightarrow \mathbb{C}$ is **algebraic at ∞** if $\pi(i_\mathbb{R}(x)) = \text{sign}(x)^m |x|^n$ for some $m \in \{0, 1\}$ and $n \in \mathbb{Z}$. We characterize all the continuous homomorphisms $\pi : \mathbb{A}_K^\times / K^\times \rightarrow \mathbb{C}^\times$ (“**Größencharacters**”) that are algebraic at ∞ .

It is enough to introduce 1 more character. Let ℓ be a prime of \mathbb{Q} . Let $|\cdot| : \mathbb{A}_\mathbb{Q}^\times / \mathbb{Q}^\times \rightarrow \mathbb{C}^\times$ denote the map $|\mathbf{x}| = \prod_{v \in V_\mathbb{Q}} |x_v|_v$, and define χ_ℓ by

$$\chi_\ell : G(\overline{\mathbb{Q}}/\mathbb{Q}) \twoheadrightarrow G(\mathbb{Q}^{\text{ab}}/\mathbb{Q}) = G(\mathbb{Q}(\zeta_\infty)/\mathbb{Q}) \xrightarrow{\cong} \widehat{\mathbb{Z}}^\times = \prod_p \mathbb{Z}_p^\times \twoheadrightarrow \text{GL}_1(\mathbb{Z}_\ell).$$

(We say χ_ℓ is “**algebraic at ℓ** .” Note there is a noncanonical field isomorphism $\overline{\mathbb{Q}}_\ell \cong \mathbb{C}$, so we can think of $\text{GL}_1(\mathbb{Z}_\ell)$ as being “inside” $\text{GL}_1(\mathbb{C})$.)

Every continuous homomorphism $\pi : \mathbb{A}_K^\times / K^\times \rightarrow \mathbb{C}^\times$ algebraic at ∞ is in the form $|\cdot|^n \cdot \chi$, where χ is a Hecke character. We can extend the correspondence in Theorem 16.7.4 by associating $|\cdot|$ with χ_ℓ :

$$\pi = |\cdot|^n \cdot \chi \leftrightarrow \chi_\ell^n \cdot (\chi \circ \phi_K^{-1})$$

where the right-hand side is now viewed in \mathbb{Q}_ℓ instead of \mathbb{C} .

7.2.2 Artin L -functions are Hecke L -functions

Associated to each Hecke character is a L -function.

Definition 16.7.7: Let χ be a Hecke character and \mathfrak{m} be the conductor of χ . The **L -function** associated to χ is

$$L(\chi, s) := \prod_{\mathfrak{p} \nmid \mathfrak{m}} \frac{1}{1 - \chi(\mathfrak{p}) \mathfrak{N}\mathfrak{p}^{-s}}.$$

Because χ admits a modulus, Hecke L -series have nice analytic properties.

Theorem 16.7.8 (Hecke, Tate). *Every Hecke L -series admits an analytic continuation to \mathbb{C} and satisfies a functional equation.*

For the details, see Tate’s thesis in [CF69].

Theorem 16.7.9. *Any 1-dimensional Artin L -function is a Hecke L -function. Hence it has analytic continuation and satisfies a functional equation.*

Proof. Let $\rho : G(\overline{K}/K) \rightarrow \mathrm{GL}_1(\mathbb{C})$ be a 1-dimensional representation. By Theorem 16.7.4, $\rho(\Phi_{\mathfrak{p}}) = \chi(\mathfrak{p})$ for some Hecke character $\chi : \mathbb{A}_K^\times/K^\times \rightarrow \mathbb{C}^\times$. Let \mathfrak{m} be the modulus of ρ ; note it is also the conductor for χ . Then

$$L(\rho, s) = \prod_{\mathfrak{p} \nmid \mathfrak{m}} \frac{1}{1 - \rho(\Phi_{\mathfrak{p}})\mathfrak{N}\mathfrak{p}^{-s}} = \prod_{\mathfrak{p} \nmid \mathfrak{m}} \frac{1}{1 - \chi(\mathfrak{p})\mathfrak{N}\mathfrak{p}^{-s}} = L(\chi, s).$$

□

This theorem is another way of saying that the Artin map factors through a modulus, and this is basically what allowed us to get all the density results in this chapter.

7.2.3 Algebraic varieties and Galois representations

We give examples of how to get Galois representations from algebraic varieties.

First consider the variety $\overline{\mathbb{Q}}^\times = \{x \in \overline{\mathbb{Q}} : x \neq 0\}$. It is a group under multiplication, and the torsion points $\overline{\mathbb{Q}}^\times[m]$ are exactly the roots of unity μ_m . We can define a Galois representation by considering the action of $G(\overline{\mathbb{Q}}/\mathbb{Q})$ on the l -power roots of unity. Define the Tate module of $\overline{\mathbb{Q}}^\times$ by

$$T_\ell(\overline{\mathbb{Q}}^\times) = \varprojlim_n \overline{\mathbb{Q}}^\times[\ell^n] = \varprojlim_n \mu_{\ell^n} \cong \mathbb{Z}_\ell.$$

Then $G(\overline{\mathbb{Q}}/\mathbb{Q})$ acts naturally on $T_\ell(\overline{\mathbb{Q}}^\times)$ so we get a representation

$$\rho : G(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \mathrm{Aut}(T_\ell(\overline{\mathbb{Q}}^\times)) \cong \mathrm{Aut}(\mathbb{Z}_\ell) \hookrightarrow \mathrm{GL}_1(\mathbb{Q}_\ell)$$

sending the element $\phi_{\mathbb{Q}}(p)$ to p . The corresponding L -function is just a translate of the ζ function, missing the factor ℓ : $\prod_{p \neq \ell} \frac{1}{1-p^{1-s}}$. This construction is a good analogy for what we will eventually do with elliptic curves, although it is a bit too “trivial” to capture any significant number theory facts.

We give another example, with equations in 1 variable, which is a bit less natural but show more of the number theory. Consider the variety defined by $f(X) = 0$ where $f \in K[X]$ is a irreducible polynomial. Let α be a root, and L be the Galois closure of $K(\alpha)$ over K . Let $\alpha_1, \dots, \alpha_n$ be the roots of f in L . $G(\overline{K}/K)$ acts by permuting the α_i , so we get a representation $G(\overline{K}/K) \rightarrow S_n$. We can embed S_n in some general linear group, to get $\rho : G(\overline{K}/K) \rightarrow \mathrm{GL}_m(k)$ for some k . Then to find how many roots f has modulo \mathfrak{p} , we can look at the trace of $\rho(\mathrm{Frob}(\mathfrak{p}))$.

For example, consider $f(X) = X^3 - X - 1$ over \mathbb{Q} . We get a representation $\rho : G(\overline{K}/K) \rightarrow S_3 \rightarrow \mathrm{GL}_2(\mathbb{C})$, where we embed $S_3 \hookrightarrow \mathrm{GL}_2(\mathbb{C})$ as follows: we have a natural permutation representation $S_3 \hookrightarrow \mathrm{GL}_3(\mathbb{C})$; now take out the trivial representation to get $S_3 \hookrightarrow \mathrm{GL}_2(\mathbb{C})$. From this description we have $N_p(f) = \mathrm{tr}(\rho(\mathrm{Frob}(\mathfrak{p}))) + 1$, so we can get the number of solutions of $X^3 - X - 1 \equiv 0 \pmod{p}$ from looking at the trace of Frobenius. Constructing the L -function, the trace of Frobenius becomes the coefficient of $\frac{1}{p^s}$. Now ρ comes from an automorphic form, so L comes from a 2-dimensional automorphic form, i.e. a modular

form. We can write this modular form explicitly using theta functions or as an eta quotient. At the end of the day, we have this striking fact: For $p \neq 23$, the number of solutions of $X^3 - X - 1 \equiv 0 \pmod{p}$ is $N_p(f) = a_p + 1$, where a_p is the coefficient of the modular form

$$q \prod_{k=1}^{\infty} (1 - q^k)(1 - q^{23k}) = \frac{1}{2} \sum_{(x,y) \in \mathbb{Z}^2} (q^{x^2+xy+6y^2} - q^{2x^2+xy+3y^2}) = \sum_{n=1}^{\infty} a_n q^n.$$

(See Serre's article [Ser03].) In this example we have traced out a relationship

$$(\text{algebraic variety}) \rightarrow (\text{Galois representation}) \rightarrow (\text{automorphic form}).$$

7.3 Elliptic curves and 2-dimensional Langlands

7.3.1 Galois representations and automorphic representations

Definition 16.7.10: A 2-dimensional automorphic form is a continuous function $\text{GL}_2(\mathbb{Q}) \backslash \text{GL}_2(\mathbb{A}_{\mathbb{Q}})$ satisfying certain conditions.

A large class of 2-dimensional automorphic forms can be related to modular forms. A holomorphic function $f(z) : \mathcal{H} \rightarrow \mathbb{C}$ is a **modular function** of weight k for a congruence subgroup $\Gamma \subseteq \text{GL}_2(\mathbb{Z})$ if

$$f(\gamma z) = (cz + d)^k f(z) \text{ for all } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma.$$

If $\Gamma = \Gamma_0(N) := \left\{ M \in \text{SL}_2(\mathbb{Z}) : M \equiv \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}$, we say f is of **level** N . Here \mathcal{H} denotes the upper half-plane $\{z : \Im(z) > 0\}$ and $\gamma z = \frac{az+b}{cz+d}$.

A modular function is a **modular form** if it is holomorphic at cusps of $\mathcal{H}^* = \mathcal{H} \cup \mathbb{P}^1(\mathbb{Q})$. A **cusp form** is a modular form that vanishes at the cusps.

There is a way to go from modular forms to Galois representations; this is better understood than going in the opposite direction. One of the biggest theorems in the 2-D case is Serre's conjecture, now a theorem, that tells us that we can go from Galois representations to modular forms in certain cases.

Definition 16.7.11: We say a Galois representation is **modular** if there exists a cusp form f of some level N and a finite set S such that

$$f = \sum_{n=1}^{\infty} a_n q^n, \quad \text{tr}(\rho(\text{Frob}(p))) = a_p \text{ for } p \notin S.$$

Theorem 16.7.12 (Serre's conjecture; Khare, Wintenberger). *Any irreducible odd Galois representation $\rho : G(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\overline{\mathbb{F}}_p)$ is modular.*

7.3.2 Elliptic curves and Galois representations

Given an elliptic curve, we can define a Galois representation by looking at its torsion points.

Definition 16.7.13: Let E be an elliptic curve over a number field K . It is known that the m -torsion points $E[m]$ over \overline{K} satisfy

$$E[m] \cong \mathbb{Z}/m \times \mathbb{Z}/m.$$

(See Silverman [Sil86, III.6.4].)

Define the ℓ -adic **Tate module** of E by

$$T_\ell E := \varprojlim_n E[\ell^n] \cong \mathbb{Z}_\ell^2.$$

As $G(\overline{K}/K)$ acts on $E[\ell^n]$ for each n , it acts on $T_\ell E$, so we get a map

$$G(\overline{K}/K) \rightarrow \text{Aut } T_\ell E = \text{GL}_2(\mathbb{Z}_\ell) \hookrightarrow \text{GL}_2(\mathbb{Q}_\ell),$$

called the ℓ -adic **Galois representation** of E .⁵

Thus we can define the L -series of an elliptic curve, by defining it as the L -series of the corresponding Galois representation. (Roughly speaking, this definition is independent of the choice of ℓ .) We'll flesh out this definition in Section 7. Thus we have the (tentative) correspondences

$$(\text{Elliptic curves}) \dashrightarrow (\text{Galois representations}) \dashrightarrow (\text{cusp forms}) \quad (16.10)$$

$$(\text{L-series of elliptic curve}) \dashrightarrow (\text{L-series of modular form}). \quad (16.11)$$

Again, more is known about L -series of modular forms since modular forms have nice analytic properties and transformation properties. The theory of Jacquet-Langlands establishes analytic continuation and functional equations for L -series coming from modular forms.

This relationships in (16.10) and (16.11) are involved in the proof of two big theorems.

1. We now know the dotted lines in (16.10) are true, thanks to the following.

Theorem 16.7.14 (Modularity Theorem; Taniyama-Shimura-Weil). *All elliptic curves are modular.*

The heart of this proof is in showing that the Galois representations associated to the elliptic curves come from modular forms. This theorem (or rather, its earlier version with semistable elliptic curves) is what allowed the proof of Fermat's last theorem: there is no nontrivial solution to $a^n + b^n = c^n$ for $n > 2$. A nontrivial solution would give rise to an elliptic curve associated to a modular form that does not exist.

⁵Alternatively, let $V_\ell E := T_\ell E \otimes \mathbb{Q}$ and consider $G(\overline{K}/K)$ as acting on $V_\ell E$.

2. By working with L -functions of the elliptic curves, and reinterpreting them as L -functions of certain automorphic forms as in (16.11), one can prove the following.

Theorem 16.7.15 (Sato-Tate conjecture; Barnet-Lamb, Geraghty, Harris, Taylor). *Let E be an elliptic curve without complex multiplication, and let $E(\mathbb{F}_p)$ denote the set of solutions to E over \mathbb{F}_p . The density of primes p with $|E(\mathbb{F}_p)| \in [p + 1 + a\sqrt{p}, p + 1 + b\sqrt{p}]$, for $-1 \leq a \leq b \leq 1$ is*

$$d(\{p : |E(\mathbb{F}_p)| \in [p + 1 + a\sqrt{p}, p + 1 + b\sqrt{p}]\}) = \frac{2}{\pi} \int_a^b \sqrt{1 - x^2} dx.$$

By the correspondence between elliptic curves and modular forms, another way to phrase this theorem is that the distribution of coefficients of certain modular forms is the same “semicircle” distribution.

This theorem is like the elliptic curve analogue of the Dirichlet’s theorem on the distribution of primes in congruence classes.

8 Problems

- 3.1 (from Serre, [Ser03]) Using Chebotarev’s Density Theorem, prove the following.

Theorem. *Let $f \in \mathbb{Z}[X]$ be an irreducible polynomial of degree $n \geq 2$. Let $N_p(f)$ denote the number of zeros of f in \mathbb{F}_p . Then the set $P_0(f)$ of primes with $N_p(f) = 0$ has a density $c_0(f)$. Moreover, $c_0(f) \geq \frac{1}{n}$, with strict inequality if n is not a prime power.*

You may use the following theorem from group theory.

Theorem (Jordan). *Let G is a group acting transitively on a finite set S with $n \geq 2$ elements. There exists $g \in G$ having no fixed point in S . If n is not a prime power, then there exist at least 2 such g .*

- 3.2 (All primes divide some coefficient of Δ) Let ℓ be a given prime, and K_ℓ be the maximal extension of \mathbb{Q} ramified only at ℓ . Given that there is a continuous homomorphism (a.k.a. Galois representation)

$$\tilde{\rho}_\ell : G(K_\ell/\mathbb{Q}) \rightarrow \mathrm{GL}_2(\mathbb{F}_\ell)$$

such that

$$\mathrm{tr}(\tilde{\rho}_\ell(\mathrm{Frob}_{K_\ell/\mathbb{Q}}(p))) = \tau(p)$$

for all $p \neq \ell$, and that there is an element in $\mathrm{im}(\tilde{\rho}_\ell)$ with trace 0, prove that a positive proportion of primes p have the property that

$$\ell \mid \tau(p).$$

Note. Here τ is *Ramanujan's tau function*, the coefficients of a certain modular form Δ . For more on the relationship between Galois representations and congruences for coefficients of modular forms, see Birch and Swinnerton-Dyer [SD72].

4.1 In Section 4, we showed that L/K is abelian iff the primes that split can be characterized by a modular condition. In this problem, we do more: given a Galois extension L/K , characterize the maximal abelian subextension by looking at the primes that split.

(a) Let \mathfrak{m} be a modulus for K , and suppose L/K is a Galois extension. Let $H_{\mathfrak{m}}$ be the subset of the ray class field $C_K(\mathfrak{m})$ defined as follows:

$$H_{\mathfrak{m}} = \{\mathfrak{K} : \text{There exists } \mathfrak{p} \in \mathfrak{K} \text{ such that } \mathfrak{p} \text{ splits completely in } L\}.$$

Show that $H_{\mathfrak{m}}$ is a subgroup of $C_K(\mathfrak{m})$.

(b) Suppose we are given the groups $H_{\mathfrak{m}}$ for all \mathfrak{m} . Characterize the maximal abelian subextension of L/K .

4.2 Let f be an irreducible polynomial. How could you find the Galois group of f by reducing it modulo different primes? Do this with $f = X^6 + 2X^5 + 3X^4 + 4X^3 + 5X^2 + 6X + 7$. (Exercise 8.3 from ??)

6.1 Prove an analogue of Theorem 16.6.6 for positive discriminants.

6.2 Let $n > 0$ be an integer such that $K = \mathbb{Q}(\sqrt{-n})$ is an imaginary quadratic field, and let

$$Q(x, y) = \begin{cases} x^2 + ny^2, & n \equiv 1 \pmod{4} \\ x^2 + xy + \frac{1-n}{2}y^2, & n \equiv 3 \pmod{4}. \end{cases}$$

(a) Find a condition on $G(H_K/K)$ so that for all but a finite number of primes, the primes represented by Q are given by a modulo condition. In other words, find all n such that there exists m and a set of residues S modulo m such that if $p \nmid m$, then p is represented by Q iff p is congruent to a residue in S . (Hint: combine the results of Section 4 with Section 6.)

(b) Find some values of n for which $|\text{Cl}(K)| \neq 1$ and such that the primes represented by Q are given by a modulo condition.

(c) Suppose $G(H_K/K)$ satisfies the property you found in part 1. Characterize all n , not necessarily prime, such that Q represents n . (For simplicity, you can just consider $n \perp m$.) Compare to the statement in Example 4.3.4.)

1. (**Genus theory**) It is useful to group the equivalence classes of quadratic forms with given discriminant into *genera* (plural of genus).

Definition 16.8.1: Define a similarity relation between primitive quadratic forms of discriminant d as follows. We say $Q_1 \sim Q_2$ if Q_1 and Q_2 represent the same values in $(\mathbb{Z}/d\mathbb{Z})^\times$. The similarity classes are called **genera**.

In this problem you will find an easy way to characterize the genera of discriminant d .

- (a) Let H be the subgroup of $C(d)$ such that $C(d)/H \cong C(d)[2]$ where $G[n]$ denotes the n -torsion subgroup of G .

Prove that $Q_1, Q_2 \in C(d)$ are in the same genera iff Q_1, Q_2 are the same in $C(d)/H$.

In particular, conclude that the number of genera is a power of 2.

- (b) Let M be the ring class field of K and let L denote the subextension of M/K such that $G(L/K) \cong C(d)[2]$. (That is, under the Galois correspondence, $L \subseteq M$ corresponds to $H \subseteq C(d) = G(M/K)$.) Prove that L/\mathbb{Q} is the maximal abelian subextension of M/\mathbb{Q} .

The fact that L/K is abelian, while M/K may not be, makes it much easier to prove results pertaining to a genus of quadratic forms rather than an equivalence class of quadratic forms.

2. (?) Let $f, g \in \mathbb{Q}[X]$ be two irreducible cubic polynomials. How can you determine algorithmically whether f, g have roots α, β such that $\mathbb{Q}(\alpha) = \mathbb{Q}(\beta)$? (from <http://math.stackexchange.com/questions/34522/cubic-polynomials-that-generate-rq=1>)

Chapter 17

Complex multiplication

In this chapter, we combine class field theory with the theory of elliptic curves, first to characterize the maximal abelian extension of K , then to illustrate the relationships in Section 16.7 for CM elliptic curves. We will assume basic facts about elliptic curves (for an introduction see Silverman [Sil86, Chapter III]).

We know that every elliptic curve over \mathbb{C} has endomorphism ring either equal to \mathbb{Z} or a quadratic order. In the second case, the elliptic curve is said to have **complex multiplication**. This gives the elliptic curve a lot more structure. On one hand, it is useful algebraically—as we will see, torsion points of a CM elliptic curve give abelian extensions of imaginary quadratic fields. In general, because of the added structure, much more is known about CM elliptic curves than other elliptic curves, and they can act as a kind of “testing ground” or “first case” of general conjectures.

On the other hand, CM elliptic curves have practical uses—for instance, if we take an CM elliptic curve corresponding to a specific endomorphism ring, we can easily compute its order. Hence we can generate an elliptic curve with near-prime order, useful in cryptography. This is much more efficient than generating random elliptic curves and using Schoof’s algorithm to find their orders.

There are several big theorems about complex multiplication. In Section 2, we specialize our knowledge about the relationship between elliptic curves over \mathbb{C} and complex tori to CM elliptic curves and build a toolbox of basic facts. However, since we are interested in number theory, we want to take curves defined over \mathbb{C} and define them over $\overline{\mathbb{Q}}$ instead—which we do in Section 3. Once we have these basics, we can then prove the big theorems.

We suppose E has CM by a quadratic order $\mathcal{O} \subset K$ (i.e. $\text{End}(E) \cong \mathcal{O}$), where K is a quadratic extension of \mathbb{Q} . Then the following hold.

1. The j -invariant $j(E)$ generates the *ring class field* of \mathcal{O} over K . In particular, if $\mathcal{O} = \mathcal{O}_K$, then $j(E)$ generates the *Hilbert class field* of K , the maximal unramified abelian extension (Theorem 17.4.4):

$$K(j(E)) = H_K.$$

2. If E is defined over H_K , and we adjoin certain functions of torsion points of E , then

we get the *maximal abelian extension* of K (Theorem 17.5.4):

$$K(j(E), h(E_{\text{tors}})) = K^{\text{ab}}.$$

Compare this with the Kronecker-Weber Theorem, which says the maximal abelian extension of \mathbb{Q} is generated by roots of unity (torsion points of $\overline{\mathbb{Q}}^\times$).

3. $j(E)$ is moreover an *algebraic integer* (We omit this; see Silverman AT, [Sil94, II.6].)
4. The action of the idele class group sending K/\mathfrak{a} to $K/\mathfrak{x}^{-1}\mathfrak{a}$ corresponds to the Galois action on the corresponding elliptic curves, where the Galois action is given by the Frobenius element of σ . This is the Main Theorem of Complex Multiplication 17.6.2, and plays an important part in taking moduli spaces initially defined only over \mathbb{C} and defining them over algebraic number fields.
5. The L -series of a CM elliptic curve is particularly easy to understand, because it is a product of 2 Hecke L -series (Theorem 17.7.5).

Two “big ideas” we’ll consistently see are the following.

1. We expect abelian extensions because for CM elliptic curves (with endomorphism ring \mathcal{O}_K , say), the image of the map $G(L/H_K) \hookrightarrow \text{Aut}(E[m])$ commutes with \mathcal{O}_K , not just \mathbb{Z} and hence must be abelian, with appropriate L .
2. We can use torsion points $E[m]$ to “keep book” on the action of Frobenius, in the same way that we used the roots of unity μ_m to keep book on the action of Frobenius on $G(\mathbb{Q}(\mu_m)/\mathbb{Q})$.

1 Elliptic curves over \mathbb{C}

The following theorem helps us understand elliptic curves over \mathbb{C} .

Theorem 17.1.1. *Let $g_2(\Lambda) = 60G_4(\Lambda)$ and $g_3(\Lambda) = 140G_6(\Lambda)$, where G_n is the Eisenstein series. Let Λ be a lattice in \mathbb{C} and \wp be the associated Weierstrass \wp -function.*

There is a complex analytic isomorphism between the complex torus \mathbb{C}/Λ and the elliptic curve over \mathbb{C} ,

$$y^2 = 4x^3 - g_2(\Lambda)x - g_3(\Lambda)$$

given by

$$\Phi(z) = (\wp(z), \wp'(z)).$$

The map Φ gives an equivalence of categories between the following.

1. *Objects: Complex tori \mathbb{C}/Λ , where Λ is a lattice in \mathbb{C} .
Maps: Multiplication-by- α $\mathbb{C}/\Lambda_1 \rightarrow \mathbb{C}/\Lambda_2$ where $\alpha\Lambda_1 \subseteq \Lambda_2$.*

2. *Objects: Elliptic curves over \mathbb{C} .*
Maps: Isogenies.

Proof. Silverman [Sil86, VI.5.1.1, 5.3] □

The endomorphism ring of a lattice $\Lambda \subset \mathbb{C}$ is either \mathbb{Z} or an imaginary quadratic order, so the same is true of an elliptic curve E over \mathbb{C} . If the endomorphism ring is a quadratic order \mathcal{O} , we say E has **complex multiplication** by \mathcal{O} .

2 Complex multiplication over \mathbb{C}

2.1 Embedding the endomorphism ring

We know the endomorphism ring $\text{End}(E)$ of a CM elliptic curve corresponds to a quadratic order \mathcal{O} but since any quadratic order has conjugation as an isomorphism, we need to specify a way to embed $\text{End}(E)$ into \mathbb{C} .

Example 17.2.1: Consider the curve $E : y^2 = x^3 + x$. We note that the endomorphisms

$$\begin{aligned}\phi_1(x, y) &= (-x, iy) \\ \phi_2(x, y) &= (-x, -iy)\end{aligned}$$

both square to -1 . Which one should we call $[i]$, multiplication by i ?

Fortunately, we have a way of embedding $\text{End}(\Lambda)$ into \mathbb{C} , where Λ is the lattice corresponding to E , because Λ itself is in \mathbb{C} . This to give a canonical way of embedding $\text{End}(E)$ into \mathbb{C} .

Proposition 17.2.2: Let E/\mathbb{C} be a CM elliptic curve with complex multiplication by \mathcal{O} . There is a unique isomorphism $[\cdot] : \mathcal{O} \xrightarrow{\cong} \text{End}(E)$ satisfying either of the following equivalent conditions.

1. $[\alpha]$ is the unique morphism making the following diagram commute, where the top map is multiplication by α .

$$\begin{array}{ccc} \mathbb{C}/\Lambda & \xrightarrow{m_\alpha} & \mathbb{C}/\Lambda \\ \downarrow \Phi & & \downarrow \Phi \\ E_\Lambda & \xrightarrow{[\alpha]} & E_\Lambda \end{array}$$

2. For any invariant differential $\omega \in \Omega_E$, $[\alpha]^*\omega = \alpha\omega$.

Moreover, we have the following.

3. Define $[\cdot]_1$ and $[\cdot]_2$ for elliptic curves E_1 and E_2 . For any morphism $\phi : E_1 \rightarrow E_2$,

$$\phi \circ [\alpha]_1 = [\alpha]_2 \circ \phi.$$

In other words, multiplication by α commutes with all morphisms.

4. For any $\sigma \in \text{Aut}(\mathbb{C})$,

$$[\alpha]_E^\sigma = [\sigma(\alpha)]_{\sigma(E)},$$

i.e. it commutes with Galois action.

The pair $(E, [\cdot])$ is called a **normalized** elliptic curve. After we prove this proposition, we will assume all CM elliptic curves are normalized.

Proof. The uniqueness and existence of $[\alpha]$ satisfying item 1 follows directly from the equivalence of categories (Theorem 17.1.1).

Define $[\alpha]$ as in item 1. For any invariant differential ω on E_Λ , since Φ is an analytic isomorphism, we can consider its pullback to \mathbb{C}/Λ ; it will be cdz for some c (The space of invariant differentials on \mathbb{C}/Λ is 1-dimensional.) Clearly, $m_\alpha^*(cdz) = cd(\alpha z) = \alpha cdz$. Transferring this to the bottom row of the commutative diagram gives $[\alpha]^*\omega = \alpha\omega$. For uniqueness, note the map

$$\begin{aligned} \text{Hom}(E_1, E_2) &\hookrightarrow \text{Hom}(\Omega_{E_2}, \Omega_{E_1}) \\ \phi &\rightarrow \phi^* \end{aligned} \tag{17.1}$$

is injective when all isogenies $E_1 \rightarrow E_2$ are separable (in particular, in characteristic 0), i.e. the action of an isogeny of elliptic curves on an invariant differential completely determines the morphism. Taking $E_1 = E_2$ and considering the preimage of multiplication-by- α gives uniqueness in item 2.

A simple diagram chase shows that $(\phi \circ [\alpha]_1)^*$ and $([\alpha]_2 \circ \phi)^*$ act the same way on $\omega \in \Omega_{E_2}$. Then (17.1) gives item 3.

The proof of item 4 is similar. □

Example 17.2.3: The definition using differentials is useful for calculations. Revisiting the above Example 17.2.1, we see that we should let

$$[i](x, y) = (-x, iy).$$

Indeed, defining $[i]$ in this way, we check that

$$[i]^* \frac{dx}{y} = \frac{d(-x)}{iy} = i \frac{dx}{y}.$$

2.2 The class group parameterizes elliptic curves

Let K be an imaginary quadratic field and \mathcal{O} an order inside K .

Definition 17.2.4: Let L be a field. Define

$$\begin{aligned} \text{Ell}_L(\mathcal{O}) &= \{\text{elliptic curves } E/L \text{ with } \text{End}(E) \cong \mathcal{O}\} \\ \mathcal{E}\text{ll}_L(\mathcal{O}) &= \frac{\{\text{elliptic curves } E/L \text{ with } \text{End}(E) \cong \mathcal{O}\}}{\text{isomorphism over } L}, \end{aligned}$$

i.e. $\mathcal{E}ll_L(\mathcal{O})$ is the set of elliptic curves over L whose endomorphism ring is \mathcal{O} . If we omit L , we assume $L = \mathbb{C}$.

If $E \in \text{Ell}(\mathcal{O})$, then its corresponding lattice Λ must be homothetic to a fractional ideal of \mathcal{O} : indeed, we can scale the lattice so that $1 \in \Lambda$; then $\mathcal{O} \subseteq \Lambda$ so $\Lambda \subseteq K$; since it is a lattice it must be a fractional \mathcal{O} -ideal. Now note an \mathcal{O} -ideal \mathfrak{a} has endomorphism ring \mathcal{O} iff \mathfrak{a} is a *proper* ideal (see Definition 4.4.4.5).¹ Hence we get a correspondence between isomorphism classes of elliptic curves $[E] \in \mathcal{E}ll(\mathcal{O})$ and proper \mathcal{O} -ideals up to homothety. However, two fractional ideals \mathfrak{a} and \mathfrak{b} are homothetic iff $\lambda\mathfrak{a} = \mathfrak{b}$ for some λ , i.e. iff they are equivalent in the class group. Thus the class group of \mathcal{O} parameterizes all isomorphism classes of elliptic curves with endomorphism ring \mathcal{O} . This is summarized in the following.

$$\mathcal{E}ll(\mathcal{O}) = \frac{\{\text{elliptic curves } E/\mathbb{C} \text{ with } \text{End}(E) \cong \mathcal{O}\}}{\text{isomorphism over } \mathbb{C}} = \frac{\{\text{proper fractional } \mathcal{O}\text{-ideal}\}}{\text{principal } \mathcal{O}\text{-ideals}} = \text{Cl}(\mathcal{O}).$$

We state this as a theorem.

Theorem 17.2.5. *We have a bijection*

$$\mathcal{E}ll(\mathcal{O}) \cong \text{Cl}(\mathcal{O})$$

where $[E] \in \mathcal{E}ll(\mathcal{O})$ is sent to a $[\mathfrak{a}]$, where \mathfrak{a} is a fractional ideal homothetic to the lattice corresponding to E .

We get much more than this, however. $\mathcal{E}ll(\mathcal{O})$ is a priori just a set; however, $\text{Cl}(\mathcal{O})$ is a group. We can define the action of $I(\mathcal{O})$ on $\text{Ell}(\mathcal{O})$ since $I(\mathcal{O})$ acts on lattices. This action will descend to an action of $\text{Cl}(\mathcal{O})$ on $\mathcal{E}ll(\mathcal{O})$, since isomorphic elliptic curves correspond to equivalent ideals.

Theorem 17.2.6. *There is a group action of $\text{Id}(\mathcal{O})$ on $\text{Ell}(\mathcal{O})$ given by*

$$\mathfrak{a}E_\Lambda = E_{\mathfrak{a}^{-1}\Lambda}$$

where E_Λ denotes the elliptic curve corresponding to the lattice Λ .

This descends to a simply transitive group action of $\text{Cl}(\mathcal{O})$ on $\mathcal{E}ll(\mathcal{O})$.

Proof. Just check that if Λ has endomorphism ring \mathcal{O} , then so does the lattice $\mathfrak{a}^{-1}\Lambda$. (Note that $\mathfrak{b}L$ is defined by $\{s\alpha : s \in \mathfrak{b}, \alpha \in L\}$.)

For the second part, note that $E_\Lambda \cong \mathfrak{a}E = E_{\mathfrak{a}^{-1}\Lambda}$ iff Λ and $\mathfrak{a}^{-1}\Lambda$ are homothetic, i.e. \mathfrak{a} is principal. □

Remark 17.2.7: Another way of saying that $\text{Cl}(\mathcal{O})$ acts simply transitively on $\mathcal{E}ll(\mathcal{O})$ is that $\mathcal{E}ll(\mathcal{O})$ is a **torsor** or **principal homogeneous space** for $\text{Cl}(\mathcal{O})$.

This action will be fundamental to our understanding of CM elliptic curves. Later on we will relate this to the Galois action. The interplay between these two actions is the source for much of the richness of CM theory.

¹When $R = \mathcal{O}_K$, all ideals are proper, so this distinction is not important. The reader unfamiliar with non-maximal orders can take $R = \mathcal{O}_K$ throughout.

2.3 Ideals define maps

For any $n \in \mathbb{Z}$ and any elliptic curve E , n defines the multiplication by n map $[n] : E \rightarrow E$. When E has CM, we saw in Theorem 17.2.2 that $\alpha \in \mathcal{O}$ defines (canonically) the multiplication by α map $[\alpha] : E \rightarrow E$. We now extend this to *ideals*: if \mathfrak{a} is a proper \mathcal{O} -ideal, \mathfrak{a} determines a “multiplication by \mathfrak{a} ” map. The only difference is that $[\mathfrak{a}]$ is now a map $E \rightarrow \mathfrak{a}E$.

Definition 17.2.8: Let $E \in \text{Ell}(\mathcal{O})$ correspond to the lattice Λ . Let \mathfrak{a} be a proper integral ideal of \mathcal{O} . We have $\mathfrak{a}\Lambda \subseteq \Lambda$, so \mathfrak{a} determines a map $\mathbb{C}/\Lambda \rightarrow \mathbb{C}/\mathfrak{a}^{-1}\Lambda$, sending $z \mapsto z$. Define the multiplication by \mathfrak{a} -map as the corresponding map on elliptic curves

$$[\mathfrak{a}] : E \rightarrow E_{\mathfrak{a}^{-1}\Lambda} = \mathfrak{a}E.$$

Proposition 17.2.9: Let $E \in \text{Ell}(\mathcal{O}_K)$. We have the following.

1. The kernel of $[\mathfrak{a}]$ (the “ \mathfrak{a} -torsion points”) is

$$E[\mathfrak{a}] := \{P \in E : [\alpha]P = 0 \text{ for all } \alpha \in \mathfrak{a}\} \cong \mathcal{O}_K/\mathfrak{a}.$$

2. The degree of $[\mathfrak{a}]$ is

$$\deg([\mathfrak{a}]) = |E[\mathfrak{a}]| = \mathfrak{N}(\mathfrak{a}),$$

and in particular, $\deg([\alpha]) = |E[\alpha]| = \text{Nm}_{K/\mathbb{Q}}(\alpha)$.

Proof. Silverman AT [Sil94, pg. 102-3]. □

3 Defining CM elliptic curves over $\overline{\mathbb{Q}}$

We show that we do not lose anything if we just consider elliptic curves over $\overline{\mathbb{Q}}$ instead of over \mathbb{C} . To do this, we look at the j -invariants.

Proposition 17.3.1: Suppose E is an elliptic curve with CM by a quadratic order \mathcal{O} . Then $j(E) \in \overline{\mathbb{Q}}$, i.e. $j(E)$ is algebraic.

Proof. Let σ be any automorphism of \mathbb{C} over \mathbb{Q} . We look at how σ acts on $j(E)$.

Note that E^σ is defined by taking any equation for E and operating on all the coefficients of E by σ , so $\sigma(j(E)) = j(E^\sigma)$.

First note that $\text{End}(E) \cong \text{End}(E^\sigma)$ by the map $\phi \mapsto \phi^\sigma$. Hence $\text{End}(\sigma(E)) = \mathcal{O}$ as well. But $\text{Cl}(\mathcal{O})$ is finite, and as $|\text{Cl}(\mathcal{O})| = |\mathcal{E}\text{ll}(\mathcal{O})|$ (Theorem 17.2.5) we see that the E^σ lie in finitely many isomorphism classes. Because isomorphic elliptic curves have the same j -invariant, there are a finite number of possibilities for $j(E^\sigma)$.

As $\{\sigma(j(E)) : \sigma \in \text{Aut}(\mathbb{C})\}$ is finite, $j(E)$ must be algebraic. □

This allows us to prove the following.

Theorem 17.3.2. *We have*

$$\mathcal{E}ll_{\mathbb{C}}(\mathcal{O}) \cong \mathcal{E}ll_{\overline{\mathbb{Q}}}(\mathcal{O}).$$

Proof. We use the following properties of the j -invariant. ([Sil86, III.1.4])

1. For every $j \in K$, there exists an elliptic curve E/K with $j(E) = j$.
2. Let K be an algebraically closed field and E_1, E_2 be elliptic curves defined over K . Then $E_1 \cong E_2$ over K iff $j(E_1) = j(E_2)$. (The backwards direction does not necessarily hold if K is not algebraically closed.)

We show that the map

$$\mathcal{E}ll_{\overline{\mathbb{Q}}}(\mathcal{O}) \rightarrow \mathcal{E}ll_{\mathbb{C}}(\mathcal{O}) \tag{17.2}$$

is an isomorphism (of sets, in fact, of $\text{Cl}(\mathcal{O})$ -modules). The map is well-defined, because any automorphism over $\overline{\mathbb{Q}}$ is an automorphism over \mathbb{C} .

By Lemma 17.3.1, if $[E] \in \mathcal{E}ll_{\mathbb{C}}(\mathcal{O})$ then $j(E) \in \overline{\mathbb{Q}}$. By item 1, there exists an elliptic curve E' defined over $\overline{\mathbb{Q}}$ with $j(E') = j(E)$. Then E' is isomorphic to E over \mathbb{C} . Thus the map (17.2) above is surjective. It is injective because if E, E' are defined over $\overline{\mathbb{Q}}$ and isomorphic over \mathbb{C} , then item 2 says $j(E) = j(E')$; and the other direction of item 2 says that $E \cong E'$ over $\overline{\mathbb{Q}}$. \square

It is also important to know what fields we can define elliptic curves and isogenies over.

Proposition 17.3.3: Suppose E is an elliptic curve with CM by $\mathcal{O} \subset K$, where K is an imaginary quadratic field.

1. If E is defined over L then endomorphisms of E can be defined over LK .
2. If E_1, E_2 are defined over L then there exists a finite extension M/L , so that every isogeny $E_1 \rightarrow E_2$ is defined over M .

Proof. For item 1, note that all endomorphisms are in the form $[\alpha]$ and use Proposition 17.2.2(4).

For item 2, first we claim that any isogeny ϕ is defined over a finite extension of L . For any $\sigma \in \text{Aut}(\mathbb{C})$ fixing L , ϕ^σ is a map $E_1 \rightarrow E_2$ having the same degree as ϕ . Any isogeny is determined by its kernel, up to automorphism of E_1 and E_2 . As E_1 has a finite number of subgroups of given index and $\deg(\phi) = \#\ker(\phi)$, there are finitely many isogenies of a given degree. Hence $\{\phi^\sigma : \sigma \in G(\mathbb{C}/L)\}$ is finite, showing ϕ is defined over a finite extension of L .

Now $\text{Hom}(E_1, E_2)$ is a finitely generated group, so we can take the field of definition for a finite set of generators. \square

4 Hilbert class field

4.1 Motivation: Class field theory for $\mathbb{Q}(\zeta_n)$ and Kronecker-Weber

4.1.1 The case of \mathbb{Q}

First we give some motivation for the next two sections by making an analogy with class field theory for $\mathbb{Q}(\zeta_n)$. We can think of μ_n , the n th roots of unity, as the analogue of $E[n]$: μ_n are the n -torsion points of the group variety $\overline{\mathbb{Q}}^\times$ under multiplication, and $E[n]$ are the n -torsion points of an elliptic curve. To emphasize this analogy, we write $K^\times[n]$ to denote the n th roots of unity in \overline{K} .

Recall how we established class field theory for $\mathbb{Q}(\zeta_n)$: given a prime p , we want to find $(p, \mathbb{Q}(\zeta_n)/\mathbb{Q})$. To do this we looked at the action of $(p, \mathbb{Q}(\zeta_n)/\mathbb{Q})$ on $\mathbb{Q}^\times[n] = \mu_n$, by taking everything modulo p . We know by definition of $(p, \mathbb{Q}(\zeta_n)/\mathbb{Q})$ how it must act on the residue field extension $\mathbb{F}_p/\mathbb{F}_p$ and hence on $\mathbb{F}_p^\times[n]$. Suppose $p \nmid n$. Because the maps

$$\begin{aligned} \mathbb{Q}^\times[n] &\hookrightarrow \mathbb{F}_p^\times[n] \\ \text{End}(\mathbb{Q}^\times[n]) &\hookrightarrow \text{End}(\mathbb{F}_p^\times[n]) \end{aligned} \tag{17.3}$$

are injective (the first is because $p \nmid n$ and the second is a direct consequence of the first), once we know how $(p, \mathbb{Q}(\zeta_n)/\mathbb{Q})$ acts on $\mathbb{F}_p^\times[n]$, we know it acts on $\mathbb{Q}^\times[n]$, so we know exactly what automorphism it is:

$$(p, \mathbb{Q}(\zeta_n)/\mathbb{Q})(\zeta_n) = \zeta_n^p.$$

In particular, since ζ_n is a n -torsion point (i.e. $\zeta_n^n = 1$) this only depends on $p \pmod n$. Hence we get the Artin map $\psi_{\mathbb{Q}(\zeta_n)/\mathbb{Q}}$ factoring through the modulus ∞n :²

$$\psi_{\mathbb{Q}(\zeta_n)/\mathbb{Q}} : I_{\mathbb{Q}}/I_{\mathbb{Q}}(1, n\infty) \xrightarrow{\cong} G(\mathbb{Q}(\zeta_n)/\mathbb{Q}).$$

Finally, since every modulus divides ∞n for some n , we get the Kronecker-Weber Theorem

$$\mathbb{Q}^{\text{ab}} = \mathbb{Q}(\zeta_\infty) = \mathbb{Q}(\mathbb{Q}^\times[\infty]).$$

In summary, we found the ray class groups and thus the maximal abelian extension by looking at how $(p, \mathbb{Q}(\zeta_n)/\mathbb{Q})$ acted on $\mathbb{Q}^\times[n]$:

$$\begin{array}{ccc} \mathbb{Q}^\times[n] & \xrightarrow[\text{reduction}]{\tilde{\bullet}} & \mathbb{F}_p^\times[n] \\ \circlearrowleft & & \circlearrowleft \\ I_{\mathbb{Q}}/P_{\mathbb{Q}}(1, n\infty) & \xrightarrow{\psi_{\mathbb{Q}(\zeta_n)/\mathbb{Q}}} & G(\mathbb{Q}(\zeta_n)/\mathbb{Q}) \xrightarrow{\tilde{\bullet}} G(\mathbb{F}_p(\zeta_n)/\mathbb{F}_p). \end{array} \tag{17.4}$$

²The ∞ is a technical detail coming from the fact that \mathbb{Q} is totally real.

4.1.2 The case of K

One big difference when we're working over an imaginary quadratic field K is that while we had $\text{Cl}_{\mathbb{Q}} = 1$, we have Cl_K is nontrivial in general. This corresponds to the fact that there is only 1 nonisomorphic "version" of $\mathbb{G}_m(\mathbb{Q}) = \mathbb{Q}^\times$, but multiple elliptic curves with endomorphism ring by the same order \mathcal{O} . Hence $G(K^{\text{ab}}/K)$ no longer operates on the same elliptic curve. Instead we have to analyze it in two steps.

1. Consider the action of $G(H_K/K)$ on $\mathcal{E}ll_{\overline{\mathbb{Q}}}(\mathcal{O})$, i.e. equivalence classes of elliptic curves with CM by \mathcal{O} .
2. Consider the action of $G(K^{\text{ab}}/H_K)$ on the torsion points E_{tors} of a single elliptic curve.

In both cases, we will understand the action by looking at how the Frobenius elements of the Galois groups act.

4.1.3 The case of K : Part 1

We have two natural actions on the set of elliptic curve $\mathcal{E}ll_{\overline{\mathbb{Q}}}(\mathcal{O}_K)$, namely the action of $G(\overline{K}/K)$ and $\text{Cl}(\mathcal{O}_K)$. Our first task is to relate these, i.e. find a dotted map that preserves the action on $\mathcal{E}ll_{\overline{\mathbb{Q}}}(\mathcal{O}_K)$:

$$\begin{array}{ccc}
 & \mathcal{E}ll_{\overline{\mathbb{Q}}}(\mathcal{O}_K) & \\
 & \circlearrowleft & \circlearrowright \\
 G(\overline{K}/K) & \xrightarrow{\hspace{10em}} & \text{Cl}(\mathcal{O}_K).
 \end{array} \tag{17.5}$$

We'll see that this map factors through $G(L/K)$ where $L = K(j(E))$. We have a map $\psi_{L/K} : I_K/P_K(1, \mathfrak{f}) \rightarrow G(L/K)$; we show that $\mathfrak{f} = 1$ and the composition of the two maps is an isomorphism, and that in fact we have

$$\begin{array}{ccc}
 & \mathcal{E}ll_{\overline{\mathbb{Q}}}(\mathcal{O}_K) & \\
 & \circlearrowleft & \circlearrowright \\
 I_K/P_K & \xrightarrow{\Psi_{L/K}} G(L/K) & \xrightarrow{\hspace{10em}} \text{Cl}(\mathcal{O}_K). \\
 & \xrightarrow{\alpha \rightarrow [\mathfrak{a}]} &
 \end{array} \tag{17.6}$$

We establish (17.6) by looking at the reduction of the elliptic curves modulo some \mathfrak{P} .

Since $G(H_K/K) \cong \text{Cl}(\mathcal{O}_K)$ this will show that $L = H_K$, the Hilbert class field of K .

4.1.4 The case of K : Part 2

We can now do the same thing we did with \mathbb{Q} , use the torsion points of elliptic curves to find the ray class fields and the maximal abelian extensions. We can't work directly over K because Cl_K is nonzero, but if we imitate the argument (with some modifications) over

\mathbb{Q} for H_K we will get the ray class fields of K . We let $L_n = K(j(E), h(E[n]))$ where h is a Weber function (to be defined).

Let l_n, l be the residue fields of L_n and H_K modulo some prime. We show L_n is the ray class field for (n) by constructing the diagram

$$\begin{array}{ccc}
 E[n] & \xrightarrow[\text{reduction}]{\bullet} & \widetilde{E}[n] \\
 \circ & & \circ \\
 \text{Nm}_{H_K/K}(I_{H_K}^n)/P_K(1, n) & \xrightarrow{\psi_{L_n/K}} & G(L_n/H_K) \xrightarrow{\bullet} G(l_n/l).
 \end{array} \tag{17.7}$$

We now carry out these two parts.

4.2 The Galois group and class group act compatibly

We establish the map in (17.5).

Theorem 17.4.1. *There exists a map $F : G(\overline{K}/K) \rightarrow \text{Cl}(\mathcal{O}_K)$ such that for any elliptic curve E ,*

$$[E^\sigma] = F(\sigma)E.$$

This map factors through $G(K^{ab}/K)$.

As a reminder, the action of $\text{Cl}(\mathcal{O}_K)$ on $\mathcal{E}\text{ll}_{\mathbb{Q}}(\mathcal{O}_K)$ is such that if $E = E_\Lambda$, then $F(\sigma)E = E_{F(\sigma)^{-1}\Lambda}$. Theorem 17.4.1 expresses a deep relationship because the left-hand side expresses an algebraic action, while the right-hand side expresses an analytic action, as it is defined on lattices and the map between E and \mathbb{C}/Λ is inherently analytic.

Proving this theorem essentially boils down to showing the Galois action commutes with the action on $\text{Cl}(\mathcal{O}_K)$.

Proposition 17.4.2: For all E ,

$$\sigma([\mathfrak{a}][E]) = [\sigma(\mathfrak{a})][\sigma(E)].$$

Proof. Suppose E corresponds to Λ , i.e. $E \cong \mathbb{C}/\Lambda\mathbb{C}$. Then we have the exact sequence

$$0 \rightarrow \Lambda \rightarrow \mathbb{C} \rightarrow E \rightarrow 0.$$

Then $\mathfrak{a}E$ corresponds to $\mathfrak{a}^{-1}\Lambda$. Take a resolution for \mathfrak{a} :

$$R^m \xrightarrow{A} R^n \rightarrow \mathfrak{a} \rightarrow 0.$$

Take a ‘‘Hom product’’ and use the Snake Lemma. See [Sil94, II.2.5]. □

Proof of Theorem 17.4.1. See [Sil94, II.2.4]. □

4.3 Hilbert class field

Before we proceed with finding the Hilbert class field, we need to show injectivity of the reduction map like in (17.3).

Theorem 17.4.3. *Suppose E_1 and E_2 are elliptic curves defined over L with good reduction at \mathfrak{P} . Then the reduction map*

$$\mathrm{Hom}(E_1, E_2) \rightarrow \mathrm{Hom}(\widetilde{E}_1, \widetilde{E}_2)$$

is injective and preserves degrees.

Proof. See Silverman AT [Sil94, pg. 124] (Also see Silverman’s errata). □

The main theorem of this section is the following.

Theorem 17.4.4 ($j(E)$ generates the Hilbert class field). *Let E be an elliptic curve with CM by \mathcal{O}_K . Then*

1. $K(j(E)) = H_K$, the Hilbert class field of K .
2. $G(\overline{K}/K)$ acts transitively on the isomorphism classes of curves in $\mathcal{E}\mathrm{ll}(\mathcal{O}_K)$.
3. For any ideal $\mathfrak{a} \in I_K$,

$$[E^{\psi_{H_K/K}(\mathfrak{a})}] = [\mathfrak{a}][E].$$

In particular, the action of Frobenius on the j -invariant is given by operating by $[\mathfrak{p}]$ on the elliptic curve:

$$[E^{(\mathfrak{p}, H_K/K)}] = [\mathfrak{p}][E].$$

Proof. Step 1: First we show the following: There exists a finite set of primes S of \mathbb{Z} such that for any $p \notin S$ that splits completely in K , $p = \mathfrak{p}\overline{\mathfrak{p}}$, we have

$$F((\mathfrak{p}, L/K)) = [\mathfrak{p}] \in \mathrm{Cl}(\mathcal{O}_K).$$

This will show the dotted map in (17.6) is the identity for a large number of primes \mathfrak{p} .

We have the map $[\mathfrak{p}] : E \rightarrow \mathfrak{p}E$. We show that this is “like” the p th power Frobenius map. To do this, we show that it is inseparable of degree p (this is why we needed p to be split)³, and then look at the j -invariants of the reduced maps modulo \mathfrak{p} .

As $\mathcal{E}\mathrm{ll}_{\overline{\mathbb{Q}}}(\mathcal{O}_K) = \mathcal{E}\mathrm{ll}_{\mathbb{C}}(\mathcal{O}_K)$ is finite, we can find a finite extension L/K and representatives E_1, \dots, E_h of classes in $\mathcal{E}\mathrm{ll}_{\mathbb{C}}(\mathcal{O}_K)$, that are defined over L . Let S be a set of primes containing the primes that satisfy one of the following conditions.

1. p ramifies in L . (Primes that ramify always cause trouble.)
2. E or some E_i has bad reduction at some prime of L lying over p .

³If \mathfrak{p} is not split, one can still show the map is inseparable of degree p^2 , with some more work.

3. $v_p(\text{Nm}_{L/\mathbb{Q}}(j(E_i) - j(E_k))) \neq 0$ for some $i \neq k$. (This allows us to know what equivalence class an elliptic curve lies in, just by looking at its reduction modulo p .)

Let Λ be the lattice such that $E(\mathbb{C}) \cong \mathbb{C}/\Lambda$, and let \mathfrak{a} be an integral ideal relatively prime to \mathfrak{p} such that $\mathfrak{a}\mathfrak{p} = (\alpha)$ is principal (This exists by Corollary 2.2.2.5). By the equivalence of categories 17.1.1, the following maps on complex tori correspond to isogenies of elliptic curves:

$$\begin{array}{ccccccc} \mathbb{C}/\Lambda & \xrightarrow{i} & \mathbb{C}/\mathfrak{p}^{-1}\Lambda & \xrightarrow{i} & \mathbb{C}/\mathfrak{p}^{-1}\mathfrak{a}^{-1}\Lambda & \xrightarrow[\cong]{[\alpha]} & \mathbb{C}/\Lambda \\ \cong \downarrow \Phi & & \cong \downarrow \Phi & & \cong \downarrow \Phi & & \cong \downarrow \Phi \\ E & \xrightarrow{\phi_1} & \mathfrak{p}E & \xrightarrow{\phi_2} & \mathfrak{a}\mathfrak{p}E & \xrightarrow[\cong]{\phi_3} & E \end{array}$$

Let the composition of the top maps be f and the composition of the bottom maps be g .

Let ω be an invariant differential on E . Then $\omega' = \Phi^*\omega$ is an invariant differential on \mathbb{C}/Λ . It is in the form cdz . The composition of the top maps is just multiplication by α , so $f^*\omega' = \alpha\omega'$. By commutativity, we get $g^*\omega = \alpha\omega$ as well.

Let $p \notin S$ and $\mathfrak{P} \mid \mathfrak{p} \mid p$ in L, K, \mathbb{Q} , respectively. Since E has good reduction at \mathfrak{P} , we can reduce the elliptic curves and maps modulo \mathfrak{P} to get

$$\tilde{g}^*\tilde{\omega} = \tilde{\alpha}\tilde{\omega} = 0$$

since $\mathfrak{P} \mid \alpha$. By a criterion for separability (g is separable iff g^* does not act as 0 on Ω_E), \tilde{g} is inseparable. Now

$$\begin{aligned} \deg(\phi_1) &= \mathfrak{N}\mathfrak{p} = p, \\ \deg(\phi_2) &= \mathfrak{N}\mathfrak{a} \perp p, \\ \deg(\phi_3) &= 1. \end{aligned}$$

An inseparable map must have degree divisible by p , and the composition of separable maps is separable, so $\tilde{\phi}_1$ must be inseparable.

Any inseparable map factors through the Frobenius map:

$$\begin{array}{ccc} \tilde{E} & \xrightarrow{\phi_p} & \tilde{E}^{(p)} \\ & \searrow \tilde{\phi}_1 & \cong \downarrow \varepsilon \\ & & \mathfrak{p}\tilde{E}. \end{array} \quad (17.8)$$

We have $p \deg(\varepsilon) = \deg(\phi_p) \deg(\varepsilon) = \deg(\tilde{\phi}_1) = p$ so $\deg(\varepsilon) = 1$. This shows ε is an isomorphism.

Thus we have

$$\mathfrak{p}\tilde{E} \cong \tilde{E}^{(p)}.$$

Now by definition of the Frobenius element (it is the p th power map modulo \mathfrak{P}), we have $j(\tilde{E}^{(p)}) = j(\tilde{E})^p = j(E)^{(\mathfrak{p}, L/K)}$ modulo \mathfrak{P} . Putting everything together,

$$j(\mathfrak{p}E) \equiv j(\tilde{E}^{(p)}) \equiv j(E^{(\mathfrak{p}, L/K)}) \pmod{\mathfrak{P}}.$$

But we chose p so that nonisomorphic curves have j -invariants that are not congruent modulo p (item 3). Therefore, $\mathfrak{p}E \cong E^{(\mathfrak{p}, L/K)}$. This shows that the action of \mathfrak{p} is the same as the action of $(\mathfrak{p}, L/K)$, i.e. $F((\mathfrak{p}, L/K)) = [\mathfrak{p}]$.

Step 2: We show that $F : G(\overline{K}/K) \rightarrow \text{Cl}(\mathcal{O}_K)$ has kernel equal to $G(\overline{K}/K(j(E)))$, and so factors through $G(K(j(E))/K) \hookrightarrow \text{Cl}(\mathcal{O}_K)$. Indeed,

$$\begin{aligned} \ker(F) &= \{\sigma : F(\sigma)E = E\} \\ &= \{\sigma : E^\sigma = E\} && \text{definition of } \sigma \\ &= \{\sigma : j(E)^\sigma = j(E)\} && j \text{ parameterizes isomorphism classes} \\ &= G(\overline{K}/K(j(E))). \end{aligned}$$

We let $L = K(j(E))$.

Step 3: Let \mathfrak{f} be the conductor of L/K . We extend Step 1 to all ideals \mathfrak{a} : for all \mathfrak{a} we have

$$F((\mathfrak{a}, L/K)) = [\mathfrak{a}] \in \text{Cl}(\mathcal{O}_K);$$

in other words $\mathfrak{f} = 1$ and the following composition is the identity map.

$$\begin{array}{ccc} I_K/P_K & \xrightarrow{\psi_{L/K}} & G(L/K) \xrightarrow{F} \text{Cl}(\mathcal{O}_K). \\ & \searrow \cong & \nearrow \\ & & \text{Id} \end{array} \tag{17.9}$$

Given $\mathfrak{a} \in I_K^\mathfrak{f}$, there are infinitely many $\mathfrak{p} \in I_K^\mathfrak{f}$ in the same class as \mathfrak{a} with degree 1 by Corollary 16.16.3.6. Choose such a prime \mathfrak{p} , that does not divide a prime in S . Note $\mathfrak{a}, \mathfrak{p}$ differ by an ideal in $P_K(1, \mathfrak{f})$ so they have the same image by the Artin symbol. Step 1 shows that

$$F((\mathfrak{a}, L/K)) = F((\mathfrak{p}, L/K)) \stackrel{\text{Step 1}}{=} [\mathfrak{p}] = [\mathfrak{a}].$$

In particular, for any principal ideal $(\alpha) \in I_K^\mathfrak{f}$, we have $F(((\alpha), L/K)) = 1$. However, by definition the conductor is the smallest \mathfrak{p} such that $\alpha \equiv 1 \pmod{\mathfrak{f}}$ implies $((\alpha), L/K) = 1$, so we must have $\mathfrak{f} = (1)$.⁴ Thus the map $F : I_K^\mathfrak{f}/P_K(1, \mathfrak{f}) \rightarrow G(L/K)$ we had originally is actually just $F : I_K/P_K \rightarrow G(L/K)$, and we get (17.9).

Step 4: Since the conductor is divisible by exactly the ramifying primes, L/K is unramified, and $L \subseteq H_K$. On the other hand, the map $F \circ \psi_{L/K} : I_K/P_K \rightarrow \text{Cl}(\mathcal{O}_K)$ is an isomorphism because $F \circ \psi_{L/K}$ is just the identity map. This gives $[L : K] = |\text{Cl}(\mathcal{O}_K)| = [H_K : K]$. Hence $L = H_K$. This shows item 1.

⁴ Technically, we only have $((\alpha), L/K) = 1$ for $(\alpha) \perp \mathfrak{f}$, and a priori $((\alpha), L/K)$ is not defined for $(\alpha) \perp \mathfrak{f}$. (We don't know $\mathfrak{f} = 1$ yet.) The proper way to conclude $\mathfrak{f} = (1)$ is transfer the problem over to ideles: We know $\psi_{L/K}(P_K^\mathfrak{f}) = 1$, so $\phi_{L/K}(K^\times \cup_K^\mathfrak{f}) = 1$. By $\mathbb{I}_K^\mathfrak{f}/K(1, \mathfrak{f}) \cup_K(1, \mathfrak{f}) \cong \mathbb{I}_K/K^\times \cup_K(1, \mathfrak{f})$ we conclude that $\phi_{L/K}(K^\times \cup_K) = 1$. Hence $\mathfrak{f} = 1$.

Step 5: Item 3 now follows immediately, since we already showed $E^{\psi_{L/K}(\mathfrak{a})} = [\mathfrak{a}]E$ and we now know $\bar{L} = H_K$. Item 2 follows since the fact that the composition in (17.9) is an isomorphism means the map $F : G(L/K) \rightarrow \text{Cl}(\mathcal{O}_K)$ is surjective. Since F transfers the action of $G(L/K)$ on $\mathcal{E}ll_{\bar{\mathbb{Q}}}(\mathcal{O}_K)$ to $\text{Cl}(\mathcal{O}_K)$, and $\text{Cl}(\mathcal{O}_K)$ acts simply transitively on $\mathcal{E}ll_{\bar{\mathbb{Q}}}(\mathcal{O}_K)$, we get that the same is true for $G(L/K)$. \square

5 Maximal abelian extension

We next carry out part 2 of our outline in Section 4.1. We construct the ray class fields for K , then take their compositum to get the maximal abelian extension.

Definition 17.5.1: Suppose E has CM by an order in K , and E is defined over H_K . A **Weber function** is an isomorphism $h : E/\text{Aut}(1) \rightarrow \mathbb{P}^1$ defined over H_K . (So if $f : E \rightarrow E'$ is an automorphism, then $h(P) = h(f(P))$.)

We can always fix a concrete Weber function.

Example 17.5.2: The simplest Weber function is the following. If E has the form

$$y^2 = x^3 + Ax + B, \quad A, B \in H_K,$$

then take

$$h(P) = \begin{cases} x, & AB \neq 0 \\ x^2, & B = 0 \\ x^3, & C = 0. \end{cases}$$

In the 3 cases, respectively, $\text{Aut}(E)$ is 1, $\mathbb{Z}/2$ or $\mathbb{Z}/4$, and $\mathbb{Z}/3$ or $\mathbb{Z}/6$.

We can define a Weber function that is “model independent,” i.e. doesn’t change under if we change to an isomorphic elliptic curve, by

$$h(f(z)) = \begin{cases} \frac{g_2(\Lambda)g_3(\Lambda)}{\Delta(\Lambda)} \wp(z, \Lambda), & j(E) \neq 0, 1728 \\ \frac{g_2(\Lambda)^2}{\Delta(\Lambda)} \wp(z, \Lambda)^2, & j(E) = 1728 \\ \frac{g_3(\Lambda)}{\Delta(\Lambda)} \wp(z, \Lambda)^3, & j(E) = 0. \end{cases}$$

This is because the expressions have “weight 0.”

The importance of the Weber function is given below. It would not be true if $h(P)$ were just defined as $h(x, y) = x$.

Lemma 17.5.3. *Let E be an elliptic curve with CM by \mathcal{O} .*

1. *The extension $K(j(E), E_{\text{tors}})/K(j(E))$ is abelian.*
2. *The extension $K(j(E), h(E_{\text{tors}}))/K$ is abelian.*

The first statement is important because it tells us $G(\overline{K}/K(j(E)))$ acts in an abelian way on E_{tors} . Thus the “Galois representation” of the Galois group on E_{tors} is abelian. Thus, as we will see, it will decompose into two Grössencharacters.

Proof. We have an injective map $G(K(j(E), E[m])/K(j(E))) \hookrightarrow \text{Aut}(E[m])$.⁵ Now, the image of G in $\text{Aut}(E[m])$ commutes with \mathcal{O}_K , so is contained in

$$\text{Aut}_{\mathcal{O}_K/m\mathcal{O}_K}(E[m]) \cong \text{Aut}_{\mathcal{O}_K/m\mathcal{O}_K}(\mathcal{O}_K/m\mathcal{O}_K) \cong (\mathcal{O}_K/m\mathcal{O}_K)^\times$$

which is abelian.

For the second, suppose $\sigma, \tau \in G(K(j(E), h(E_{\text{tors}}))/K)$. We show that $\sigma\tau = \tau\sigma$. Since $K(j(E))/K$ is abelian, $\sigma\tau\sigma^{-1}\tau^{-1}$ fixes $j(E)$. Now $\sigma\tau\sigma^{-1}\tau^{-1}$ gives an automorphism of $E' = \tau\sigma(E)$ because

$$(\sigma\tau\sigma^{-1}\tau^{-1})\tau\sigma(E) = \sigma\tau(E) \cong \tau\sigma(E),$$

as the Galois action factors through $G(K^{\text{ab}}/K)$ and hence is abelian (Theorem 17.4.1) (alternatively, because $\sigma\tau\sigma^{-1}\tau^{-1}$ fixes $j(E)$). As E is defined over H_K , we actually have equality.

Since h is invariant under automorphism, for any $P \in E_{\text{tors}}$,

$$h(P) = h(\sigma\tau\sigma^{-1}\tau^{-1}P) = \sigma\tau\sigma^{-1}\tau^{-1}h(P).$$

(We know h is defined over H_K and $\sigma\tau\sigma^{-1}\tau^{-1}$ fixes $H_K = K(j(E))$.) Hence $\sigma\tau\sigma^{-1}\tau^{-1}$ fixes $h(E_{\text{tors}})$ as well, and $\sigma\tau\sigma^{-1}\tau^{-1} = 1$. \square

Theorem 17.5.4. *Suppose K is a quadratic imaginary field and E has CM by \mathcal{O}_K .*

1. *For an integral ideal \mathfrak{a} of \mathcal{O}_K , $L_{\mathfrak{a}} := H_K(h(E[\mathfrak{a}])) = K(j(E), h(E[\mathfrak{a}]))$ is the ray class field of K modulo \mathfrak{a} .*
2. *The maximal abelian extension of K is*

$$K(j(E), h(E_{\text{tors}})).$$

Proof. Step 1: We need the following lemma.

Lemma 17.5.5. *Suppose E is an elliptic curve defined over L with CM by \mathcal{O}_K , and has good reduction at \mathfrak{P} . Let \tilde{E} be the reduction modulo \mathfrak{P} . Let $\theta : \text{End}(E) \rightarrow \text{End}(\tilde{E})$ be the reduction map on endomorphisms. Then for any $\gamma \in \text{End}(\tilde{E})$,*

$$\gamma \in \text{im}(\theta) \iff \gamma \text{ commutes with every element in } \text{im}(\theta).$$

Proof. Since E has good reduction, the map $\text{End}(E) \hookrightarrow \text{End}(\tilde{E})$ is injective. Consider 2 cases.

⁵Since $E[m] = \mathbb{Z}/m \times \mathbb{Z}/m$, if we choose a basis for $E[m]$, we have $\text{Aut}(E[m]) \cong \text{GL}_2(\mathbb{Z}/m)$, so we have a Galois representation.

1. $\text{End}(\widetilde{E})$ is a quadratic order. Then $\text{End}(E) = \text{End}(\widetilde{E})$ (as $\text{End}(E)$ is a maximal order) so this case is clear.
2. $\text{End}(\widetilde{E})$ is an order in a quaternion algebra. Then $\text{End}(E) \otimes \mathbb{Q}$ is its own centralizer in the quaternion algebra $\text{End}(\widetilde{E}) \otimes \mathbb{Q}$, by the Double Centralizer Theorem 13.13.4.11.

□

Step 2: We show that in general, we can lift the Frobenius map.

Proposition 17.5.6: Suppose E has CM by \mathcal{O}_K and is defined over H_K . Let $\mathfrak{P} \mid \mathfrak{p} \mid p$ in H_K, K, \mathbb{Q} , respectively, with \mathfrak{p} having degree 1 and $p \notin S$, S being defined as in the proof of Theorem 17.4.4. Then the p th power Frobenius map can be lifted to a map on E , i.e. there is λ making the following commute:

$$\begin{array}{ccc} E & \xrightarrow{\lambda} & E^{(\mathfrak{p}, H_K/K)} \\ \downarrow & & \downarrow \\ \widetilde{E} & \xrightarrow{\widetilde{\lambda}=\phi_p} & \widetilde{E}^{(p)}. \end{array}$$

Moreover, if E corresponds to the complex torus \mathbb{C}/Λ , then up to isomorphism, λ corresponds to the map $\mathbb{C}/\Lambda \rightarrow \mathbb{C}/\mathfrak{p}^{-1}\Lambda$. (Recall that $E^{(\mathfrak{p}, H_K/K)} \cong \mathfrak{p}E$ by Theorem 17.4.4.)

Proof. We need to show ϕ_p is the reduction of some map; we do this by first reducing the problem to showing a certain endomorphism is in the image of θ and then showing the conditions of the previous lemma hold.

Again we use (17.8): $\widetilde{\phi}_1 : \widetilde{E} \rightarrow \widetilde{\mathfrak{p}E}$ is “like” the Frobenius map. We know $\widetilde{\phi}_1$ is the reduction of a map, namely the map $\phi_1 : E \rightarrow \mathfrak{p}E$. Now note $\widetilde{\mathfrak{p}E} \cong \widetilde{E^{(\mathfrak{p}, L/K)}} = \widetilde{E}^{(p)}$, the first from Thm 17.4.4 and the second from definition of the Frobenius element.

Let $\sigma = (\mathfrak{p}, L/K)$. It remains to show that $\varepsilon : \widetilde{E}^\sigma \rightarrow \widetilde{\mathfrak{p}E} \cong \widetilde{E}^\sigma$ is the reduction of a map ε' , because then $\varepsilon'^{-1} \circ \phi_1$ will be the desired map. Let $[\widetilde{\alpha}] \in \text{Aut}(\widetilde{E}^\sigma)$ be the reduction of a map $[\alpha]$. To show ε commutes with $[\alpha]$, we consider $\phi_1 = \varepsilon \circ \phi_p$, and consider how $[\alpha]$ “commutes” with $\widetilde{\phi}_1$ and ϕ_p .

1. $\widetilde{\phi}_1$: By normalization (Proposition 17.2.2(3)), we know

$$\phi_1 \circ [\alpha]_E = [\alpha]_{E^\sigma} \circ \phi_1.$$

2. ϕ_p : Note that for any morphism of varieties $f : V \rightarrow W$ over a field of characteristic p , the following commutes, where ϕ_V, ϕ_W are the p th power Frobenius maps on V and W :

$$\begin{array}{ccc} V & \xrightarrow{f} & W & \phi_W \circ f = f^\sigma \circ \phi_V. \\ \downarrow \phi_V & & \downarrow \phi_W & \\ V^{(p)} & \xrightarrow{f^\sigma} & W^{(p)} & \end{array}$$

Applying this to $[\alpha]_E$,

$$\phi_p \circ \widetilde{[\alpha]_E} = \widetilde{[\alpha]_E^\sigma} \circ \phi_p = \widetilde{[\alpha]_{E^\sigma}} \circ \phi_p,$$

where in the last step we used Theorem 17.2.2(4), noting $\sigma(\alpha) = \alpha$ since $\alpha \in K$ and $\sigma \in G(H_K/K)$.

Hence

$$\widetilde{[\alpha]_{E^\sigma}} \circ \underbrace{\varepsilon \circ \phi_p}_{\phi_1} \stackrel{1}{=} \varepsilon \circ \phi_p \circ \widetilde{[\alpha]_E} \stackrel{2}{=} \varepsilon \circ \widetilde{[\alpha]_{E^\sigma}} \circ \phi_p.$$

Cancelling ϕ_p gives $\widetilde{[\alpha]_{E^\sigma}} \circ \varepsilon = \varepsilon \circ \widetilde{[\alpha]_{E^\sigma}}$, so Lemma 17.5.5 shows ε is the reduction of some ε' , as needed.

To finish, note that ϕ_1 does indeed correspond to $\mathbb{C}/\Lambda \rightarrow \mathbb{C}/\mathfrak{p}^{-1}\Lambda$. Hence λ corresponds to $\mathbb{C}/\Lambda \rightarrow \mathbb{C}/\mathfrak{p}^{-1}\Lambda$, up to some automorphism. \square

Step 3: When $(\mathfrak{p}, H_K/K) = 1$, λ is just an endomorphism of E , hence equals $[\alpha]$ for some α . In fact, the following proposition shows it is $[\pi]$ for some π generating \mathfrak{p} , so that multiplication by π corresponds to the p th power Frobenius in the reduction.

Proposition 17.5.7: Suppose E has CM by \mathcal{O}_K and is defined over H_K . For all but finitely many degree 1 prime ideals \mathfrak{p} with $(\mathfrak{p}, H_K/K) = 1$ (equivalently, such that \mathfrak{p} is principal), there exists a unique π such that $\mathfrak{p} = (\pi)$ and the following commutes.

$$\begin{array}{ccc} E & \xrightarrow{[\pi]} & E \\ \downarrow & & \downarrow \\ \widetilde{E} & \xrightarrow{\phi_p} & \widetilde{E}. \end{array}$$

Proof. Since $(\mathfrak{p}, H_K/K) = 1$, Proposition 17.5.6 gives a diagram

$$\begin{array}{ccc} E & \xrightarrow{\lambda} & E \\ \downarrow & & \downarrow \\ \widetilde{E} & \xrightarrow{\phi_p} & \widetilde{E}. \end{array}$$

for some λ . We know λ is in the form $[\pi]$, and show π satisfies the desired conditions. We have by Proposition 17.2.9 that

$$\mathrm{Nm}_{K/\mathbb{Q}}(\pi) = \deg([\pi]) = \deg(\phi) = p = \mathfrak{N}\mathfrak{p}$$

so either $(\pi) = \mathfrak{p}$ or $(\pi) = \bar{\mathfrak{p}}$. As always, when we're deciding between conjugates, normalization comes to the rescue. Take $\omega \in \Omega_E$ whose reduction modulo \mathfrak{P} is nonzero. Normalization says that $[\pi]^*\omega = \pi\omega$ so

$$\tilde{\pi}\tilde{\omega} = \widetilde{[\pi]^*\omega} = \phi_p^*\tilde{\omega} = 0,$$

the last step since the Frobenius map is inseparable. We get $\mathfrak{P} \mid \pi$, forcing $(\pi) = \mathfrak{p}$.

For uniqueness, note the map

$$\mathcal{O}_K \xrightarrow[\cong]{[\cdot]} \text{End}(E) \xrightarrow{\tilde{E}} \text{End}(\tilde{E})$$

is injective for E having good reduction at \mathfrak{P} (Theorem 17.4.3). □

Step 4: Consider (17.7). We need to show that $P_K(1, \mathfrak{a})$ is exactly the kernel of the Artin map $\psi_{L_{\mathfrak{a}}/K}$. Note that $P_K(1, \mathfrak{a})$ and $\ker(\psi_{L_{\mathfrak{a}}/K})$ are both subgroups of $P_K^{\mathfrak{a}} = \ker(\psi_{H_K/K}) = \ker(\psi_{L_{\mathfrak{a}}/K}(\bullet)|_{H_K})$. It suffices to show that for all but finitely many primes \mathfrak{p} of degree 1 such that $(\mathfrak{p}, H_K/K) = 1$, we have $\mathfrak{p} \in P_K(1, \mathfrak{a})$ iff $\mathfrak{p} \in \ker(\psi_{L_{\mathfrak{a}}/K})$.

Let \mathfrak{p} satisfy the conditions of Proposition 17.5.7. Since the reduction of $\psi_{L/K}(\mathfrak{p})$ is the Frobenius map, we get that $\psi_{L/K}(\mathfrak{p}) = [\pi]$, for some π such that $(\pi) = \mathfrak{p}$.⁶ Since $(\mathfrak{p}, H_K/K) = 1$, we have the commutative diagram

$$\begin{array}{ccc} \psi_{L/K}(\mathfrak{p})=[\pi] \\ \tilde{E} & \xrightarrow{\quad} & \tilde{E} \\ \downarrow & & \downarrow \\ \tilde{E} & \xrightarrow{\phi_p} & \tilde{E} \end{array} \tag{17.10}$$

We have the following string of equivalences, for all but finitely many degree 1 primes \mathfrak{p} with $(\mathfrak{p}, H_K/K) = 1$,

1. $\mathfrak{p} \in P_K(1, \mathfrak{a})$.
2. $\mathfrak{p} = (\pi)$ where $\pi = u\alpha$ where u is a unit and $\alpha \equiv 1 \pmod{\mathfrak{a}}$.
3. For all \mathfrak{a} -torsion points $P \in E[\mathfrak{a}]$, $h([\pi]P) = h(P)$.
- 3'. For all \mathfrak{a} -torsion points $P \in \tilde{E}[\mathfrak{a}]$, $\tilde{h}([\pi]\tilde{P}) = \tilde{h}(\tilde{P})$.
4. $(\mathfrak{p}, L_{\mathfrak{a}}/K)$ fixes $h(E[\mathfrak{a}])$.
5. $\mathfrak{p} \in \ker(\psi_{L_{\mathfrak{a}}/K})$.

(1) \iff (2) is clear.

For (2) \implies (3), note that for all \mathfrak{a} torsion points $P \in E[\mathfrak{a}]$,

$$\begin{aligned} h([\pi]P) &= h([u][\alpha]P) \\ &= h([\alpha]P) && h \text{ is Aut}(E)\text{-invariant} \\ &= h(P) && \alpha \equiv 1 \pmod{\mathfrak{a}} \text{ and } P \in E[\mathfrak{a}]. \end{aligned}$$

Note it is important that h be $\text{Aut}(E)$ -invariant.

⁶Note the analogy with the cyclotomic case. $\psi_{L/K}(\mathfrak{p})$ acts on torsion points as $[\pi]$, just as in the cyclotomic case it acted as the p th power map, that corresponds to $[p]$ if we consider the natural map $\mathbb{Z} \rightarrow \text{End}(\mathbb{Q}(\zeta_n))$.

For (3') \implies (2), let $P \in E[\mathfrak{a}]$ be a torsion point. By [Sil86, VII.3.1b], $E[\mathfrak{a}] \hookrightarrow \widetilde{E}[\mathfrak{a}]$ is injective for $\mathfrak{p} \nmid \mathfrak{a}$ and E with good reduction at \mathfrak{p} . Since h is an isomorphism (in particular, an injection) $E/\text{Aut}(E) \rightarrow \mathbb{P}^1$, we get that $[\pi]P = [u]P$ for some $[u] \in \text{Aut}(E)$. But $E[\mathfrak{a}] \cong \mathcal{O}_K/\mathfrak{a}$, so we can choose u such that $\pi \equiv u \pmod{\mathfrak{a}}$. Then there exists α such that $\pi = u\alpha$, with $\alpha \equiv 1 \pmod{\mathfrak{a}}$.

For (3) \implies (4), we calculate the action of $(\mathfrak{p}, L/K)$ on a torsion point $P \in E[\mathfrak{a}]$, in the reduced curve:

$$\widetilde{P^{(\mathfrak{p}, L/K)}} = \phi_p(\widetilde{P}) = \widetilde{[\pi]P},$$

the second equality from Proposition 17.5.7. This allows us to understand the action on the nonreduced curve, since $E[\mathfrak{a}] \hookrightarrow \widetilde{E}[\mathfrak{a}]$ is injective for $\mathfrak{p} \nmid \mathfrak{a}$ and \mathfrak{p} of good reduction. We get

$$P^{(\mathfrak{p}, L/K)} = [\pi]P.$$

Thus (3) implies

$$\begin{aligned} h(P)^{(\mathfrak{p}, L/K)} &= h(P^{(\mathfrak{p}, L/K)}) && (\mathfrak{p}, L/K) \text{ fixes } H_K \text{ and } E \text{ defined over } H_K \\ &= h([\pi]P) \\ &= h(P) && \text{by (3)}. \end{aligned}$$

Now we prove (4) \implies (3'). Let $\sigma \in G(\overline{K}/K)$ be an automorphism such that $\sigma|_{K^{\text{ab}}} = (\mathfrak{p}, K^{\text{ab}}/K)$. Then for any $P \in E[\mathfrak{a}]$,

$$\widetilde{h}([\pi]\widetilde{P}) \stackrel{(17.10)}{=} \widetilde{h}(\phi(\widetilde{P})) = \widetilde{h}(P^\sigma) = \widetilde{h}(P)^\sigma = \widetilde{h}(\widetilde{P}),$$

the last two equalities since $\sigma|_H = 1$, h is defined over H , and $\sigma|_{L_a}$ fixes $h(E[\mathfrak{a}])$ by assumption. Thus (3') holds.

Now (4) \iff (5) comes from the fact that (\mathfrak{p}, L_a, K) already fixes $K(j(E))$, so to fix L_a it only needs to fix $h(E[\mathfrak{a}])$.

Step 7: The maximal abelian extension is the union of the all ray class fields. Note every \mathfrak{c} divides n for some n so we can just restrict to ray class fields corresponding to (n) for some $n \in \mathbb{N}$:

$$K^{\text{ab}} = \bigcup_n K(j(E), h(E[n])) = K(j(E), h(E_{\text{tors}})).$$

□

6 The Main Theorem of Complex Multiplication

Given $\sigma \in \text{Aut}(\mathbb{C}/K)$, consider the map $\sigma : E(\mathbb{C}) \rightarrow E^\sigma(\mathbb{C})$. We would like to know how this map acts on torsion points. This is since to get Galois representations of elliptic curves, we look at how σ acts on torsion points—often specializing to torsion points that are a power of a prime.

Because we are considering CM elliptic curves, we can identify the torsion points with K/\mathfrak{a} , for some ideal \mathfrak{a} . Namely, given an analytic isomorphism $f : \mathbb{C}/\mathfrak{a} \xrightarrow{\cong} E(\mathbb{C})$, we can restrict it to K/\mathfrak{a} to get

$$f|_{K/\mathfrak{a}} : K/\mathfrak{a} \xrightarrow{\cong} E_{\text{tors}} \hookrightarrow E(\mathbb{C}).$$

The main theorem of complex multiplication tells us we can transfer the map $\sigma : E(\mathbb{C}) \rightarrow E^\sigma(\mathbb{C})$ via an *analytic isomorphism* to a multiplication-by-an-idele map $[\mathbf{x}^{-1}] : K/\mathfrak{a} \rightarrow K/\mathbf{x}^{-1}\mathfrak{a}$, where \mathbf{x} and σ are related in terms of the Artin map (to be made precise).

Definition 17.6.1: Let $\mathbf{x} = \prod_{\mathfrak{p} \in V_K^0} \mathfrak{p}^{m(\mathfrak{p})} \prod_{v \in V_K^\infty} v^{m(v)} \in \mathbb{I}_K$ be an idele. Let \mathfrak{a} be an ideal, and define $\mathbf{x}\mathfrak{a}$ by

$$\mathbf{x}\mathfrak{a} = p(\mathbf{x})\mathfrak{a} = \left(\prod_{\mathfrak{p} \in V_K} \mathfrak{p}^{m(\mathfrak{p})} \right) \mathfrak{a}.$$

Define the map

$$[\mathbf{x}] : K/\mathfrak{a} \rightarrow K/\mathbf{x}\mathfrak{a} \tag{17.11}$$

as follows. Note $K/\mathfrak{a} \cong \prod_{\mathfrak{p}} K_{\mathfrak{p}}/\mathfrak{a}K_{\mathfrak{p}}$ by the Chinese Remainder Theorem, where x is just identified with its images in the $K_{\mathfrak{p}}/\mathfrak{a}K_{\mathfrak{p}}$: $(x_{\mathfrak{p}})_{\mathfrak{p} \in V_K^0}$. Then (17.11) sends

$$(a_{\mathfrak{p}}) \mapsto (x_{\mathfrak{p}}a_{\mathfrak{p}}) \text{ where } \mathbf{x} = (x_{\mathfrak{p}}). \tag{17.12}$$

Theorem 17.6.2 (Main Theorem of Complex Multiplication). *Suppose E is an elliptic curve with CM by \mathcal{O}_K . Let $\sigma \in \text{Aut}(\mathbb{C}/K)$ and $\mathbf{x} \in \mathbb{I}_K$ be such that*

$$\sigma|_{K^{ab}} = \phi_K(\mathbf{x}).$$

Fix an analytic isomorphism $f : \mathbb{C}/\mathfrak{a} \xrightarrow{\cong} E(\mathbb{C})$. Then there exists a unique analytic isomorphism $f' : K/\mathbf{x}^{-1}\mathfrak{a} \rightarrow E^\sigma(\mathbb{C})$ such that the following commutes:

$$\begin{array}{ccc} K/\mathfrak{a} & \xrightarrow{\mathbf{x}^{-1}} & K/\mathbf{x}^{-1}\mathfrak{a} \\ \downarrow f & & \downarrow f' \\ E(\mathbb{C}) & \xrightarrow{\sigma} & E^\sigma(\mathbb{C}). \end{array}$$

Remark 17.6.3: The map (17.12) can be a bit weird to think about: For instance, consider the simpler case $K = \mathbb{Q}$, $\mathfrak{a} = \mathbb{Z}$. Take the idele \mathbf{x} with 1's everywhere except $x_5 = 2$. Then $[\mathbf{x}]$ sends $\frac{1}{2} \mapsto \frac{1}{2}, \frac{1}{3} \mapsto \frac{1}{3}, \frac{1}{7} \mapsto \frac{1}{7}$ and so forth but sends $\frac{1}{5} \mapsto \frac{2}{5}$. So it is surprising that $\mathbf{x}^{-1} : K/\mathfrak{a} \rightarrow K/\mathbf{x}^{-1}\mathfrak{a}$ can be related analytically to $E(\mathbb{C}) \rightarrow E^\sigma(\mathbb{C})$.

Compare this theorem to Proposition 17.5.7. Rather than just dealing with the Frobenius element of a prime, we deal with the Artin map of an idele.

Proof. Note uniqueness follows from the fact that topologically, the closure of $K/\mathbf{x}^{-1}\mathfrak{a}$ is $\mathbb{C}/\mathbf{x}^{-1}\mathfrak{a}$, and any continuous function is determined by its values on a dense set.

First we prove this for E defined over $\mathbb{Q}(j(E))$ and \mathfrak{a} integral. We do this in 2 steps. Step 1: Approximate σ by a field automorphism λ that is the Frobenius element of a prime \mathfrak{p} . (The Frobenius element is something much more concrete to work with than the abstract Artin map of an idele.) We will take better and better approximations, which determine the action on $E[m]$ for larger and larger m , and take an inverse limit.

So let L'_m be the Galois closure of $K(j(E), E[m])/K$. By Corollary ??, there are infinitely many primes with $\mathfrak{P} \mid \mathfrak{p}$ in K and L such that

$$(\mathfrak{P}, L/K) = \sigma|_{L'_m}, \quad \mathfrak{N}(\mathfrak{p}) = 1.$$

We can furthermore choose \mathfrak{p} satisfying the following, because each condition excludes only finitely many primes.

1. \mathfrak{p} is unramified in L'_m .
2. $\mathfrak{p} \notin S$, where S is defined as in the proof of Theorem 17.4.4.
3. $\mathfrak{p} \nmid m$.

By Proposition 17.5.6, there exists a map $\lambda : E \rightarrow E^\sigma$ that reduces to ϕ_p modulo \mathfrak{P} . On $\widetilde{E}[m]$, both λ and σ act as ϕ_p . Because $\mathfrak{P} \nmid m$ by item 3, the reduction map modulo \mathfrak{P} , $E[m] \rightarrow \widetilde{E}[m]$, is injective. Hence λ and σ act the same on $E[m]$:

$$\lambda|_{E[m]} = \sigma|_{E[m]} : E[m] \rightarrow E^\sigma[m]. \tag{17.13}$$

But we know how the map λ acts: Proposition 17.5.6 tells us that the map $\lambda : E \rightarrow E^\sigma$ corresponds to the map on complex tori $i : \mathbb{C}/\mathfrak{a} \rightarrow \mathbb{C}/\mathfrak{p}^{-1}\mathfrak{a}$.⁷ Hence we have the commutative diagram

$$\begin{array}{ccc} \mathbb{C}/\mathfrak{a} & \xrightarrow{i} & \mathbb{C}/\mathfrak{p}^{-1}\mathfrak{a} \\ \downarrow f & & \downarrow f'' \\ E(\mathbb{C}) & \xrightarrow{\lambda} & E^\sigma(\mathbb{C}) \end{array} \tag{17.14}$$

for some analytic isomorphism f'' .

Step 2: By Theorem 17.5.4, the ray class group modulo m is $K_m = K(j(E), h(E[m]))$. Note $\overline{K_m} \subseteq L'_m$. Now by assumption, \mathfrak{p} was chosen so that the images of \mathfrak{p} and \mathfrak{x} under the Artin map both project to $\sigma|_{K_m}$:

$$\phi_{K_m/K}(\mathfrak{x}) = \sigma|_{K_m} = \psi_{K_m/K}(\mathfrak{p}) = \phi_{K_m/K}(i_{\mathfrak{p}}(\pi))$$

where ψ, ϕ denote the Artin map on ideals and on ideles, respectively, and π is the uniformizer of \mathfrak{p} in $K_{\mathfrak{p}}$. We have

$$\ker \psi_{K_m/K} = K^\times \mathbb{U}_K(1, m).$$

⁷The map σ and \mathfrak{x}^{-1} appearing in the theorem statement are bijections, while λ and i are not. This is okay, though, because we only use λ, i to approximate σ on m -torsion, and λ, i are injective on m -torsion, since $\mathfrak{P} \nmid m$.

(See Definition 11.11.5.8 for notation.) This follows from the definition of the ray class field and from the correspondence between ray class groups in Definition 11.11.4.5 and idele class groups in Example 11.11.5.10. We have $\mathbf{x} \in i_{\mathfrak{p}}(\pi) \ker \phi_{K_m/K}$, giving

$$\mathbf{x} = \alpha \cdot i_{\mathfrak{p}}(\pi) \cdot \mathbf{u}, \quad \alpha \in K^\times, \quad \mathbf{u} \in \mathbb{U}_K(1, m).$$

We now compose (17.14) with the homothety α^{-1} , and note $(\mathbf{x}) = (\alpha)\mathfrak{p}$, to get the desired map $\mathbb{C}/\mathbf{x}^{-1}\mathfrak{a} \rightarrow E^\sigma(\mathbb{C})$:

$$\begin{array}{ccccc} \mathbb{C}/\mathfrak{a} & \xrightarrow{i} & \mathbb{C}/\mathfrak{p}^{-1}\mathfrak{a} & \xrightarrow{\alpha^{-1}} & \mathbb{C}/\mathbf{x}^{-1}\mathfrak{a} \\ \downarrow f & & \downarrow f'' & \swarrow f'_m & \\ E(\mathbb{C}) & \xrightarrow{\lambda} & E^\sigma(\mathbb{C}) & & \end{array} \quad (17.15)$$

Here, $f'_m(z) := f''(\alpha z)$.

This isn't quite what we want yet, though, because the top row is the map α^{-1} rather than the map \mathbf{x}^{-1} . We need to show that for m -torsion points, α^{-1} acts the same as \mathbf{x}^{-1} . Then we would have

$$\sigma(f(t)) = \lambda(f(t)) = f'_m(\alpha^{-1}t) = f'_m(\mathbf{x}^{-1}t), \quad t \in m^{-1}\mathfrak{a}/\mathfrak{a}.$$

The first equality is since σ, λ were by construction the same on $E[m]$ (17.13), so $\sigma \circ f$ and $\lambda \circ f$ are the same on $m^{-1}\mathfrak{a}/\mathfrak{a}$. The second is by commutativity of (17.15).

To show the third equality, we note that

$$\begin{array}{llll} & f'_m(\alpha^{-1}t) = f'_m(\mathbf{x}^{-1}t) & \text{for all } t \in m^{-1}\mathfrak{a}/\mathfrak{a} \\ (f'_m \text{ bijective}) & \iff \alpha^{-1}t - \mathbf{x}^{-1}t \in \mathfrak{a} & \text{for all } t \in m^{-1}\mathfrak{a} \\ & \iff \alpha^{-1}t_{\mathfrak{q}} - x_{\mathfrak{q}}^{-1}t_{\mathfrak{q}} \in \mathfrak{a}_{\mathfrak{q}} & \text{for all } t \in m^{-1}\mathfrak{a}, \mathfrak{q} \\ (\text{multiplying by } x_{\mathfrak{q}} = \alpha[i_{\mathfrak{p}}(\pi)]_{\mathfrak{q}}u_{\mathfrak{q}}) & \iff [i_{\mathfrak{p}}(\pi)]_{\mathfrak{q}}u_{\mathfrak{q}}t - t \in \mathfrak{a}_{\mathfrak{q}} & \text{for all } t \in m^{-1}\mathfrak{a}_{\mathfrak{q}} \\ & \iff ([i_{\mathfrak{p}}(\pi)]_{\mathfrak{q}}u_{\mathfrak{q}} - 1)\mathfrak{a}_{\mathfrak{q}} \subseteq m\mathfrak{a}_{\mathfrak{q}} \\ u_{\mathfrak{q}} \in \mathbb{U}_K(1, m) & \iff ([i_{\mathfrak{p}}(\pi)]_{\mathfrak{q}} - 1)\mathfrak{a}_{\mathfrak{q}} \subseteq m\mathfrak{a}_{\mathfrak{q}}. \end{array}$$

Consider 2 cases.

1. $\mathfrak{q} \neq \mathfrak{p}$: In this case, $[i_{\mathfrak{p}}(\pi)]_{\mathfrak{q}} = 1$, so this is trivial.
2. $\mathfrak{q} = \mathfrak{p}$: $[i_{\mathfrak{p}}(\pi)]_{\mathfrak{p}} = \pi$, and $\pi - 1$ is a unit. By assumption $\mathfrak{p} \nmid m$, hence $(\pi - 1)\mathfrak{a} = \mathfrak{a} = m\mathfrak{a}$.

Step 3: We now show that the maps f'_m are all actually the same for $m \geq 3$. Indeed, $f'_m|_{E[m]} = f'_{mn}|_{E[m]}$ by construction, so f'_m, f'_{mn} differ by an automorphism that fixes $E[m]$. This automorphism must be $[\zeta]$ for some element of norm 1 in K , and $f'_m = [\zeta] \circ f'_{mn}$. Since f'_m, f'_{mn} are isomorphisms, this says

$$E[m] \subseteq \ker[1 - \zeta]$$

The only possibilities are ζ a 4th or 6th root of unity, and if $\zeta \neq 1$, then $[1 - \zeta]$ has norm at most 4. So for $m \geq 3$, $\zeta = 1$, and $f'_m = f'_{mn}$.

Step 4: Finally, we show the theorem holds for general E/L . Any elliptic curve E has a model E' defined over $M' = \mathbb{Q}(j(E))$, corresponding to a complex torus \mathbb{C}/\mathfrak{a}' with \mathfrak{a}' an integral ideal (see the left face below). Let $E \rightarrow E'$ be an isomorphism and $K/\mathfrak{a} \rightarrow K/\mathfrak{a}'$ be the corresponding map on torsion. Then the existence of $f'_{E'}$ for E'/L gives the existence of f'_E for E/L , by choosing f'_E to make the below diagram commute.

$$\begin{array}{ccccc}
 K/\mathfrak{a} & \xrightarrow{\mathbf{x}^{-1}} & K/\mathbf{x}^{-1}\mathfrak{a} & & \\
 \downarrow f_E & \searrow \cong & \downarrow \cong & & \\
 & & K/\mathfrak{a}' & \xrightarrow{\mathbf{x}^{-1}f'_E} & K/\mathbf{x}^{-1}\mathfrak{a}' \\
 & & \downarrow \sigma_{f_{E'}} & \downarrow \cong & \downarrow f'_{E'} \\
 E(\mathbb{C}) & \xrightarrow{\sigma_{f_{E'}}} & E^\sigma(\mathbb{C}) & & \\
 \searrow & \downarrow & \downarrow \cong & & \\
 & & E'(\mathbb{C}) & \xrightarrow{\quad} & E'^\sigma(\mathbb{C}).
 \end{array}$$

□

6.1 The associated Grössencharacter

The Main Theorem involved 2 different elliptic curves, and 2 different analytic isomorphisms. In the special case that σ fixes E , the curves will be the same, and by nudging the map upstairs by a constant depending on \mathbf{x} , we can restate the theorem using a consistent choice of f . (Compare to how we specialized from Proposition 17.5.6 to 17.5.7.) The action of $\phi_L(\mathbf{x})$ on the elliptic curve will “essentially” correspond to multiplication by $\chi_{E/L}$ on K/\mathfrak{a} .

Theorem 17.6.4 (Grössencharacter of an elliptic curve). *Let E/L be an elliptic curve with complex multiplication by \mathcal{O}_K , and suppose $K \subseteq L$. Let $\mathbf{x} \in \mathbb{I}_L$ and $\mathbf{y} = \text{Nm}_{L/K}(\mathbf{x}) \in \mathbb{I}_K$. Then there exists a unique $\alpha = \alpha_{E/L}(\mathbf{x}) \in K^\times$ with the following properties.*

1. $\alpha\mathcal{O}_K = (\mathbf{y})$.
2. For any fractional ideal $\mathfrak{a} \subseteq K$ and any analytic isomorphism $f : \mathbb{C}/\mathfrak{a} \rightarrow E(\mathbb{C})$, the following commutes.

$$\begin{array}{ccc}
 K/\mathfrak{a} & \xrightarrow{\alpha\mathbf{y}^{-1}} & K/\mathfrak{a} \\
 \downarrow f & & \downarrow f \\
 E(L^{ab}) & \xrightarrow{\phi_L(\mathbf{x})} & E(L^{ab}).
 \end{array}$$

Moreover, defining $\chi_{E/L} : \mathbb{I}_L \rightarrow \mathbb{C}^\times$ by

$$\chi_{E/L}(\mathbf{x}) := \alpha_{E/L}(\mathbf{x})[\mathrm{Nm}_{L/K}(\mathbf{x}^{-1})]_\infty,$$

$\chi_{E/L}$ is a Grössencharacter of K , and $\chi_{E/L}$ is ramified at \mathfrak{P} (i.e. $\chi_{E/L}(U_{\mathfrak{P}})$ is not identically 1) iff E has bad reduction at \mathfrak{P} .

Proof. Part 1: Since f is an isomorphism, uniqueness is clear. To construct α , choose any $\sigma \in \mathrm{Aut}(\mathbb{C}/L)$ such that $\sigma|_{L^{\mathrm{ab}}} = \phi_L(\mathbf{x})$. We use Theorem 17.6.2 with σ and $\mathbf{y} \in \mathbb{I}_K$, noting the following points.

1. $E^\sigma = E$ since E is defined over L and σ fixes L .
2. The image of f is contained in $E(L^{\mathrm{ab}})$ as $E_{\mathrm{tors}} \in E(L^{\mathrm{ab}})$ by Lemma 17.5.3.
3. By compatibility of the Artin map, $\phi_L(\mathbf{x})|_{K^{\mathrm{ab}}} = \phi_K(\mathrm{Nm}_{L/K} \mathbf{x}) = \phi_K(\mathbf{y})$.

We obtain an analytic map f' making the following commute.

$$\begin{array}{ccc} K/\mathfrak{a} & \xrightarrow{\mathbf{y}^{-1}} & K/\mathbf{y}^{-1}\mathfrak{a} \\ \downarrow f & & \downarrow f' \\ E(L^{\mathrm{ab}}) & \xrightarrow{\phi_L(\mathbf{x})} & E(L^{\mathrm{ab}}). \end{array}$$

Because

$$\mathbb{C}/\mathbf{y}^{-1}\mathfrak{a} \cong E^\sigma(\mathbb{C}) \cong E(\mathbb{C}) \cong \mathbb{C}/\mathfrak{a},$$

we have that $\mathbf{y}^{-1}\mathfrak{a}$ is homothetic to \mathfrak{a} , i.e. there exists β so that β takes $K/\mathbf{y}^{-1}\mathfrak{a}$ back to K/\mathfrak{a} . Defining $f''(x) = f'(\beta^{-1}x)$, we have that it differs from f by some automorphism $[\zeta]$: $f \circ [\zeta] = f''$. Let $\alpha = \beta\zeta$. Then we can extend the above diagram as follows.

$$\begin{array}{ccccc} K/\mathfrak{a} & \xrightarrow{\mathbf{y}^{-1}} & K/\mathbf{y}^{-1}\mathfrak{a} & \xrightarrow{\alpha} & K/\mathfrak{a} \\ \downarrow f & & \downarrow f' & \swarrow f & \\ E(L^{\mathrm{ab}}) & \xrightarrow{\phi_L(\mathbf{x})} & E(L^{\mathrm{ab}}) & & \end{array}$$

As $\alpha\mathbf{y}^{-1}\mathfrak{a} = \mathfrak{a}$, we get $(\alpha) = (\mathbf{y})$.

To see that α is independent of f and the ideal \mathfrak{a} , let f' be another analytic isomorphism $K/\mathfrak{a}' \rightarrow E(L^{\mathrm{ab}})$. Let the map $K/\mathfrak{a}' \rightarrow K/\mathfrak{a}$ be multiplication-by- γ . Then $f(\gamma x)$ is also an analytic isomorphism $K/\mathfrak{a}' \rightarrow E(L^{\mathrm{ab}})$. Hence $\gamma^{-1}f^{-1} \circ f'$ is an automorphism $[\zeta]$ of K/\mathfrak{a}' , i.e. $f'(x) = f([\zeta]\gamma x)$. Thus $\phi_L(\mathbf{x})[f'(x)] = f'(\alpha\mathbf{y}^{-1}x)$ as well.

Part 2: $\alpha_{E/L}$ and hence $\chi_{E/L}$ is a homomorphism since it's clear that $\phi_L(\mathbf{x}\mathbf{x}') \circ f = f \circ \alpha\alpha'\mathbf{y}\mathbf{y}'^{-1}$, and $\phi_L(\mathbf{x}^{-1}) \circ f = f \circ \alpha^{-1}\mathbf{y}$.

We need to check that $\chi_{E/L}(L^\times) = 1$ and that $\chi_{E/L}$ factors through a modulus.

For the first point, note $\phi_L(L^\times) = 1$, the identity element of $G(L^{\text{ab}}/L)$. Let $i : K^\times \rightarrow \mathbb{I}_K$, $L^\times \rightarrow \mathbb{I}_L$ be the diagonal maps, and suppose $\mathbf{x} = i(x)$. We have $\mathbf{y} = \text{Nm}_{L/K}(i(x)) = i(\text{Nm}_{L/K}(x))$. Then α is just the element such that $\alpha \text{Nm}_{L/K}(x)^{-1}$ induces the identity map, i.e. $\alpha = \text{Nm}_{L/K}(x) = [\text{Nm}_{L/K} \mathbf{x}]_\infty$, so $\chi_{E/L}(\mathbf{x}) = 1$.

For the second point, fix $m \geq 3$ ($m = 3$ works fine). We'll show that for any idele \mathbf{x} in a small enough open subset of finite index, $\phi_L(\mathbf{x})$ acts just like multiplication by $\alpha_{E/L}(\mathbf{x})$ and fixes $E[m]$, without the extra $\text{Nm}_{L/K}(\mathbf{x})_\infty$ factor, so that α will actually be 1.

Let B_m be the kernel of the Artin map $\mathbb{I}_L \rightarrow G(L(E[m])/L)$ (abelian by Lemma 17.5.3), so that it induces an isomorphism

$$\phi_{L(E[m])/L} : \mathbb{I}_L/B_m \xrightarrow{\cong} G(L(E[m])/L). \tag{17.16}$$

We show that

$$U_m := B_m \cap L^\times (\text{Nm}_{L/K}^{-1} \mathbb{U}_K(1, m)) \subseteq \ker \chi_{E/L}.$$

This is of finite index in \mathbb{I}_L since B_m is open of finite index in \mathbb{I}_L and $K^\times \mathbb{U}_K(1, m)$ is open of finite index in \mathbb{I}_K .

Fixing an analytic isomorphism $f : \mathbb{C}/\mathfrak{a} \xrightarrow{\cong} E(\mathbb{C})$, we get that for any $t \in m^{-1}\mathfrak{a}/\mathfrak{a}$ and any $\mathbf{x} \in U_m$, $f(t) \in E[m]$ so

$$\begin{aligned} f(t) &= f(t)^{\phi_L(\mathbf{x})} && \text{by (17.16) and } \mathbf{x} \in B_m \\ &= f(\alpha \text{Nm}_{L/K}(\mathbf{x})^{-1}t) && \text{by the Main Theorem 17.6.2} \\ &= f(\alpha t) && t \in m^{-1}\mathfrak{a}/\mathfrak{a} \text{ and } \text{Nm}_{L/K}(\mathbf{x})_{\mathfrak{p}} \equiv 1 \pmod{m\mathcal{O}_{K_{\mathfrak{p}}}} \text{ for all } \mathfrak{p}. \end{aligned}$$

Thus multiplication by α fixes $m^{-1}\mathfrak{a}/\mathfrak{a}$, i.e. $\alpha \equiv 1 \pmod{m\mathcal{O}_K}$. Note $\text{Nm}_{L/K}(\mathbf{x})^{-1} \in \mathbb{U}_K(1, m)$, so

$$(\alpha) = (\mathbf{y}) = (\text{Nm}_{L/K}(\mathbf{x})) = \mathcal{O}_K$$

and α is a unit. Together with $\alpha \equiv 1 \pmod{m\mathcal{O}_K}$, we get $\alpha = 1$.⁸

Part 3: The relationship between ramification and bad reduction hinges on the Néron-Ogg-Shafarevich Criterion. See [Sil94, pg. 169-170]. □

Note that if $\chi_{E/L}$ is unramified at \mathfrak{P} , then $\chi_{E/L}(i_{\mathfrak{P}}(U_{\mathfrak{P}})) = 1$, so it makes sense to talk about $\chi_{E/L}(\mathfrak{P})$ (defined as $\chi_{E/L}(i_{\mathfrak{P}}(\pi))$ for any uniformizer π).

Proposition 17.6.5: Let E/L be an elliptic curve with CM by \mathcal{O}_K , with $K \subseteq L$. Let \mathfrak{P} be a prime of L of good reduction, let \tilde{E} be the reduction of E modulo \mathfrak{P} . Let $\phi_{\mathfrak{P}}$ be the Frobenius on \tilde{E} . Then the following commutes.

$$\begin{array}{ccc} E & \xrightarrow{[\chi_{E/L}(\mathfrak{P})]} & E \\ \downarrow & & \downarrow \\ \tilde{E} & \xrightarrow{\phi_{\mathfrak{P}}} & \tilde{E} \end{array}$$

⁸Any number in the form $m\tau + 1$, $\tau \in \mathcal{O}_K$ with norm 1 has norm at least $(\text{Nm}_{K/\mathbb{Q}}(m) - 1)^2 - 1$, by the triangle inequality. In order for it to have norm 1, $\tau = 0$.

Proof. Let π be a uniformizer of $L_{\mathfrak{P}}$, and let $\varpi = i_{\mathfrak{P}}(\pi)$. Note that $\varpi_{\infty} = 1$. Hence $\text{Nm}_{L/K}(\varpi)_{\infty} = 1$, giving

$$\chi_{E/L}(\mathfrak{P}) = \chi_{E/L}(\varpi) = \alpha_{E/L}(\varpi).$$

If m is an integer such that $\mathfrak{P} \nmid m$, then $\text{Nm}_{L/K}(\varpi)$ fixes $m^{-1}\mathfrak{a}/\mathfrak{a}$ (since it is 1 at all \mathfrak{Q} with $\mathfrak{Q} \mid m$). Then

$$\begin{aligned} f(t)^{\phi_L(\varpi)} &= f([\alpha_{E/L}(\varpi)] \text{Nm}_{L/K}(\varpi)^{-1}t) && \text{definition of } \alpha_{E/L} \\ &= f([\chi_{E/L}(\mathfrak{P})] \text{Nm}_{L/K}(\varpi)^{-1}t) \\ &= [\chi_{E/L}(\mathfrak{P})]f(\text{Nm}_{L/K}(\varpi)^{-1}t) && f \text{ preserves the action of } \mathcal{O}_K \\ &= [\chi_{E/L}(\mathfrak{P})]f(t) && \text{Nm}_{L/K}(\varpi) \text{ fixes } m^{-1}\mathfrak{a}/\mathfrak{a}. \end{aligned}$$

Modulo \mathfrak{P} , $\phi_L(\varpi)$ is just the q th power Frobenius map, so we get

$$\phi_{\mathfrak{P}}|_{\tilde{E}[m]} = \overline{[\chi_{E/L}(\mathfrak{P})]}|_{E[m]}.$$

Since an isogeny is determined by its action on $E[m]$ for $m \rightarrow \infty$ (the kernel of a nonzero isogeny is finite), we get that this is true for E , not just $E[m]$, as needed. \square

To study the Galois representation $G(\overline{K}/H_K) \rightarrow \text{Aut } E_{\text{tors}}$ of E , we reduce modulo a prime \mathfrak{P} of L , and show that on this reduced curve, the q th power Frobenius acts exactly as multiplication by the Grössencharacter. In particular, the q th power Frobenius is represented by multiplication by $\chi_{E/L}(\mathfrak{P})$ when we think of E_{tors} as K/\mathfrak{a} . Thinking of E_{tors} as a 2-dimensional space \mathbb{Q}^2 , this says exactly that the eigenvalues of the Frobenius acting on E_{tors} is exactly $\chi_{E/L}(\mathfrak{P})$ and $\overline{\chi_{E/L}(\mathfrak{P})}$. Typically we just restrict our attention to ℓ -power torsion points for some ℓ .

7 L -series of CM elliptic curve

7.1 Defining the L -function

We define the L -series of an elliptic curve as the L -series of the corresponding Galois representation.

Definition 17.7.1: Let E be an elliptic curve defined over K , and ρ_{ℓ} the associated Galois representation $G(\overline{K}/K) \rightarrow \text{Aut } V_{\ell}E \cong \text{GL}_2(\mathbb{Q}_{\ell})$.

Define the **local L -factor** of E at a prime \mathfrak{p} of K as follows. Choose ℓ such that $\mathfrak{p} \nmid \ell$, and let

$$L_{\mathfrak{p}}(E, s) := L_{\mathfrak{p}}(\rho_{\ell}, s) = \det(1 - q^{-s} \text{Frob}(\mathfrak{p})|(V_{\ell}E)^{I_{\mathfrak{p}}})^{-1},$$

where $q = \mathfrak{N}\mathfrak{p}$ and $I_{\mathfrak{p}}$ is the inertia subgroup of $G(\overline{K}/K)$. (Choose an embedding $\mathbb{Q}_{\ell} \hookrightarrow \mathbb{C}$.) The L -series of E is the product of local factors

$$L_{\mathfrak{p}}(E/K, s) := \prod_{\mathfrak{p}} L_{\mathfrak{p}}(E, s).$$

Remark 17.7.2: This is (almost) the same as saying: fix a prime ℓ and let $L(E/K, s) := L(\rho_\ell, s)$. The only difference is that we run into trouble with the local factor $L_{\mathfrak{p}}(\rho_\ell, s)$ on the right hand side, so we have to choose a different ℓ' and let this local factor be $L_{\mathfrak{p}}(\rho_{\ell'}, s)$ instead.

The following is an equivalent definition (that is more concrete).

Definition 17.7.3: Let N be the conductor⁹ of the elliptic curve E . Define the local L -factor by

$$L_{\mathfrak{p}}(E, s) = 1 - a_q q^{-s} + \chi(q) q q^{-2s}, \quad a_q = q + 1 - |E(\mathbb{F}_q)|, \quad \chi(q) = \begin{cases} 1, & m \perp N \\ 0, & \text{else} \end{cases}$$

where $q = \mathfrak{N}\mathfrak{p}$. Thus

$$L_v(E, s) = \begin{cases} 1 - a_q q^{-s} + q q^{-2s}, & \text{good reduction} \\ 1 - q^{-s}, & \text{split multiplicative reduction} \\ 1 + q^{-s}, & \text{non-split multiplicative reduction} \\ 1, & \text{additive reduction.} \end{cases}$$

Note that a_q , the “trace of Frobenius,” is related to the number of points of E over \mathbb{F}_q . Hence the L -function contains information about the number of points of E over each \mathbb{F}_q .

Showing that these two definitions are equivalent requires us to show that $(V_\ell E)^{I_{\mathfrak{p}}}$ is 2, 1, or 0-dimensional when E has good, multiplicative, and additive reduction, respectively. The general idea is that the action of $I_{\mathfrak{p}}$ on $V_\ell E$ contains exactly the information lost by looking at the reduced elliptic curve, since $I_{\mathfrak{p}}$ is exactly the kernel of $D_{\mathfrak{p}}(\overline{K}/K) \rightarrow G(\overline{k}/k)$, so nontrivial action of $I_{\mathfrak{p}}$ corresponds to bad reduction.

In the CM case, we cannot have multiplicative reduction, so the L -series is particularly simple. We will show that the two definitions are equivalent in this case.

Theorem 17.7.4. *Let E/K be a CM elliptic curve. Then E cannot have multiplicative reduction at any prime.*

Proof. An elliptic curve E has potential good reduction iff its j -invariant is integral [Sil86, VII.5.5]. CM have integral j -invariants, so have potential good reduction, i.e. have good or multiplicative reduction. \square

Proof that Definitions 17.7.1 and 17.7.3 are equivalent in the CM case. Suppose E has CM by an order \mathcal{O} in K , and E is defined over L . By Néron-Ogg-Shafarevich, $I_{\mathfrak{p}}$ acts trivially on $V_\ell E$ iff E has good reduction at \mathfrak{p} . Let $q = \mathfrak{N}\mathfrak{p}$.

In the case of good reduction we need to show $\det(1 - q^{-s} \text{Frob}(\mathfrak{p})|V_\ell E) = 1 - a_q q^{-s} + q q^{-2s}$. Every endomorphism ϕ on E satisfies $\phi^2 - \text{tr}(\phi)\phi + \text{deg}(\phi) = 0$, where $\text{tr}(\phi) = 1 + \text{deg}(\phi) -$

⁹ N is divisible by exactly the primes of bad reduction

$\deg(1 - \phi)$. Since $\text{Frob}(\mathfrak{p})$ acts as the Frobenius morphism $\phi_{\mathfrak{p}}$, its characteristic polynomial is

$$\det(\lambda - \text{Frob}(\mathfrak{p})) = \lambda^2 - \text{tr}(\phi_{\mathfrak{p}})\lambda + \deg(\phi_{\mathfrak{p}}).$$

But

$$\begin{aligned} \deg(\phi_{\mathfrak{p}}) &= q \\ \text{tr}(\phi_{\mathfrak{p}}) &= 1 + \deg(\phi_{\mathfrak{p}}) - \deg(1 - \phi_{\mathfrak{p}}) \\ &= q + 1 - \ker(1 - \phi_{\mathfrak{p}}) \\ &= q + 1 - |E(\mathbb{F}_q)|. \end{aligned}$$

(This part of the proof doesn't use the fact that E has CM.)

Since E has no multiplicative reduction by Theorem 17.7.4, it remains to prove that $W := (V_{\ell}E)^{I_{\mathfrak{p}}} = 0$ when E has multiplicative reduction. We know by Néron-Ogg-Shafarevich that $\dim(W) \leq 1$. But because E is CM, $V_{\ell}E \cong (\varprojlim^n \ell^{-n}\mathfrak{a}/\mathfrak{a}) \otimes \mathbb{Q}$ has the structure of a $\mathcal{O}_K \otimes \mathbb{Q}_{\ell}$ -vector space. If $a \in W$, then for any $\alpha \in \overleftarrow{K}$, $\alpha a \in W$ because $[\alpha]$ commutes with the Galois action. Hence W is not just a \mathbb{Q}_{ℓ} -subspace of V , but also a $\mathcal{O}_K \otimes \mathbb{Q}_{\ell}$ -subspace. Hence its dimension over \mathbb{Q}_{ℓ} is even, and must be 0. \square

7.2 Analytic continuation

Theorem 17.7.5 (Deuring). *Let E/L be an elliptic curve with CM by \mathcal{O}_K with $K \subseteq L$. Then*

$$L(E/L, s) = L(s, \psi_{E/L})L(s, \overline{\psi_{E/L}}).$$

Corollary 17.7.6 (Analytic continuation of L -function for CM elliptic curves). *Let E/L be an elliptic curve with CM by \mathcal{O}_K . Then L admits an analytic continuation to \mathbb{C} and satisfies a functional equation relating its values at s and $2 - s$.*

This theorem for general elliptic curves is very deep (it follows from the Modularity Theorem and the analytic properties of L -functions associated to modular forms).

Proof of Theorem 17.7.5. By Theorem 17.7.4, E has no multiplicative reduction. Let \mathfrak{P} be a prime, and consider 2 cases.

1. E has good reduction at \mathfrak{P} . Choose any ℓ not dividing \mathfrak{P} . The characteristic polynomial of the action of $\phi_{\mathfrak{P}}$ on $V_{\ell}E$ is $\det(\lambda - \phi_{\mathfrak{P}}|V_{\ell}E)$. However, if we make the identification $E_{\text{tors}} \cong K/\mathfrak{a}$, we have

$$V_{\ell}E = \varprojlim \ell^{-n}\mathfrak{a}/\mathfrak{a},$$

and we know that $\phi_{\mathfrak{P}}$ acts on $E_{\text{tors}} \cong K/\mathfrak{a}$ as multiplication by $\chi_{E/L}(\mathfrak{P})$. Therefore, the eigenvalues of the action of $\phi_{\mathfrak{P}}$ on $V_{\ell}E$ are just $\chi_{E/L}(\mathfrak{P})$ and $\overline{\chi_{E/L}(\mathfrak{P})}$, and

$$\det(\lambda - \phi_{\mathfrak{P}}|V_{\ell}E) = (\lambda - \chi_{E/L}(\mathfrak{P}))(\lambda - \overline{\chi_{E/L}(\mathfrak{P})}).$$

Taking $\lambda = p^s$ and dividing by p^{2s} gives

$$L_{\mathfrak{P}}(E/L, s) = \det(1 - p^{-s}\phi_{\mathfrak{P}}|V_{\ell}E) = L_{\mathfrak{P}}(s, \chi_{E/L})L(s, \overline{\chi_{E/L}}).$$

2. E has bad reduction at \mathfrak{P} . Then $\chi_{E/L}(\mathfrak{P}) = 0$ by definition, and $L_{\mathfrak{P}}(E/L, s) = 1 = (1 - \chi_{E/L}(\mathfrak{P}))(1 - \bar{\chi}_{E/L}(\mathfrak{P})) = L_{\mathfrak{P}}(s, \chi_{E/L})L(s, \bar{\chi}_{E/L})$.

Multiplying together all the local factors gives the result. □

Proof of Corollary 17.7.6. The L -functions of Grössencharacters have analytic continuation (Theorem 16.16.7.8, which works for Grössencharacters as well). Thus the result follows directly from Theorem 17.7.5. □

Thus we have carried out the program in Section 16.7 for CM elliptic curves, to get the correspondences.

$$(\text{CM Elliptic curves}) \rightarrow (\text{Galois representation}) \rightarrow (2 \text{ Grössencharacters})$$

Remember Grössencharacters are 1-dimensional automorphic representations. If we wanted a modular form, we can use the technique of *automorphic induction* to construct a modular form from 2 Grössencharacters.

Part III
Analytic Number Theory

Chapter 18

Elementary estimates for primes

1 Chebyshev's Theorem

Today we prove some asymptotic results about the distribution of prime numbers. Specifically, we derive estimates for the *prime-counting functions*

$$\begin{aligned}\vartheta(x) &= \sum_{p \leq x} \ln(p) \\ \psi(x) &= \sum_{p^k \leq x} \ln(p) \\ \pi(x) &= \sum_{p \leq x} 1\end{aligned}$$

Note that we will always use p to denote a prime.

Lacking the tools of complex analysis, it is difficult to find the exact asymptotic formulas; however, our elementary methods suffice to determine the asymptotics up to a constant multiple. Our main result is Chebyshev's Theorem:

Theorem 18.1.1. [?, Theorem 6.3] *There exist positive constants c_1 and c_2 such that*

$$c_1x \leq \vartheta(x) \leq \psi(x) \leq \pi(x) \ln(x) \leq c_2x. \quad (18.1)$$

for all $x \geq 2$. Moreover,

$$\liminf_{x \rightarrow \infty} \frac{\vartheta(x)}{x} = \liminf_{x \rightarrow \infty} \frac{\psi(x)}{x} = \liminf_{x \rightarrow \infty} \frac{\pi(x) \ln(x)}{x} \geq \ln(2) \quad (18.2)$$

$$\limsup_{x \rightarrow \infty} \frac{\vartheta(x)}{x} = \limsup_{x \rightarrow \infty} \frac{\psi(x)}{x} = \limsup_{x \rightarrow \infty} \frac{\pi(x) \ln(x)}{x} \leq 2 \ln(2) \quad (18.3)$$

We will prove this in three steps.

1.1 Comparing the three functions

Since all terms in the sum defining $\vartheta(x)$ are included in the sum defining $\psi(x)$, $\vartheta(x) \leq \psi(x)$. For a given p there are $\left\lfloor \frac{\ln(x)}{\ln(p)} \right\rfloor$ choices for k so that $p^k \leq x$, so

$$\psi(x) = \sum_{p^k \leq x} \ln(p) = \sum_{p \leq x} \left\lfloor \frac{\ln(x)}{\ln(p)} \right\rfloor \ln(p) \leq \sum_{p \leq x} \ln(x) = \pi(x) \ln(x).$$

This shows the middle two inequalities in (18.1).

Given $\vartheta(x) \leq \psi(x) \leq \pi(x) \ln(x)$, to show that the three quantities in (18.2) and (18.3) are equal it suffices to show that

$$\liminf_{x \rightarrow \infty} \frac{\vartheta(x)}{x} \geq \liminf_{x \rightarrow \infty} \frac{\pi(x) \ln(x)}{x}, \quad \limsup_{x \rightarrow \infty} \frac{\vartheta(x)}{x} \geq \limsup_{x \rightarrow \infty} \frac{\pi(x) \ln(x)}{x} \quad (18.4)$$

To compare $\vartheta(x) = \sum_{p \leq x} \ln(p)$ and $\pi(x) \ln(x) = \sum_{p \leq x} \ln(x)$, note that for p “close” to x , we have $\ln(p)$ “close” to $\ln(x)$ and relatively large, while the terms for small p will not contribute much to either sum. Thus we can just consider the terms with $p > x^{1-\delta}$, where $\delta \in (0, 1)$.

$$\begin{aligned} \vartheta(x) &\geq \sum_{x^{1-\delta} < p \leq x} \ln(p) \\ &\geq \sum_{x^{1-\delta} < p \leq x} \ln(x^{1-\delta}) \\ &= \ln(x^{1-\delta})(\pi(x) - \pi(x^{1-\delta})) \\ &= (1 - \delta) \ln(x)(\pi(x) - \pi(x^{1-\delta})) \\ &\geq (1 - \delta) \ln(x)(\pi(x) - x^{1-\delta}) \end{aligned}$$

Hence

$$\frac{\vartheta(x)}{x} \geq \frac{(1 - \delta)\pi(x) \ln(x)}{x} - \frac{(1 - \delta) \ln(x)}{x^\delta}.$$

Letting $\delta \rightarrow 0$ gives (18.4).

1.2 Upper Bound

We show that $\vartheta(x) \leq 2x \ln(x)$. Instead of thinking about bounding $\vartheta(x)$, it is easier to think about bounding $e^{\vartheta(x)} = \prod_{p \leq x} p$.

Lemma 18.1.2. [AD08, 3.?] For any $x \in \mathbb{N}$,

$$\prod_{p \leq x} p \leq 4^{x-1} \quad (18.5)$$

Proof. Use strong induction on x . For $x = 1, 2$ the statement holds. The induction step from odd $x > 1$ to $x + 1$ is obvious, since $x + 1$ is not a prime.

Consider the induction step from even x to $x + 1$. Let $x = 2n$. The key idea is that there cannot be “too many” primes between $n + 2$ and $2n + 1$, because...

1. These primes all divide $\binom{2n+1}{n} = \frac{(2n+1)!}{n!(n+1)!}$.
2. $\binom{2n+1}{n}$ can easily be bounded from above:

$$\binom{2n+1}{n} = \frac{1}{2} \left(\binom{2n+1}{n} + \binom{2n+1}{n+1} \right) \leq \frac{1}{2} \sum_{i=0}^n \binom{2n+1}{i} = 4^n.$$

Then

$$\prod_{p \leq x+1} p = \prod_{p \leq n+1} p \prod_{n+2 \leq p \leq 2n+1} p \leq 4^n \cdot \binom{2n+1}{n} \leq 4^{2n}.$$

□

Taking the logarithm of both sides of (18.5) gives $\vartheta(x) \leq (x-1) \ln(4) \leq 2x \ln(x)$.

1.3 Lower Bound

We show that $\liminf_{x \rightarrow \infty} \frac{\pi(x) \ln(x)}{x} \geq \ln(2)$. First consider when x is even, say equal to $2n$. Like in Section 3, we consider a binomial coefficient, this time $\binom{2n}{n}$. We show that each prime cannot appear as a factor in $\binom{2n}{n}$ “too many” times, so it can be bounded above by $(2n)^{\pi(2n)}$. We can easily bound $\binom{2n}{n}$ below:

$$\binom{2n}{n} \geq \frac{2^{2n}}{2n}$$

since it is the largest among $2, \binom{2n}{1}, \dots, \binom{2n}{2n-1}$. Putting these two bounds together will give the desired bound for $\pi(2n)$.

We need the following to count the highest prime powers dividing $\binom{2n}{n}$:

Lemma 18.1.3. [*AD08, Lemma 6.3*] For every positive integer n ,

$$v_p(n!) = \sum_{k=1}^{\lfloor \frac{\ln(n)}{\ln(p)} \rfloor} \left\lfloor \frac{n}{p^k} \right\rfloor,$$

where $v_p(m)$ denotes the largest integer i such that $p^i | m$.

Proof. There are $\lfloor \frac{n}{p^k} \rfloor$ multiples of p^k less than or equal to n . In the sum $\sum_{k \geq 1} \lfloor \frac{n}{p^k} \rfloor$, each multiple of p^k less than n is counted k times, once each as a multiple of p, p^2, \dots, p^k . □

From Lemma 18.1.3, we get

$$v_p \left(\binom{2n}{n} \right) = v_p \left(\frac{(2n)!}{n!^2} \right) = v_p((2n)!) - 2v_p(n!) = \sum_{k=1}^{\lfloor \frac{\ln(2n)}{\ln(p)} \rfloor} \left\lfloor \frac{2n}{p^k} \right\rfloor - 2 \left\lfloor \frac{n}{p^k} \right\rfloor$$

Since each term of the sum is at most 1,

$$v_p \left(\binom{2n}{n} \right) \leq \left\lfloor \frac{\ln(2n)}{\ln(p)} \right\rfloor \leq \frac{\ln(2n)}{\ln(p)}.$$

Thus

$$\frac{2^{2n}}{2n} \leq \binom{2n}{n} = \prod_{p \leq 2n} p^{v_p \left(\binom{2n}{n} \right)} \leq (2n)^{\pi(2n)}.$$

Taking logs and remembering $x = 2n$ gives $x \ln(2) - \ln(x) \leq \pi(x) \ln(x)$, which gives the desired bound. For odd x , the value of $\frac{\pi(x) \ln(x)}{x}$ can be compared to the value for $x - 1$.

Finally, (18.2) and (18.3), and the fact that all the prime-counting functions are positive for $x \geq 2$, show the existence of c_1 and c_2 in (18.1). This finishes the proof of Theorem 18.1.1.

1.4 The n th prime

We found an estimate for the number of primes less than or equal to a given number; we can use this bound to find an estimate for the n th prime number.

Theorem 18.1.4. *Let p_n denote the n th prime number. Then there exist constants c_3, c_4 such that*

$$c_3 n \ln(n) \leq p_n \leq c_4 n \ln(n)$$

for all $n \geq 2$.

Proof. From Theorem 18.1.1,

$$\frac{c_1 p_n}{\ln(p_n)} \leq \pi(p_n) = n \leq \frac{c_2 p_n}{\ln(p_n)}, \quad (18.6)$$

so

$$\frac{n \ln(p_n)}{c_2} \leq p_n \leq \frac{n \ln(p_n)}{c_1}.$$

The LHS is at least $c_3 n \ln(n)$ by the trivial bound $n \leq p_n$. On the RHS, use the LHS of (18.6) again to get $\ln(p_n) \leq \ln \left(\frac{n \ln(p_n)}{c_1} \right)$, giving $\ln(p_n) \leq c \ln(n)$ for some c . \square

Chapter 19

Crash course in complex analysis

Complex analysis is calculus on the complex numbers. The main functions of study are complex differentiable functions.

Reference books: Lang or Ahlfors

1 Holomorphic functions

Definition 19.1.1: Let $U \subseteq \mathbb{C}$ be an open set and $f : U \rightarrow \mathbb{C}$ be a function. The **derivative** of f is

$$f'(z) := \lim_{\Delta z \rightarrow 0} \frac{f(z + \Delta z) - f(z)}{\Delta z}$$

if it exists. f is **holomorphic** if its derivative exists at every point of U . f is **meromorphic** if it is defined and holomorphic on U except at a discrete set of points.

Write $f(x + iy) = u(x, y) + iv(x, y)$. Note that f being differentiable is a much *stronger* condition than being simply u and v being differentiable, because the limit of f as $\Delta z \rightarrow 0$ along the real and complex directions must be equal:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial x} = \frac{1}{i} \left(\frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right).$$

Thus we get the Cauchy-Riemann criteria: If f is differentiable as a function of (x, y) , then f is holomorphic iff

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Another way to think about complex differentiability is that holomorphic maps preserve angles (i.e. are *conformal*); we have

$$f(z + re^{i\theta}) - f(z) \approx re^{i\theta} f'(z).$$

Because complex differentiability is such a strong property, holomorphic functions have many nice properties. Hence it is often useful to take functions defined on the reals and

extend them as far as possible on \mathbb{C} . Some of the good properties are the following (to be explained in the rest of the chapter); note they are not necessarily true for real differentiable functions!

- A function is holomorphic iff it is analytic (has a power series expansion).
- A sequence of holomorphic functions with good convergence properties converges to a holomorphic function.
- A bounded entire function is constant.
- If two holomorphic functions agree on a set containing a limit point, then they are equal. Thus analytic continuations are unique.
- Bounds on a function give bounds on the derivative. Hence we can “differentiate” asymptotic formulas.
- We can expand holomorphic functions into products or sums depending on their poles and zeros—in much the same way that rational functions can be expanded into partial fractions or factored.

2 Complex integration

We now give two definitions of the integral.

Definition 19.2.1: A **path** is a continuous function $\gamma : [a, b] \rightarrow \mathbb{C}$. It is called a **loop** if $\gamma(a) = \gamma(b)$. Let f be a holomorphic function on U and γ be a path in U .

1. If γ is differentiable (except possibly at a finite number of points), define

$$\int_{\gamma} f(z) dz = \int_a^b f(\gamma(t))\gamma'(t) dt.$$

2. Define an (indefinite) **integral** of f on a set V to be a function F on V such that $F'(z) = f(z)$. Given holomorphic f , choose points t_0, \dots, t_n such that there exist open sets $U_j \supseteq f(\gamma([t_{j-1}, t_j]))$ such that f has an integral F_j on U_j . Define

$$\int_{\gamma} f(z) dz = \sum_{k=1}^n [F_j(\gamma(t_j)) - F_j(\gamma(t_{j-1}))].$$

Note that unlike in the real case, indefinite integrals may not exist globally, for example, $\ln t$ is locally an integral for $\frac{1}{t}$ but cannot be extended holomorphically to $\mathbb{C} \setminus \{0\}$. We need to establish the well-definedness of the second definition.

Theorem 19.2.2 (Cauchy's Theorem, version 1). *Let f be holomorphic on a closed rectangle R , with boundary ∂R . Then (using the first definition),*

$$\int_{\partial R} f = 0.$$

From this one can show that integrals exist locally by defining

$$F(z) = \int_{z_0}^z f(s) ds$$

where the integral is along horizontal and vertical lines; moreover one gets well-definedness in the second definition.

We can now define the logarithm of a function.

Definition 19.2.3: Let f be a holomorphic function on a simply connected set U (see Definition 19.3.1), with $f(z) \neq 0$ on U . Choose $z_0 \in U$ and a_0 such that $e^{a_0} = z_0$.

$$(\ln f)(z) = \int_z^{z_0} \frac{f'}{f}(z) dz.$$

Note different definitions of the logarithm will differ by integer multiples of $2\pi i$, and $e^{(\ln f)(z)} = f(z)$. The motivation comes from the fact that one would expect the derivative of $\ln f(z)$ to be $\frac{f'}{f}(z)$. We write $(\ln f)(z)$ to emphasize that this is *not* simply a composite of functions: We could have $f(z_1) = f(z_2)$ but $(\ln f)(z_1) \neq (\ln f)(z_2)$.¹

We seek a generalization of Theorem 19.2.2 to meromorphic functions and arbitrary paths.

3 Cauchy's Theorem

Definition 19.3.1: Two paths γ and $\eta : [a, b] \rightarrow \mathbb{C}$ are **homotopic** if there exists a continuous map

$$\gamma_s(t) : [0, 1] \times [a, b] \rightarrow \mathbb{C}$$

such that $\gamma_0(t) = \gamma(t)$ and $\gamma_1(t) = \eta(t)$.

A subset of \mathbb{C} is simply connected if it is pathwise connected and every loop in \mathbb{C} is homotopic to a point.

Theorem 19.3.2. *Let U be a simply connected open set containing z_0 . Every path γ around z_0 in $U \setminus \{z_0\}$ is homotopic to a circle going around z_0 n times for some $n \in \mathbb{Z}$. This n can be calculated by*

$$n = W(\gamma, z_0) := \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z - z_0} dz$$

and is called the **winding number**.

¹Consider, for example, the case where $f(z) = z^2$ on $\mathbb{C} \setminus \mathbb{R}_{\leq 0}$, and $z_1 = i$, $z_2 = -i$.

Theorem 19.3.3 (Global Cauchy's formula). *Let U be a simply connected open set and $f : U \rightarrow \mathbb{C}$ be holomorphic. Suppose γ is a loop in U . Then*

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z - z_0} dz = W(\gamma, z_0) f(z_0).$$

4 Power series and Laurent series

As complex differentiability is a much stronger condition than differentiability for real functions, holomorphic functions enjoy nicer properties. The most important one is the following.

Definition 19.4.1: A function $f : U \rightarrow \mathbb{C}$ is **analytic** at z_0 if it can be written as a power series in a neighborhood around z_0 :

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n.$$

If f is given by its power series representation then we must have $a_n = \frac{f^{(n)}(z_0)}{n!}$.

Theorem 19.4.2. *A function $f : U \rightarrow \mathbb{C}$ is analytic iff and only iff it is holomorphic.*

Note this is not true for real functions: for example, $e^{-\frac{1}{x^2}}$ has Taylor expansion equal to 0 at 0, but is not the zero function. This kind of irregularity does not happen for holomorphic functions.

Corollary 19.4.3. *A holomorphic function has infinitely many derivatives.*

The following theorem says that for holomorphic functions, the radius of convergence is “as large as it could possibly be.”

Theorem 19.4.4. *Suppose f is holomorphic on a disc $N_r(z_0)$ of radius r around z_0 . Then the Taylor series around z_0 converges absolutely to f on $N_r(z_0)$.*

Proof. Estimate coefficients using Cauchy's theorem. Complex Analysis, Lang III.7.3. \square

We can generalize power series to allow terms with negative exponents.

Theorem 19.4.5. *Suppose f is defined on an annulus $A = \{z : r < |z - z_0| < R\}$. Let C be the circle of radius $r' \in (r, R)$ around z_0 . Then f has a Laurent expansion on A :*

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n, \quad a_n = \frac{1}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^{n+1}} dz.$$

If f is defined on $\{z : |z - z_0| < R\}$ then

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^{n+1}} dz.$$

The coefficient a_{-1} is called the **residue** of f at z_0 :

$$\operatorname{Res}_{z_0}(f) = a_{-1}.$$

The following theorem controls the size of the derivatives of a complex analytic function by its values of the function in a circle. Note that in the real analytic case we can't make such a statement!

Corollary 19.4.6. *Suppose f is defined on $\{z : |z - z_0| < R\}$, and let C be a circle of radius $r < R$ around z_0 . Then*

$$|f^{(n)}(z)| \leq \frac{n!}{r^n} \max_{z \in C} |f(z)|$$

and the n th coefficient in the power series expansion satisfies

$$a_n \leq \frac{1}{r^n} \max_{z \in C} |f(z)|.$$

Proof. Simply note that in the integral $\int_C \frac{f(z)}{(z-z_0)^{n+1}} dz$, the denominator has constant absolute value r^{n+1} , the numerator is bounded by $\max_{z \in C} |f(z)|$, and the arc length is $2\pi r$. \square

Corollary 19.4.7 (Liouville). *A bounded entire function is constant.*

Proof. We can take $r \rightarrow \infty$ in the inequality for $n = 1$ to find that $f'(z) = 0$ everywhere. \square

4.1 Cauchy's residue formula

Using residues, we can state the most comprehensive form of Cauchy's formula:

Theorem 19.4.8 (Residue formula). *Suppose f is meromorphic on simply connected open U , and γ is a loop in U . Then*

$$\int_{\gamma} f(s) ds = 2\pi i \sum_{z \text{ pole of } f} W(\gamma, z) \operatorname{Res}_z(f).$$

One useful application of this is counting zeros and poles of a function f .

Definition 19.4.9: Define the **order** of f at z_0 to be the least integer so that the Laurent expansion of f at z_0 has $a_m \neq 0$:

$$\operatorname{ord}_f(z_0) = m.$$

Note that $\operatorname{ord}_f(z_0) > 0$ signals a zero and $\operatorname{ord}_f(z_0) < 0$ signals a pole.

Corollary 19.4.10. *Suppose f is meromorphic on simply connected open U , and γ is a loop in U . Then*

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'}{f}(s) ds = \sum_{\rho} W(\gamma, \rho) \operatorname{ord}_f(\rho).$$

Proof. If f has Laurent expansion $a_m(z - z_0)^m + \dots$ at z_0 then $\frac{f'}{f}$ has Laurent expansion

$$\frac{ma_m(z - z_0)^{m-1} + \dots}{a_m(z - z_0)^m + \dots} = m(z - z_0)^{-1} + \dots$$

\square

5 Convergence

Unlike in the real case, holomorphic functions behave nicely under infinite sums and pointwise convergence. This is because by Cauchy's theorem we can write f as an integral, and integrals preserve convergence.

Theorem 19.5.1 (Holomorphic functions converge to holomorphic functions). *Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of holomorphic functions on U .*

1. *Suppose $f_n \rightarrow f$ uniformly on compact subsets of U . Then f is holomorphic.*
2. *Suppose $\sum_{n=1}^{\infty} f_n = f$ converges absolutely and uniformly on compact subsets of U . Then f is holomorphic.*

6 Series and product developments

We know that locally, we can write a meromorphic function f as a Laurent series $\sum_{n=-\infty}^{\infty} a_n x^n$. There are two other representations that are useful, depending on what information we have about the function f .

1. If we know the *poles* of f , we can write f as a sum of rational functions

$$f(z) = \sum_{n=1}^{\infty} \left[P_n \left(\frac{1}{z - z_n} \right) - Q_n(z) \right] + g(z).$$

2. If f is entire and we know the *zeros* of f , we can write f as an infinite product

$$f(z) = z^m e^{g(z)} \prod_{n=1}^{\infty} \left(1 - \frac{z}{z_n} \right) e^{P_n \left(\frac{z}{z_n} \right)}.$$

(Think of this as “factoring” f , much like a polynomial can be factored as in the fundamental theorem of algebra.) These representations come about from convergence properties of holomorphic functions—so we can be sure the infinite products converge to holomorphic functions—and by Liouville's theorem—if we engineer a function that is close enough to f then it must be equal to f .

Theorem 19.6.1 (Mittag-Leffler). *Let z_n be a sequence with $\lim_{n \rightarrow \infty} |z_n| = \infty$ (or a finite sequence), and P_n polynomials without constant term.*

1. *(Existence) There is a meromorphic function f with poles exactly at z_n , with Laurent expansion $P_n \left(\frac{1}{z - z_n} \right) + \cdots$ at z_n .*
2. *(Uniqueness) All such functions f are in the form*

$$\sum_{n=1}^{\infty} \left(P_n \left(\frac{1}{z - z_n} \right) - Q_n(z) \right) + g(z)$$

where Q_n is a polynomial and $g(z)$ is analytic.

Proof. See Ahlfors [Ahl79, p. 187]. □

Warning: this does not converge for all P_n . Typically we take Q_n to be the first terms of the Laurent expansion of $P_n\left(\frac{1}{z-z_n}\right)$, to ensure cancellation of high-order terms.

Definition 19.6.2: The **order** of an entire function f is the smallest $\alpha \in [0, \infty]$ such that

$$|f(z)| \lesssim_\varepsilon e^{|z|^{\alpha+\varepsilon}}$$

for all $\varepsilon > 0$.

Theorem 19.6.3 (Product development). *Let z_n be a sequence with $\lim_{n \rightarrow \infty} |z_n| = \infty$. If f is entire with order $\alpha < \infty$ with zeros z_1, z_2, \dots (with multiplicity, not including 0), then it has a product formula*

$$f(z) = z^r e^{g(z)} \prod_{n=1}^{\infty} \left(1 - \frac{z}{z_n}\right) e^{\frac{z}{z_n} + \frac{1}{2}\left(\frac{z}{z_n}\right)^2 + \dots + \frac{1}{m}\left(\frac{z}{z_n}\right)^m}, \quad (19.1)$$

where

- $m = \lfloor \alpha \rfloor$,
- r is the order of vanishing of f at 0, and
- g is a polynomial of degree at most a .

The product converges uniformly locally. Moreover,

$$|\{k : z_k < R\}| \lesssim_\varepsilon R^{\alpha+\varepsilon}. \quad (19.2)$$

Conversely, if $a = \lfloor \alpha \rfloor$ and z_k is a sequence satisfying (19.2), then the RHS of (19.1) defines an entire function of order at most α .

Proof. See Ahlfors [Ahl79, p. 195]. □

Hence the order of an entire function gives an asymptotic bound for the number of zeros.²

7 Gamma function

To prove basic properties of the zeta function in the next chapter, we need to know the properties of the gamma function.

Definition 19.7.1: Define the **gamma function** by

$$\Gamma(s) = \int_0^\infty x^s e^{-x} \frac{dx}{x}, \quad \Re s > 0.$$

²A function which grows faster is allowed to have more zeros—much like a polynomial with lots of zeros grows fast simply because it has higher degree.

We will begin by analytically continuing the gamma function and giving its basic properties.

Proposition 19.7.2 (Facts about Γ):

1. $\Gamma(s)$ can be analytically continued to a meromorphic function with poles $-n, n \in \mathbb{N}$, with residue $\frac{(-1)^n}{n!}$.
2. $\Gamma(s) = \lim_{n \rightarrow \infty} \frac{n^n n!}{s(s+1)\cdots(s+n)}$ when $s \notin -\mathbb{N}$.
3. $\frac{1}{\Gamma(s)} = s e^{Cs} \prod_{n=1}^{\infty} \left(1 + \frac{s}{n}\right) e^{-\frac{s}{n}}$.
4. $\Gamma(s+1) = s\Gamma(s)$ so $\Gamma(n+1) = n!, n \in \mathbb{N}_0$.
5. $\Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin \pi s}$.
6. $\Gamma(s)\Gamma\left(s + \frac{1}{m}\right) \cdots \Gamma\left(s + \frac{m-1}{m}\right) = (2\pi)^{\frac{m-1}{2}} m^{\frac{1}{2}-ms} \Gamma(ms)$. In particular, $\Gamma(s)\Gamma\left(s + \frac{1}{2}\right) = \pi^{\frac{1}{2}} 2^{1-2s} \Gamma(2s)$.

From the product development 19.6.3 we get the following.

Theorem 19.7.3 (Product development of Γ). *We have*

$$\Gamma(s) = \frac{e^{-\gamma s}}{s} \prod_{k=1}^{\infty} \frac{e^{\frac{s}{k}}}{1 + \frac{s}{k}}.$$

In the region

$$R_\varepsilon = \mathbb{C} \setminus (\{s : \arg(s) \in [\pi - \varepsilon, \pi + \varepsilon]\} \cup \{0\}),$$

i.e. \mathbb{C} with a wedge containing $\mathbb{R}_{\leq 0}$ deleted, we can define the function $(\ln \Gamma)(s)$. By the product formula, it equals

$$(\ln \Gamma)(s) = -\gamma s - \ln s + \sum_{k=1}^{\infty} \left(\frac{s}{k} - \ln \left(1 + \frac{s}{k}\right) \right).$$

The following asymptotic formulas will be useful.

Theorem 19.7.4 (Stirling's approximation). *Let $P_1(t) = \{t\} - \frac{1}{2}$. For $s \in R_\varepsilon$,*

$$\begin{aligned} (\ln \Gamma)(s) &= \left(s - \frac{1}{2}\right) \ln s - s + \frac{1}{2} \ln(2\pi) - \int_0^\infty \frac{P_1(t)}{z+t} \\ &= \left(s - \frac{1}{2}\right) \ln s - s + \frac{1}{2} \ln(2\pi) + O_\varepsilon(|s|^{-1}) \\ \frac{\Gamma'(s)}{\Gamma(s)} &= \ln s - \frac{1}{2s} + O_\varepsilon(|s|^{-2}) \\ \Gamma(s) &\sim s^{s-\frac{1}{2}} e^{-s} \sqrt{2\pi} \end{aligned}$$

Chapter 20

Dirichlet series

For proofs see [Apo94].

1 Dirichlet series, convergence

Dirichlet series are the “power series of number theory.” As such, we will first need to get acquainted with their analytic properties.

Definition 20.1.1: A **Dirichlet series** is a series of the form

$$F(s) = \sum_{n=1}^{\infty} \frac{f(n)}{n^s}$$

where $f(n)$ is an arithmetical function. Following convention, we let $s = \sigma + it$, with σ, t real.

Let $\{\lambda(n)\}$ be a sequence strictly increasing to ∞ . A **general Dirichlet series** with exponents $\{\lambda(n)\}_{n=1}^{\infty}$ is in the form

$$F(s) = \sum_{n=1}^{\infty} f(n)e^{-s\lambda(n)}.$$

An ordinary Dirichlet series has $\lambda(n) = \ln(n)$.¹

Theorem 20.1.2 (Half-plane of convergence). **Convergence:** *If the series $\sum_{n=1}^{\infty} |f(n)e^{-s\lambda(n)}|$ does not converge or diverge for all n , then there exists a real number σ_c , called the **abscissa of convergence**, such that $\sum_{n=1}^{\infty} f(n)n^{-s}$*

- converges locally uniformly for $\sigma > \sigma_c$, but
- does not converge for $\sigma < \sigma_c$.

¹A further generalization is given by the Laplace-Stieltjes transform, $\int_0^{\infty} e^{-st} d\alpha(t)$, where α is a measure. The “step” part of α gives a Dirichlet while the continuous part gives a Laplace transform.

In fact, if the series diverges for all s with $\sigma < 0$, then

$$\sigma_c = \limsup_{n \rightarrow \infty} \frac{\ln |\sum_{k=1}^n a(k)|}{\lambda(n)}.$$

Absolute convergence: If the series $\sum_{n=1}^{\infty} |e^{-s\lambda(n)}|$ does not converge or diverge for all n , then there exists a real number σ_a , called the **abscissa of absolute convergence**, such that $\sum_{n=1}^{\infty} f(n)n^{-s}$

- converges locally uniformly absolutely for $\sigma > \sigma_a$, but
- does not converge absolutely for $\sigma < \sigma_a$.

In fact, if the series diverges for all s with $\sigma < 0$, then

$$\sigma_a = \limsup_{n \rightarrow \infty} \frac{\ln \sum_{k=1}^n |a(k)|}{\lambda(n)}.$$

In particular, for ordinary Dirichlet series (that diverge when $\sigma < 0$),

$$\sigma_a = \limsup_{n \rightarrow \infty} n^{\sum_{k=1}^n |a(k)|}.$$

2 Basic properties

Proposition 20.2.1 (General facts): Let $F(s) = \sum_{n=1}^{\infty} f(n)n^{-s}$.

1. $\lim_{\sigma \rightarrow \infty} F(\sigma + it) = f(1)$ uniformly
2. (Uniqueness) If $F(s) = G(s)$ are absolutely convergent for $\sigma > \sigma_a$ and are equal for s in an infinite sequence $\{s_k\}$ with $\sigma_k \rightarrow \infty$, then $f(n) = g(n)$.
3. (Non-vanishing in half-plane) Suppose $F(s) \neq 0$ for some s with $\sigma > \sigma_a$. Then there is a half-plane $\sigma > c \geq \sigma_a$ in which $F(s)$ is never 0.

Proposition 20.2.2: (Operations on Dirichlet series) Let $F(s) = \sum_{n=1}^{\infty} f(n)n^{-s}$ and $G(s) = \sum_{n=1}^{\infty} g(n)n^{-s}$. Then

$$F(s)G(s) = \sum_{n=1}^{\infty} \frac{h(n)}{n^s}$$

where

$$h(n) = (f * g)(n) = \sum_{d|n} f(d)g\left(\frac{n}{d}\right).$$

Proof. Formally, by grouping together terms where mn is constant,

$$\begin{aligned} F(s)G(s) &= \sum_{m,n \in \mathbb{N}} \frac{f(m)}{n^s} \frac{g(n)}{n^s} \\ &= \sum_{k=1}^{\infty} \left(\sum_{m,n \in \mathbb{N}, mn=k} f(m)g(n) \right) \frac{1}{k^s}. \end{aligned}$$

Since the sums for F and G converge absolutely, so does the double sum above, and the rearrangement of terms is valid. \square

Theorem 20.2.3 (Euler products). *Let f be a multiplicative arithmetical function such that $\sum_{n=1}^{\infty} f(n)n^{-s}$ converges absolutely. Then when $\Re s > \sigma_a$,*

$$\sum_{n=1}^{\infty} f(n)n^{-s} = \prod_p \left(1 + \frac{f(p)}{p} + \frac{f(p^2)}{p^2} + \cdots \right).$$

If f is completely multiplicative,

$$\sum_{n=1}^{\infty} \frac{f(n)}{n^s} = \prod_p \frac{1}{1 - f(p)p^{-s}}.$$

Proposition 20.2.4 (Derivatives): The derivative is

$$F'(s) = - \sum_{n=1}^{\infty} \frac{f(n) \ln n}{n^s}.$$

Theorem 20.2.5 (Landau). *Suppose $F(s)$ is a holomorphic function that can be represented in $\sigma > c$ by the Dirichlet series*

$$F(s) = \sum_{n=1}^{\infty} f(n)n^{-s}$$

with $f(n) \geq 0$ for all $n \geq n_0$. If $F(s)$ is analytic in some disc of radius r around $s = c$, then $F(s)$ converges in $\sigma > \sigma - \varepsilon$ for some $\varepsilon > 0$.

Hence, $F(s)$ has a singularity at $s = \sigma_c$.

Proof. We reinterpret in terms of power series and apply Theorem 19.4.4.

Take $a = c + \frac{r}{2}$. Since F is analytic at in $N_r(a) \subseteq N_r(c) \cup \{z : \Re z > c\}$, it equals its Taylor expansion there:

$$F(s) = \sum_{k=1}^{\infty} \frac{F^{(k)}(a)}{k!} (s - a)^k.$$

From Proposition 20.2.4, $F^{(k)}(a) = (-1)^k \sum_{n=1}^{\infty} f(n)(\ln n)^k n^{-s}$. Plugging in and noting that

the sum converges absolutely (since $f(n) \geq 0$ for large n), we have, for $s \in N_r(a)$,

$$\begin{aligned} F(s) &= \sum_{k=0}^{\infty} \left[\left(\frac{(-1)^k}{k!} \sum_{n=1}^{\infty} f(n) (\ln n)^k n^{-a} \right) (s-a)^k \right] \\ &= \sum_{n=1}^{\infty} \left[\left(\sum_{k=0}^{\infty} \frac{(s-a)^k (\ln n)^k}{k!} \right) n^{-a} \right] \\ &= \sum_{n=1}^{\infty} f(n) e^{(a-s) \ln n} n^{-a}. \end{aligned}$$

This converges for $c - \varepsilon \in N_r(a)$. But because it has nonnegative real coefficients, this shows $\sigma_c > c - \varepsilon$. \square

Proposition 20.2.6 (Logarithms): Assume $f(1) \neq 0$. if $F(s) \neq 0$ for $\sigma > \sigma_0 \geq \sigma_a$, then for $\sigma > \sigma_0$,

$$\ln F(s) = \ln f(1) + \sum_{n=1}^{\infty} \frac{f' * f^{-1}(n)}{\ln n} n^{-s}.$$

Also talk about log diff of Euler product

3 Dirichlet generating functions

Definition 20.3.1: Let $f : \mathbb{N} \rightarrow \mathbb{C}$ be an arithmetic function. The **Dirichlet generating function** of f is

$$F(s) = \sum_{n=1}^{\infty} \frac{f(n)}{n^s}.$$

To get the generating function of $g(n) = \sum_{d|n} f(n)$, by Proposition 20.2.2, we simply multiply by $\zeta(s)$:

$$F(s)\zeta(s) = \left(\sum_n \frac{f(n)}{n^s} \right) \left(\sum_n \frac{1}{n^s} \right) = \sum_n \left(\sum_{d|n} f(d) \right) \frac{1}{n^s}.$$

Note that the inverse of $\zeta(s)$ is

$$\prod_p (1 - p^{-s}) = \sum_{n=1}^{\infty} \frac{\mu(n)}{n^s}.$$

Hence by matching coefficients of

$$(F(s)\zeta(s)) \frac{1}{\zeta(s)}$$

we get the Mobius inversion formula.

4 Summing coefficients


Lemma 20.4.1. For $y, c, T > 0$,²

$$\begin{aligned} \left| \frac{1}{2\pi i} \int_{c-iT}^{c+iT} y^s \frac{ds}{s} \right| &\leq y^c \min \left(\frac{1}{\pi T |\ln y|}, \frac{1}{2} \right), & 0 < y < 1 \\ \left| \frac{1}{2\pi i} \int_{c-iT}^{c+iT} y^s \frac{ds}{s} - \frac{1}{2} \right| &\leq \frac{y^c}{\pi T}, & y = 1 \\ \left| \frac{1}{2\pi i} \int_{c-iT}^{c+iT} y^s \frac{ds}{s} - 1 \right| &\leq y^c \min \left(\frac{1}{\pi T |\ln y|}, 1 \right), & y > 1 \end{aligned}$$

Proof. First suppose $y < 1$. Take $d > c$. By Cauchy's theorem, since $\frac{y^s}{s}$ is analytic in the region below, we have

$$\int_{c-iT}^{c+iT} y^s \frac{ds}{s} + \int_{c+iT}^{d+iT} y^s \frac{ds}{s} + \int_{d+iT}^{d-iT} y^s \frac{ds}{s} + \int_{d-iT}^{c-iT} y^s \frac{ds}{s} = 0$$

where the path of integrations are those shown in the picture.



Hence,

$$\begin{aligned} \left| \int_{c-iT}^{c+iT} y^s \frac{ds}{s} \right| &= \left| \int_{c+iT}^{d+iT} y^s \frac{ds}{s} + \int_{d-iT}^{c-iT} y^s \frac{ds}{s} + \int_{d+iT}^{d-iT} y^s \frac{ds}{s} \right| \\ &\leq 2 \int_c^d y^\sigma \frac{d\sigma}{T} + \left| \int_{d+iT}^{d-iT} y^s \frac{ds}{s} \right|. \end{aligned}$$

Note that the last integral goes to 0 as $d \rightarrow \infty$, because $|y^s| = |y^d| \rightarrow 0$. Hence, taking $d \rightarrow \infty$ gives

$$\left| \int_{c-iT}^{c+iT} y^s \frac{ds}{s} \right| \leq 2 \int_c^\infty \frac{y^\sigma}{T} d\sigma = -\frac{2y^c}{T \ln y} = \frac{2y^c}{T |\ln y|}.$$

This gives $\left| \frac{1}{2\pi i} \int_{c-iT}^{c+iT} y^s \frac{ds}{s} \right| \leq \frac{y^c}{\pi T} |\ln y|$.

²The integral $\frac{1}{2\pi i} \int_{c-\infty i}^{c+\infty i} f(s) \frac{ds}{s}$ is called the *Mellin transform* of f .

By Cauchy's theorem applied to the smaller segment bounded by $\Re s = c$ and the circle with radius $R = \sqrt{c^2 + T^2}$, we have

$$\begin{aligned} \left| \int_{c-iT}^{c+iT} y^s \frac{ds}{s} \right| &= \left| \int_C y^s \frac{ds}{s} \right| \\ &\leq \pi R \frac{y^c}{R} = \pi y^c, \end{aligned}$$

since $y < 1$ and $\Re s > c$ on the arc. Hence $\left| \frac{1}{2\pi i} \int_{c-iT}^{c+iT} y^s \frac{ds}{s} \right| \leq \frac{y^c}{2}$.

For $y > 1$, take $d < 0$. Note $\frac{y^s}{s}$ is analytic in the region below except for a simple pole at 0 with residue 1 (since $y^s = 1$ when $s = 0$). Hence by Cauchy's Theorem,

$$\int_{c-iT}^{c+iT} y^s \frac{ds}{s} + \int_{c+iT}^{d+iT} y^s \frac{ds}{s} + \int_{d+iT}^{d-iT} y^s \frac{ds}{s} + \int_{d-iT}^{c-iT} y^s \frac{ds}{s} = 2\pi i.$$

Then

$$\begin{aligned} \left| \int_{c-iT}^{c+iT} y^s \frac{ds}{s} - 1 \right| &= \left| \int_{c+iT}^{d+iT} y^s \frac{ds}{s} + \int_{d-iT}^{c-iT} y^s \frac{ds}{s} + \int_{d+iT}^{d-iT} y^s \frac{ds}{s} \right| \\ &\leq 2 \int_d^c y^\sigma \frac{d\sigma}{T} + \left| \int_{d+iT}^{d-iT} y^s \frac{ds}{s} \right|. \end{aligned}$$

The last term goes to 0 as $d \rightarrow -\infty$, so the same argument applies as in the first part to show $\left| \frac{1}{2\pi i} \int_{c-iT}^{c+iT} y^s \frac{ds}{s} - 1 \right| \leq \frac{y^c}{\pi T \ln y}$.

By Cauchy's theorem applied to the larger segment bounded by $\Re s = c$ and the circle with radius $R = \sqrt{c^2 + T^2}$, we have

$$\begin{aligned} \int_{c-iT}^{c+iT} y^s \frac{ds}{s} + \int_C y^s \frac{ds}{s} &= 2\pi i \\ \left| \int_{c-iT}^{c+iT} y^s \frac{ds}{s} - 1 \right| &\leq \left| \int_C y^s \frac{ds}{s} \right| \\ &\leq 2\pi R \frac{y^c}{R} = 2\pi y^c, \end{aligned}$$

since $y > 1$ and $\Re s < c$ on the arc. Hence $\left| \frac{1}{2\pi i} \int_{c-iT}^{c+iT} y^s \frac{ds}{s} \right| \leq y^c$.

Proof for $y = 1$ omitted. □

Corollary 20.4.2. *The partial sum of the coefficients of a Dirichlet series is given by*

$$\sum_{n < x} a_n + \frac{a_x}{2} (x \in \mathbb{N}_0) = \frac{1}{2\pi i} \lim_{T \rightarrow \infty} \int_{c-iT}^{c+iT} x^s f(s) \frac{ds}{s}.$$

The error from truncating the integral is

$$\left| \left(\sum_{n < x} a_n + \frac{a_x}{2} (x \in \mathbb{N}_0) \right) - \left(\frac{1}{2\pi i} \int_{c-iT}^{c+iT} x^s f(s) \frac{ds}{s} \right) \right| \leq \sum_{n=1}^{\infty} \left(\frac{x}{n} \right)^c a_n \min \left(1, \frac{1}{T \left| \ln \left(\frac{x}{n} \right) \right|} \right).$$

Chapter 21

Zeta functions and the prime number theorem

1 Prime number theorem: Outline

Definition 21.1.1: Define the prime-counting function

$$\pi(x) = |\{p \leq x : p \text{ prime}\}|.$$

Our goal in this chapter is to prove the following famous theorem (in all its error-bounded glory).

Theorem 21.1.2 (Prime number theorem). *There is an effective constant $C > 0$ such that*

$$\pi(x) = \text{li}(x) + O(xe^{-C\sqrt{\ln x}})$$

for all $x \geq 1$.

Here $\text{li}(x)$ denotes the **logarithmic integral**

$$\text{li}(x) = \int_2^x \frac{dt}{\ln t}.$$

Note that $\text{li}(x) = \frac{x}{\ln x} + O\left(\frac{x}{(\ln x)^2}\right)$ as $x \rightarrow \infty$, since integration by parts gives

$$\begin{aligned} \text{li}(x) &= \int_2^x \frac{dy}{\ln y} + O(1) = \frac{x}{\ln x} + \int_2^x \frac{dy}{(\ln y)^2} + O(1) \\ &= \frac{x}{\ln x} + O\left(\frac{x}{(\ln x)^2}\right). \end{aligned} \tag{21.1}$$

1.1 The big picture

We recommend Andrew Granville's article IV.2 Analytic Number Theory in [GBGL10] for an overview.

How might we guess at the asymptotics for $\pi(x)$? (In particular, why is it closer to $\text{li}(x)$ than $\frac{x}{\ln x}$?) By studying tables of primes up to 3 million, Gauss hypothesized that the density of primes at around x is around $\frac{1}{\ln x}$, and hence that the number of primes up to x would be the integral $\text{li}(x) = \int_2^x \frac{dt}{\ln t}$. Making a table of $\pi(x)$ and the difference $\text{li}(x) - \pi(x)$, we find that the difference is slightly more than on the order of \sqrt{x} , so this seems to be a good estimate.

It is a common theme in analytic number theory to make conjectures about the distribution of primes (or other subsets of interest) by assuming they are randomly distributed according to some probability model. Often a simple model works for simple asymptotics up to x , and the model needs to be refined or corrected when dealing with more complicated quantities such as number of primes in a small interval, or spacing between primes.

Model 21.1.3 (Gauss-Cramér model): For $n \geq 3$, let X_n be the random variable such that

$$\begin{aligned} X_n &= 1 \text{ with probability } \frac{1}{\ln n} \\ X_n &= 0 \text{ with probability } 1 - \frac{1}{\ln n}. \end{aligned}$$

Then the sequence X_n behaves similarly to the sequence

$$\begin{aligned} a_n &= 1 \text{ if } n \text{ is prime} \\ a_n &= 0 \text{ otherwise.} \end{aligned}$$

The Gauss-Cramér model exactly predicts $\pi(x) \sim \text{li}(x)$. The model gives more than just the asymptotics of $\pi(x)$, though, it can also be used to think about primes in short intervals $\pi(x+y) - \pi(x)$.

Problem 21.1.4: What are the shortcomings of the Gauss-Cramér model?

1.2 Main steps

The main steps in the proof are as follows.

1. When we have a Dirichlet series

$$F(s) = \sum_{n=0}^{\infty} a_n n^{-s},$$

we can get estimates for $\sum_{n=0}^N a_n$ by “plucking out” those coefficients: The equation

$$\frac{1}{2\pi i} \lim_{T \rightarrow \infty} \int_{c-iT}^{c+iT} y^s \frac{ds}{s} = \begin{cases} 1, & \text{if } y > 1 \\ \frac{1}{2}, & \text{if } y = 1 \\ 0, & \text{if } y < 1. \end{cases}$$

gives

$$\frac{1}{2\pi i} \lim_{T \rightarrow \infty} \int_{c-iT}^{c+iT} x^s f(s) \frac{ds}{s} = \sum_{n < x} a_n + \frac{a_x}{2} (x \in \mathbb{N}_0).$$

We use the more precise statement giving error bounds (Corollary 1).

We want a Dirichlet series where the sum of the first N terms is related to $\pi(N)$. Let

$$\zeta(s) = \prod_{p \text{ prime}} \frac{1}{1 - p^{-s}} = \sum_{n=1}^{\infty} \frac{1}{n^s}.$$

We consider the function

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_{p \text{ prime}} \frac{(\ln p)p^{-s}}{1 - p^{-s}} = \sum_{n=1}^{\infty} \Lambda(n)n^{-s}.$$

We use this function because $\psi(x) := \sum_{n < x} \Lambda(n)$ gives information on $\pi(x)$, and $-\frac{\zeta'}{\zeta}$ continues into a meromorphic function on \mathbb{C} (since ζ does). We now have the estimate

$$\psi(x) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} -\frac{\zeta'(s)}{\zeta(s)} x^s \frac{ds}{s} + (\text{error}).$$

2. We know ζ has analytic continuation (Theorem 21.2.2). Hence we can move the path of integration to $c < 0$. From Cauchy’s integral formula, we get extra terms from the horizontal integrals (integrals involving $-\frac{\zeta'}{\zeta}$) and terms $\frac{x^\rho}{\rho}$ from Cauchy’s integral theorem from the zeros of $\zeta(s)$. *This is why we care about its zeros!* Zeros with large real part contribute large error terms. We will need the following.

- (a) We apply the product development (Theorem 1) on $\xi(s) = \pi^{-\frac{s}{2}} \zeta(s) \Gamma\left(\frac{s}{2}\right)$ to obtain

$$\frac{\zeta'(s)}{\zeta(s)} = \sum_{\rho \text{ zero of } \zeta} \left(\frac{1}{s - \rho} + \frac{1}{\rho} \right) + \dots$$

(Theorem 21.2.5).

- (b) Using the above equation for $\frac{\zeta'}{\zeta}$, we calculate the asymptotics of $N(T)$, the number of zeros in $\{\sigma + it : (\sigma, t) \in [0, 1] \times [-T, T]\}$ (Theorem 21.3.2).
- (c) From (a) to (b) we get a zero-free region for ζ (which includes $\Re s \geq 1$) (Theorems 21.3.1 and 21.3.3).

From the zero-free region we get a bound for $\sum \frac{x^{\rho}}{\rho}$, as well as the horizontal integrals. If the Riemann hypothesis is true, then we can enlarge our zero-free region to $\Re s > \frac{1}{2}$, which is even better.

3. Finally we use the estimate for $\psi(x)$ to get an estimate for $\pi(x)$ (Lemma 21.4.2).

2 Riemann zeta function

Definition 21.2.1: The **Riemann zeta function** is defined by

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

when $\Re s > 1$. This will be generalized to L -functions $L(s, \chi)$ in Definition 1.

By Theorem 20.2.3 and by unique factorization in \mathbb{Z} , we can write

$$\zeta(s) = \prod_{p \text{ prime}} \frac{1}{1 - p^{-s}}.$$

By taking the logarithmic derivative, we have

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_p \frac{d}{ds} \ln(1 - p^{-s}) = \sum_p (\ln p) \frac{p^{-s}}{1 - p^{-s}} = \sum_p \ln p \sum_{k=1}^{\infty} p^{-ks}.$$

Interchanging order of summation gives

$$-\frac{\zeta'(s)}{\zeta(s)} = \sum_{n=1}^{\infty} \Lambda(n) n^{-s}, \quad \Re s > 1, \quad (21.2)$$

where the von Mangoldt function $\Lambda(n)$ is defined as

$$\Lambda(n) = \begin{cases} \ln p, & n = p^r, p \text{ prime}, r \in \mathbb{N}. \\ 0, & \text{else} \end{cases}$$

The most important property of ζ is its analytic continuation and functional equation.

Theorem 21.2.2 (Analytic continuation and functional equation for ζ). $\zeta(s)$ can be analytically continued to a meromorphic function with a simple pole at $s = 0, 1$. It satisfies the functional equation

$$\zeta(s) = 2(2\pi)^{s-1} \Gamma(1-s) \sin\left(\frac{\pi s}{2}\right) \zeta(1-s).$$

Letting $\xi(s) = \pi^{-\frac{s}{2}} \zeta(s) \Gamma\left(\frac{s}{2}\right)$, we have¹

$$\xi(s) = \xi(1-s).$$

Moreover, $\zeta(s)$ has zeros $-2\mathbb{N}$ (the trivial zeros); all other zeros are in the critical strip $0 \leq \Re s \leq 1$.

¹The factor $\Gamma\left(\frac{s}{2}\right)$ can be thought of as coming from the infinite place—see Chapter ??.

To prove this, we first need the transformation law for the theta function; we will show the functional equation for ζ by writing it in terms of θ . As we will prove a more generalized transformation law, we will postpone the proof for θ .

Definition 21.2.3: Define the **theta function** by

$$\theta(u) = \sum_{n \in \mathbb{Z}} e^{-\pi n^2 u}, \quad \Re u > 0.$$

Proposition 21.2.4 (Transformation law for θ): For all u with $\Re u > 0$,

$$\theta\left(\frac{1}{u}\right) = u^{\frac{1}{2}}\theta(u).$$

This is a special case of Proposition 22.2.4.

Proof of Theorem 21.2.2. We first analytically continue ζ to $\Re s > 0$, show the functional equation is true for $0 < \Re s < 1$, and use it to establish analytic continuation to \mathbb{C} .

Note

$$\zeta(s) = \frac{1}{s-1} + \sum_{n=1}^{\infty} \left[n^{-s} - \int_n^{n+1} x^{-s} dx \right] = \frac{1}{s-1} + \sum_{n=1}^{\infty} \int_n^{n+1} (n^{-s} - x^{-s}) dx \quad (21.3)$$

Since for $n \leq x \leq n+1$ we have

$$\begin{aligned} |n^{-s} - x^{-s}| &= \left| \int_n^x s x^{-s-1} dx \right| \leq |s| n^{-s-1} \\ \left| \int_n^{n+1} n^{-s} - x^{-s} dx \right| &\leq |s| n^{-s-1}, \end{aligned} \quad (21.4)$$

the sum (21.3) converges uniformly locally for $\Re s > 0$ and extends ζ to an analytic function for $\Re s > 0$.

We claim that

$$2\xi(s) = \int_0^{\infty} (\theta(u) - 1) u^{\frac{s}{2}} \frac{du}{u}, \quad \Re s > 1 \quad (21.5)$$

Indeed, we have

$$\begin{aligned} \int_0^{\infty} (\theta(u) - 1) u^{\frac{s}{2}} \frac{du}{u} &= \int_0^{\infty} 2 \sum_{n=1}^{\infty} e^{-\pi n^2 u} u^{\frac{s}{2}} \frac{du}{u} \\ &= 2 \sum_{n=1}^{\infty} \int_0^{\infty} e^{-\pi n^2 u} u^{\frac{s}{2}} \frac{du}{u} \\ &= 2 \sum_{n=1}^{\infty} \int_0^{\infty} e^{-u} \left(\frac{u}{\pi n^2} \right)^{\frac{s}{2}} \frac{du}{u} && u \leftarrow \frac{u}{\pi n^2} \\ &= 2\pi^{-\frac{s}{2}} \left(\sum_{n=1}^{\infty} \frac{1}{n^s} \right) \left(\int_0^{\infty} e^{-u} u^{\frac{s}{2}} \frac{du}{u} \right) \\ &= 2\pi^{-\frac{s}{2}} \zeta(s) \Gamma\left(\frac{s}{2}\right) = 2\xi(s). \end{aligned}$$

The theta transformation law 21.2.4 give that for $\Re s > 1$,

$$\begin{aligned} 2\xi(s) &= \int_0^1 (\theta(u) - 1)u^{\frac{s}{2}} \frac{du}{u} + \int_1^\infty (\theta(u) - 1)u^{\frac{s}{2}} \frac{du}{u} \\ &= \int_1^\infty \left(\theta\left(\frac{1}{u}\right) - 1 \right) u^{\frac{s}{2}} \frac{du}{u} + \int_1^\infty (\theta(u) - 1)u^{\frac{s}{2}} \frac{du}{u} && u \leftrightarrow \frac{1}{u} \\ &= \int_1^\infty \left(u^{-\frac{1}{2}} \theta\left(\frac{1}{u}\right) - 1 \right) u^{\frac{1-s}{2}} \frac{du}{u} + \int_1^\infty (u^{\frac{1-s}{2}} - u^{-\frac{s}{2}}) \frac{du}{u} + \int_1^\infty (\theta(u) - 1)u^{\frac{s}{2}} \frac{du}{u} \\ &= -\frac{2}{s} - \frac{2}{1-s} + \int_1^\infty (\theta(u) - 1)u^{\frac{1-s}{2}} \frac{du}{u} + \int_1^\infty (\theta(u) - 1)u^{\frac{s}{2}} \frac{du}{u}. \end{aligned}$$

The last expression converges for all $\Re s > 0$, so in fact equals $2\zeta(s)$ for all $\Re s > 0$ by uniqueness of analytic continuation. Since the last expression is symmetric under $1 - s \mapsto s$, the functional equation for ξ follows.

The functional equation for ξ gives

$$\begin{aligned} \zeta(s) &= \pi^{\frac{s}{2}} \Gamma\left(\frac{s}{2}\right)^{-1} \pi^{-\frac{1-s}{2}} \Gamma\left(\frac{1-s}{2}\right) \zeta(1-s) \\ &= \pi^{s-\frac{1}{2}} \frac{\Gamma\left(\frac{1-s}{2}\right)}{\Gamma\left(\frac{s}{2}\right)} \zeta(1-s) \\ &= \pi^{s-\frac{1}{2}} \Gamma\left(\frac{1-s}{2}\right) \Gamma\left(1-\frac{s}{2}\right) \frac{\sin\left(\frac{\pi s}{2}\right)}{\pi} \zeta(1-s) && \text{by Proposition 19.7.2(5)} \\ &= 2(2\pi)^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s) && \text{by Proposition 19.7.2(6)} \end{aligned}$$

Finally, the statement about zeros follows from the fact that ζ has no zeros with $\Re s > 1$ (as $\frac{\zeta'}{\zeta}$ is holomorphic there) and the functional equation, noting $\sin\left(\frac{\pi s}{2}\right) = 0$ exactly when s is an even integer, with the zero at $s = 0$ cancelled by the pole at 1 of ζ . \square

Theorem 21.2.5 (Product development of ξ). *The function $(s^2 - s)\xi(s)$ is entire of order 1, and $\xi(s)$ has the product expansion*

$$\xi(s) = \frac{e^{A+Bs}}{s^2 - s} \prod_{\rho \text{ zero of } \zeta} \left(1 - \frac{s}{\rho}\right) e^{\frac{s}{\rho}}.$$

Then $\frac{\zeta'}{\zeta}(s)$ has the partial-fraction expansion

$$\frac{\zeta'}{\zeta}(s) = B - \frac{1}{s-1} + \frac{1}{2} \ln(\pi) - \frac{1}{2} \frac{\Gamma'}{\Gamma}\left(\frac{s}{2} + 1\right) + \sum_{\rho \text{ nontrivial zero of } \zeta} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right).$$

From now on, unless otherwise specified, when we say zero of ζ we mean *nontrivial* zero.

Proof. Note $(s^2 - s)\xi(s)$ is entire because ξ only has 2 simple poles at 0, 1. To show it has order 1 we need two inequalities.

Step 1: There is no constant C so that $(s^2 - s)\xi(s) \lesssim e^{C|s|}$: Indeed, for real s and any constant C , by Stirling's approximation 19.7.4 we have

$$\begin{aligned}(s^2 - s)\xi(s) &= (s^2 - s)\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)\zeta(s) \\ &\gtrsim s^{-\frac{1}{2}}\left(\frac{s}{2e\pi}\right)^{\frac{s}{2}} \gtrsim e^{Cs}.\end{aligned}$$

Step 2: There is a constant C so that $(s^2 - s)\xi(s) \lesssim e^{C|s|\ln|s|}$: $e^{|s|\ln|s|} \geq 1$ for all s so it suffices to prove this for sufficiently large s . By the integral and sum formulas for Γ and ξ , and the fact that $|x^s| = |x^{\Re s}|$, we have

$$|\xi(\sigma + ti)| \leq \pi^{-\frac{\sigma}{2}}\Gamma\left(\frac{\sigma}{2}\right)\zeta(\sigma), \quad \sigma > 1.$$

By symmetry of ξ it suffices to consider $\sigma \geq \frac{1}{2}$. (“Nudging” $|s|$ in $e^{C|s|\ln|s|}$ by a constant changes it by at most a constant factor.) Consider 2 cases.

1. $\sigma > 2$: Then $\pi^{-\frac{\sigma}{2}} < 1$ and $\zeta(\sigma) < \zeta(2)$ so by Stirling's approximation 19.7.4,

$$|\xi(\sigma + ti)| \lesssim \Gamma\left(\frac{\sigma}{2}\right) = e^{(\ln\Gamma)(\sigma)} = e^{\left(\frac{\sigma}{2}-1\right)\ln\frac{\sigma}{2}-\frac{\sigma}{2}+O(1)}$$

from which the result follows.

2. $\frac{1}{2} \leq \sigma \leq 2$: From (21.4), we have for s bounded away from 1,

$$\zeta(s) \leq O(1) + |s| \sum_{n=1}^{\infty} n^{-\frac{3}{2}} = O(|s|).$$

This time $\Gamma\left(\frac{\sigma}{2}\right) = O(1)$ so

$$|(s^2 - s)\xi(s)| \leq \left|s^2\pi^{-\frac{\sigma}{2}}\zeta(s)\Gamma\left(\frac{\sigma}{2}\right)\right| = O(|s|^3) \lesssim e^{C|s|\ln|s|}.$$

This shows $(s^2 - s)\xi(s)$ has order 1.

Step 3: By the product development 19.6.3, noting the the zeros of $(s^2 - s)\xi$ are the nontrivial zeros of ζ (since Γ has no zeros and trivial zeros of ζ come from the poles of Γ in the definition of ξ), we get

$$(s^2 - s)\xi(s) = e^{A+Bs} \prod_{\rho \text{ zero of } \zeta} \left(1 - \frac{s}{\rho}\right) e^{\frac{s}{\rho}}.$$

Dividing by $s^2 - s$ and log-differentiating gives

$$\frac{\xi'}{\xi}(s) = B - \frac{1}{s} - \frac{1}{s-1} + \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right).$$

Since $\zeta(s) = \pi^{\frac{s}{2}} \Gamma\left(\frac{s}{2}\right)^{-1} \xi(s)$, we get

$$\begin{aligned} \frac{\zeta'(s)}{\zeta(s)} &= \frac{1}{2} \ln \pi + \frac{1}{2} \frac{\Gamma'}{\Gamma}\left(\frac{s}{2}\right) + B - \frac{1}{s} - \frac{1}{s-1} + \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right) \\ &= \frac{1}{2} \ln \pi + \frac{1}{2} \frac{\Gamma'}{\Gamma}\left(\frac{s}{2} + 1\right) + B - \frac{1}{s-1} + \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right), \quad \Gamma(z) = \frac{\Gamma(z+1)}{z} \square \end{aligned}$$

3 Zeros of zeta

Note that from the function equation, $\zeta(s)$ has simple zeros at $-2\mathbb{N}$. We call these trivial zeros. More importantly for us are the zeros with real part in $[0, 1]$.

Denote by $N(T)$ be the number of zeros of ζ in $\{\sigma + it : (\sigma, t) \in [0, 1] \times [-T, T]\}$, counting multiplicity. We first give asymptotics on the vertical distribution of zeros of ζ (von Mangoldt's formula, Theorem 21.3.2), then give a zero-free region for ζ (Theorem 21.3.3).

Lemma 21.3.1. *Define $\mathcal{L}(t) = \ln(|t| + 2)$. For $s = \sigma + it$ with $\sigma \in [-1, 2]$, we have²*

$$\begin{aligned} \frac{\zeta'(s)}{\zeta(s)} &= -\frac{1}{s-1} + \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right) + O(\mathcal{L}) \\ &= -\frac{1}{s-1} + \sum_{|\Im(s-\rho)| < 1} \frac{1}{s-\rho} + O(\mathcal{L}). \end{aligned} \tag{21.6}$$

Moreover, there are $O(\mathcal{L})$ zeros ρ with $|\Im(s-\rho)| < 1$, i.e. the number of zeros with imaginary part in $[t, t+1]$ is $O(\ln t)$, as $t \rightarrow \infty$.

Note this gives $N(T) = O(T \ln T)$. The next theorem will give an improvement of this estimate.

Proof. Our strategy is this: at a point where we know $\frac{\zeta'}{\zeta}$ is bounded ($s = 2 + it$), we use Theorem 21.2.5 to get information on how many zeros of ζ can be close to s . Then we use compare $\frac{\zeta'}{\zeta}(\sigma + it)$ with $\frac{\zeta'}{\zeta}(2 + it)$ to get the general estimate.

Step 1: Theorem 21.2.5 gives us

$$\frac{\zeta'(s)}{\zeta(s)} = -\frac{1}{s-1} + \underbrace{B + \frac{1}{2} \ln \pi}_{O(1)} - \frac{1}{2} \underbrace{\frac{\Gamma'}{\Gamma}\left(\frac{s}{2} + 1\right)}_{(A)} + \underbrace{\sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right)}_{(B)}. \tag{21.7}$$

From Stirling's approximation 1, (A) equals

$$\ln \left| \frac{\sigma}{2} + 1 + i \frac{t}{2} \right| + O(1) = O(\mathcal{L}) \tag{21.8}$$

²Note $\frac{1}{s-1} = O(1)$ when s is bounded away from 1.

These two equations show (21.6).

Now suppose $s = 2 + it$. Note that

$$\left| \frac{\zeta'(2 + it)}{\zeta(2 + it)} \right| = \left| \sum_{n=1}^{\infty} \Lambda(n)n^{-2-it} \right| \leq \left| \sum_{n=1}^{\infty} (\ln n)n^{-2} \right| < \infty,$$

so the LHS of (21.7) is $O(1)$. Hence (21.7) becomes

$$O(\mathcal{L}) = \sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{\rho} \right). \tag{21.9}$$

We estimate the terms with $|\Im(s - \rho)| < 1$ by a constant to show that there aren't too many of them. From (21.9) and (21.8),

$$\begin{aligned} O(\mathcal{L}) &= \Re \sum_{\rho} \left(\frac{1}{2 + it - \rho} + \frac{1}{\rho} \right) \\ &\geq \Re \sum_{\rho} \left(\frac{(2 - \Re \rho) - (t - \Im \rho)i}{(2 - \Re \rho)^2 + (t - \Im \rho)^2} \right) && \text{since } \Re \left(\frac{1}{\rho} \right) > 0 \\ &\geq \sum_{\rho} \frac{1}{4 + (t - \Im \rho)^2} && \text{since } 0 \leq \Re \rho \leq 1 \\ &\geq \frac{1}{5} |\{\rho : |\Im(s - \rho)| < 1\}| + \frac{1}{5} \sum_{|\Im(s - \rho)| \geq 1} \frac{1}{(t - \Im \rho)^2}. \end{aligned} \tag{21.10}$$

This proves the second part of the lemma.

Step 2: Now we consider general $s = \sigma + it$, by comparing it to $2 + it$. We have by (21.7) and (21.8) that

$$\begin{aligned} &\frac{\zeta'(s)}{\zeta(s)} - \underbrace{\frac{\zeta'(2 + it)}{\zeta(2 + it)}}_{O(1)} \\ &= -\frac{1}{s - 1} + O(1) + \underbrace{\frac{1}{2} \left(\ln \left| \frac{\sigma}{2} + 1 + \frac{t}{2}i \right| - \ln \left| 2 + \frac{t}{2}i \right| \right)}_{O(1)} + \sum_{\rho} \left(\frac{1}{s - \rho} - \frac{1}{2 + it - \rho} \right) \\ &= -\frac{1}{s - 1} + O(1) + \sum_{|\Im(s - \rho)| < 1} \frac{1}{s - \rho} - \underbrace{\sum_{|\Im(s - \rho)| < 1} \frac{1}{2 + it - \rho}}_{O(\mathcal{L})} + \underbrace{\sum_{|\Im(s - \rho)| \geq 1} \frac{(2 - \sigma)}{(s - \rho)(2 + it - \rho)}}_{O(\mathcal{L})}. \end{aligned}$$

The first $O(\mathcal{L})$ is because there are at most $O(\mathcal{L})$ terms and each term is at most 1 in absolute value; the second is from

$$\sum_{|\Im(s - \rho)| \geq 1} \frac{2 - \sigma}{(s - \rho)(2 + it - \rho)} = O \left(\sum_{|\Im(s - \rho)| \geq 1} \frac{1}{\Im(s - \rho)^2} \right) = O(\mathcal{L});$$

the first equality is from $2 - \sigma = O(1)$ and $\Im(s - \rho) = \Im(2 + it - \rho)$; the second is by (21.10). \square

Theorem 21.3.2 (von Mangoldt). (*) As $T \rightarrow \infty$,

$$N(T) = \frac{T}{\pi} \ln \left(\frac{T}{2\pi} \right) - \frac{T}{\pi} + O(\ln T).$$

Proof. As ζ has only a countable number of zeros, we may assume T is not the imaginary part of any zero.

Let

$$\mathcal{R} = \{\sigma + it : (s, t) \in [-1, 2] \times [-T, T]\}$$

and let C be the boundary of \mathcal{R} . From $\xi(s) = \pi^{-\frac{s}{2}} \zeta(s) \Gamma(\frac{s}{2})$, we see that ξ has the same zeros as ζ in this region, and simple poles at 0 and 1. Hence by Cauchy's residue formula 1,

$$\frac{1}{2\pi i} \oint_C \frac{\xi'(s)}{\xi(s)} ds = 2N(T) - 2.$$

Noting that $\xi(\bar{s}) = \overline{\xi(s)}$ and $\xi(s) = \xi(1-s)$, changes of variable show that the integral on each of the sections of C between $2, \frac{1}{2} + iT, -1,$ and $\frac{1}{2} - iT$ are the same.³ Let C' be the part from 1 to $\frac{1}{2} + iT$. Thus the above equals

$$\begin{aligned} \frac{2}{\pi i} \int_{C'} \frac{\xi'(s)}{\xi(s)} ds &= \frac{2}{\pi i} \int_{C'} -\frac{\ln \pi}{2} + \frac{\zeta'(s)}{\zeta(s)} + \frac{(\Gamma(\frac{s}{2}))'}{\Gamma(\frac{s}{2})} ds && \frac{(\prod_{k=1}^n f_k)'}{\prod_{k=1}^n f_k} = \sum_{k=1}^n \frac{f_k'}{f_k} \\ &= \frac{2}{\pi} \Im \int_{C'} -\frac{\ln \pi}{2} + \frac{\zeta'(s)}{\zeta(s)} + \frac{(\Gamma(\frac{s}{2}))'}{\Gamma(\frac{s}{2})} ds && \text{(expression is real).} \end{aligned}$$

We break this up into 3 integrals and estimate each part separately.

1. $\Im \int_{C'} -\frac{\ln \pi}{2} ds = -\frac{T}{2} \ln \pi$.
2. Using the estimate for $\frac{\zeta'}{\zeta}$ in Lemma 21.3.1, we evaluate the second integral. Note that $\ln \zeta$ is defined for $\Re s > 1$ and is uniformly bounded for $\Re s = 2$:

$$\begin{aligned} (\ln \zeta)(s) &= \sum_{p \text{ prime}} \ln(1 - p^{-s}) \\ |(\ln \zeta)(2 + it)| &\leq \sum_{p \text{ prime}} 2p^{-2}. \end{aligned}$$

(Just bound \ln linearly near 1, or expand in Taylor series.) Note $\ln(x - \rho)$ is well-defined on C' for any ρ . Hence by Theorem 21.3.1,

$$\begin{aligned} \Im \int_{C'} \frac{\zeta'}{\zeta}(s) ds &= (\Im(\ln \zeta)(2 + iT) - \Im(\ln \zeta)(2)) + \int_{2+iT}^{\frac{1}{2}+iT} \frac{\zeta'}{\zeta}(s) ds \\ &= O(1) + \int_{2+iT}^{\frac{1}{2}+iT} \Im \left(\sum_{|\Im(s-\rho)| < 1} \frac{1}{s - \rho} \right) + O(\ln T) ds \\ &= O(\ln T) + \sum_{|\Im(s-\rho)| < 1} \Im(\ln(x - \rho)) \Big|_{2+Ti}^{\frac{1}{2}+Ti} \\ &\leq O(\ln T) + 2\pi O(\ln T) \end{aligned}$$

³We used ξ because its symmetry allows us to do this.

since there are at most $\ln T$ terms in the sum.

3. We estimate the last integral using Stirling’s formula 1. (Note that $\ln \Gamma$ is well-defined for $s \in C'$.)

$$\begin{aligned} \int_{C'} \frac{\left(\Gamma\left(\frac{s}{2}\right)\right)'}{\Gamma\left(\frac{s}{2}\right)} &= \left[\Im(\ln \Gamma)\left(\frac{s}{2}\right)\right]_2^{\frac{1}{2}+Ti} \\ &= \Im(\ln \Gamma)\left(\frac{1}{4} + \frac{T}{2}i\right) \\ &= \Im\left[\left(-\frac{1}{4} + \frac{T}{2}i\right) \ln\left(\frac{1}{4} + \frac{T}{2}i\right) - \left(\frac{1}{4} + \frac{T}{2}i\right) + O(1)\right] \\ &= \frac{T}{2} \ln\left(\frac{T}{2}\right) - \frac{T}{2} + O(1). \end{aligned} \quad \square$$

Now put everything together to get

$$\begin{aligned} N(T) - 2 &= \frac{2}{\pi} \left(-\frac{T}{2} \ln \pi + O(\ln T) + \left(\frac{T}{2} \ln\left(\frac{T}{2}\right) - \frac{T}{2} + O(1) \right) \right) \\ N(T) &= \frac{T}{\pi} \ln\left(\frac{T}{2\pi}\right) - \frac{T}{\pi} + O(\ln T). \end{aligned}$$

Theorem 21.3.3 (Zero-free region for ζ). *There are no zeros of ζ with $\Re s \geq 1$. Moreover, there is a constant $c > 0$ such that for $|t| > 2$, every zero $\sigma + it$ satisfies*

$$\sigma < 1 - \frac{c}{\ln |t|}.$$

Proof. We already noted ζ has no zero for $\Re s > 1$ (Theorem 21.2.2), so for the first part it suffices to prove that no zero has real part 1.

If ζ had a zero $1 + it$, then $\frac{\zeta'}{\zeta}$ would have a pole of positive residue at $1 + it$. For $s = \sigma + it$, $\sigma > 1$ we have $-\frac{\zeta'}{\zeta}(s) = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s}$, so this means that as $\sigma \rightarrow 1^+$, many of the important terms would have n^{-it} “close” to -1 , to make it blow up in the negative direction. For those terms, we have n^{-2it} “close” to 1. This would force $-\frac{\zeta'}{\zeta}(\sigma + 2ti)$ to have a pole of positive residue at $1 + 2ti$, i.e ζ to have a pole at $1 + 2ti$, contradicting the fact that it is analytic there.

We now make this idea precise. What we want is an inequality between some function of an angle and its double, so that if one is small it forces the other to be large. So we consider

$$0 \leq 2(1 + \cos \theta)^2 = 3 + 4 \cos \theta + \cos 2\theta.$$

This gives

$$0 \leq 3 + 4\Re(n^{-it}) + \Re(n^{-2it}).$$

Multiplying by $\Lambda(n)n^{-\sigma}$ and summing, we get

$$0 \leq 3 \left(-\frac{\zeta'}{\zeta}(\sigma) \right) + 4\Re \left(-\frac{\zeta'}{\zeta}(\sigma + ti) \right) + \Re \left(-\frac{\zeta'}{\zeta}(\sigma + 2ti) \right), \quad \sigma > 1. \tag{21.11}$$

Letting r be the degree of the zero at $1 + ti$, we have by Lemma 21.3.1

$$0 \leq \left(\frac{3}{\sigma - 1} + O(1) \right) - \left(\frac{4r}{\sigma - 1} + O(\mathcal{L}) \right) + \Re \left(-\frac{\zeta'}{\zeta}(\sigma + 2ti) \right) \text{ as } \sigma \rightarrow 1^+.$$

If $r \geq 1$, then this gives $-\frac{\zeta'}{\zeta}(\sigma + 2ti) \rightarrow \infty$ as $\sigma \rightarrow 1^+$, contradiction. Hence $r = 0$; $1 + it$ is not a zero.

For the second statement, we have to use the partial fraction decomposition 21.2.5. Suppose $\rho = (1 - \delta) + it$ is a zero. By Lemma 21.3.1, we have

$$-\frac{\zeta'(s)}{\zeta(s)} = O(\ln |t|) - \sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{\rho} \right) \leq O(\ln |t|) - \frac{1}{s - \rho}.$$

Then

$$\begin{aligned} -\Re \frac{\zeta'}{\zeta}(\sigma + ti) &\leq O(\ln |t|) - \frac{1}{\sigma + \delta - 1} \\ -\Re \frac{\zeta'}{\zeta}(\sigma + 2ti) &\leq O(\ln |2t|) = O(\ln |t|). \end{aligned}$$

For $\sigma > 1$, plugging this into (21.11) gives

$$\begin{aligned} 0 &\leq \frac{3}{\sigma - 1} + O(\ln |t|) - \frac{4}{\sigma + \delta - 1} \\ \implies \frac{4}{\sigma + \delta - 1} &< \frac{3}{\sigma - 1} + C_1 \ln |t| \end{aligned}$$

for some C_1 . Now take $\sigma = 1 + 4\delta$ to get

$$\frac{4}{5\delta} < \frac{3}{4\delta} + C_1 \ln |t|,$$

giving

$$\delta > \frac{1}{20C_1 \ln |t|}$$

as needed. □

4 Prime number theorem: proof

Now we gather everything together to prove the prime number theorem. We first show the following.

Theorem 21.4.1 (von Mangoldt's formula). *For an integer $x > 2$ and $x \geq T$,*

$$\psi(x) = x - \sum_{|\Im(\rho)| < T} \frac{x^\rho}{\rho} + O\left(\frac{x(\ln x)^2}{T}\right). \quad (21.12)$$

Proof. Step 1: We estimate $\psi(x)$ using Theorem 20.4.2. Suppose x is an integer; the theorem gives

$$\begin{aligned} \left| \psi(x) - \left(\int_{c-iT}^{c+iT} x^s \left(-\frac{\zeta'}{\zeta}(s) \frac{ds}{s} \right) \right) \right| &\leq \Lambda(x) + \sum_{n \geq 1, n \neq x} \left(\frac{x}{n} \right)^c \Lambda(n) \frac{1}{T \left| \ln \left(\frac{x}{n} \right) \right|} \\ &\leq \ln(x) + \sum_{n \geq 1, n \neq x} \left(\frac{x}{n} \right)^c \frac{\ln(n)}{T \left| \ln \left(\frac{x}{n} \right) \right|}. \end{aligned}$$

Take

$$c = 1 + \frac{1}{\ln x}.$$

Note that this makes $x^c = ex = O(x)$. To estimate the sum we split it into several parts.

1. $1 \leq n < \frac{x}{e}$: We have

$$\begin{aligned} \sum_{1 \leq n < \frac{x}{e}} \left(\frac{x}{n} \right)^c \frac{\ln n}{T \left| \ln \left(\frac{x}{n} \right) \right|} &\lesssim \frac{x \ln x}{T} \sum_{1 \leq n < x} \frac{1}{n} \\ &\sim \frac{x(\ln x)^2}{T}. \end{aligned}$$

2. $\frac{x}{e} \leq n < ex$: We have

$$\begin{aligned} \sum_{\frac{x}{e} \leq n < ex, n \neq x} \left(\frac{x}{n} \right)^c \ln n \frac{1}{T \left| \ln \left(\frac{x}{n} \right) \right|} &\lesssim \sum_{\frac{x}{e} \leq n < ex, n \neq x} e^{1 + \frac{1}{\ln x}} \frac{\ln n}{T \left| \ln \left(\frac{x}{n} \right) \right|} \\ &\lesssim \frac{1}{T} \sum_{\frac{x}{e} \leq n < ex, n \neq x} \frac{\ln x}{\left| 1 - \frac{x}{n} \right|} && \text{using } \ln x \sim x - 1 \text{ when } x \approx 1 \\ &\lesssim \frac{x \ln x}{T} \sum_{\frac{x}{e} \leq n < ex, n \neq x} \frac{1}{|n - x|} \\ &\lesssim \frac{x \ln x}{T} \sum_{1 \leq n < (e-1)x} \frac{1}{n} \\ &\sim \frac{x(\ln x)^2}{T}. \end{aligned}$$

3. $n \geq ex$: We have

$$\begin{aligned} \sum_{n \geq ex} \left(\frac{x}{n} \right)^c \frac{\ln n}{T} &< \frac{x}{T} \int_{ex-1}^{\infty} \frac{\ln y}{y^c} dy && \frac{\ln y}{y^c} \text{ decreasing for } y > e \\ &= \frac{x}{T} \left[\frac{-y^{-c+1} \ln y}{c-1} - \frac{y^{-c+1}}{(c-1)^2} \right]_{ex-1}^{\infty} \\ &\sim \frac{x(\ln x)^2}{T}. \end{aligned}$$

Putting everything together gives

$$\left| \psi(x) - \left(\int_{c-iT}^{c+iT} x^s \left(-\frac{\zeta'}{\zeta}(s) \right) \frac{ds}{s} \right) \right| = O\left(\frac{x(\ln x)^2}{T} + \ln x \right). \tag{21.13}$$

Step 2: We move the line of integration to $\Re s = -1$. Assuming that T is not the imaginary part of any root, by Cauchy's residue theorem 19.4.8

$$\int_{c-iT}^{c+iT} \frac{x^s}{s} \frac{\zeta'}{\zeta}(s) ds + \underbrace{\int_{c+iT}^{-1+iT} \frac{x^s}{s} \frac{\zeta'}{\zeta}(s) ds}_{I_{h,1}} + \underbrace{\int_{-1+iT}^{-1-iT} \frac{x^s}{s} \frac{\zeta'}{\zeta}(s) ds}_{I_v} + \underbrace{\int_{-1-iT}^{c-iT} \frac{x^s}{s} \frac{\zeta'}{\zeta}(s) ds}_{I_{h,2}} = \frac{\zeta'}{\zeta}(0) - x + \sum_{|\Im \rho| < T} \frac{x^\rho}{\rho}.$$

Here $\frac{x^\rho}{\rho}$ are the residues at the zeros, $-x$ comes from the pole of ζ at 1, and $\frac{\zeta'}{\zeta}(0)$ comes from the pole of $\frac{1}{s}$. Then

$$\int_{c-iT}^{c+iT} \frac{x^s}{s} \left(-\frac{\zeta'}{\zeta}(s) \right) ds - x = 1 + I_{h,1} + I_{h,2} + I_v - \sum_{\Im \rho < T} \frac{x^\rho}{\rho}. \tag{21.14}$$

We estimate each summand.

1. For the horizontal integrals, we use the estimate 21.3.1 to get

$$\begin{aligned} \left| \frac{\zeta'}{\zeta}(s) \right| &= \left| \sum_{|\Im(s-\rho)| < 1} \frac{1}{s-\rho} \right| + O(\ln T), \quad s = \sigma + Ti \\ &\leq \sum_{|\Im(s-\rho)| < 1} \frac{1}{\Im(s-\rho)} + O(\ln T). \end{aligned}$$

We would like to bound $\Im(s-\rho)$ away from 0. To do this, note that there are $O(\ln T)$ roots in with $\Im \rho \in [T, T+1]$ by Lemma 21.3.1. Hence by tweaking T slightly⁴, we can assume $|\Im(s-\rho)| > \frac{C}{\ln T}$ for all ρ . Also by Lemma 21.3.1 there are at most $O(\ln T)$ terms in the sum, so the sum is $O((\ln T)^2)$. Integrating gives

$$\begin{aligned} \left| \int_{c \pm Ti}^{-1 \pm Ti} \frac{x^s}{s} \frac{\zeta'}{\zeta}(s) ds \right| &= O((\ln T)^2) O\left(\frac{1}{T} \right) \int_c^{-1} |x^s| ds \\ &= O\left(\frac{(\ln T)^2}{T} \right) O(x) \\ &= O\left(\frac{x(\ln x)^2}{T} \right). \end{aligned}$$

⁴Changing T by a constant does not change the error term of (21.12); moreover the change in the LHS sum is $O\left(\frac{x}{T} \ln T\right) = O\left(\frac{x(\ln x)^2}{T}\right)$.

2. For the vertical integral, we use the same estimate, this time noting that $|s - \rho| > 1$ for every root ρ , since every zero satisfies $\Re \rho > 0$. This gives that $\frac{\zeta'}{\zeta}(s) = O(\ln T)$, and

$$\begin{aligned} \left| \int_{-1+Ti}^{-1-Ti} \frac{x^s}{s} \frac{\zeta'}{\zeta}(s) ds \right| &= O(\ln T) \int_{-1+Ti}^{-1-Ti} \frac{x^{-1}}{|s|} ds \\ &= O\left(\frac{\ln T}{x}\right) \int_{-T}^T \frac{1}{\sqrt{t^2+1}} dt \\ &= O\left(\frac{\ln T}{x}\right) \int_1^{T+1} \frac{1}{t} dt \\ &= O\left(\frac{(\ln T)^2}{x}\right) = O\left(\frac{x(\ln x)^2}{T}\right). \end{aligned}$$

Equations (21.13) and (21.14) together with the above two estimates give the theorem. \square

The final ingredient in the proof of the Prime Number Theorem is the estimate for $\sum_{|\Im(\rho)| < T} \frac{x^\rho}{\rho}$ using the zero-free regions for ζ and the estimate for number of zeros of ζ .

Proof of Theorem 21.1.2. First, note there can only be a finite number of zeros of ζ with $|\Im(\rho)| < 2$, so $\sum_{|\Im(\rho)| < 2} \frac{x^\rho}{\rho} = O(x^r)$ for some fixed $r < 1$.⁵ We estimate $\sum_{2 \leq |\Im(\rho)| < T} \frac{x^\rho}{\rho}$ in two steps.

1. By Theorem 21.3.3, there is c such that for ρ with $2 \leq |\Im(\rho)| < T$,

$$|x^\rho| = x^{\Re \rho} \leq x^{1 - \frac{c}{\ln T}} = x e^{-\frac{c \ln x}{\ln T}}.$$

2. Using $N(T) = O(T \ln T)$ (Theorem 21.3.2 or the weaker remark after Lemma 21.3.1),

$$\begin{aligned} \sum_{2 \leq |\Im(\rho)| < T} \frac{1}{|\rho|} &\leq \sum_{2 \leq |\Im(\rho)| < T} \frac{1}{\Im(\rho)} \\ &\leq \int_2^T \frac{dN(t)}{t} && \text{(Riemann-Stieltjes integral)} \\ &= \frac{N(T)}{T} - \frac{N(2)}{2} + \int_2^T \frac{N(t)}{t^2} dt && \text{integration by parts} \\ &= O(\ln T) + \int_2^T O\left(\frac{\ln t}{t}\right) dt \\ &= O(\ln T) + O((\ln T)^2) = O((\ln T)^2). \end{aligned} \tag{21.15}$$

Putting these two estimates together,

$$\begin{aligned} \left| \sum_{|\Im(\rho)| < T} \frac{x^\rho}{\rho} \right| &\leq O(x^r) + \max_{2 \leq |\Im(\rho)| < T} (|x^\rho|) \sum_{2 \leq |\Im(\rho)| < T} \frac{1}{|\rho|} \\ &\leq O(x^r) + O\left(x e^{-\frac{c \ln x}{\ln T}} (\ln T)^2\right). \end{aligned} \tag{21.16}$$

⁵In fact, there are zero such zeros.

Combining with Theorem 21.4.1, and setting $T = e^{\sqrt{\ln x}}$ (so that $xe^{-\frac{\ln x}{T}} = \frac{x}{T}$), we get

$$\begin{aligned} |\psi(x) - x| &= O\left(x^r + xe^{-\frac{c \ln x}{\ln T}} (\ln T)^2 + \frac{x(\ln x)^2}{T}\right) \\ &= O\left(x^r + xe^{-c\sqrt{\ln x}} \ln x + x(\ln x)^2 e^{-\sqrt{\ln x}}\right) \\ &= O(xe^{-C\sqrt{\ln x}}), \end{aligned}$$

for some $C > 0$. This shows

$$\psi(x) = x + O(xe^{-C\sqrt{\ln x}}). \tag{21.17}$$

Finally, we extract the asymptotics of π from the following.

Lemma 21.4.2. *We have the following estimates:*

$$\begin{aligned} \pi(x) &= \frac{\psi(x)}{\ln x} + \int_2^x \psi(y) \frac{dy}{y(\ln y)^2} + O(x^{\frac{1}{2}}), \\ \psi(x) &= \pi(x) \ln x - \int_2^x \frac{\pi(y)}{y} dy + O(x^{\frac{1}{2}} \ln x). \end{aligned}$$

Proof. Define

$$\gamma(n) = \begin{cases} 1, & n \text{ prime,} \\ 0, & n \text{ not prime,} \end{cases} \quad \Lambda_1(n) = \begin{cases} \ln n, & n \text{ prime,} \\ 0, & n \text{ not prime,} \end{cases}$$

and

$$\psi_1(x) = \sum_{n \leq x} \Lambda_1(n).$$

First note

$$\begin{aligned} |\psi(x) - \psi_1(x)| &= \sum_{2 \leq r \leq \log_2(x)} \sum_{p|p^r \leq x} \ln p \\ &\leq \sum_{2 \leq r \leq \log_2(x)} x^{\frac{1}{r}} \ln x \\ &= O(x^{\frac{1}{2}} \ln x + x^{\frac{1}{3}} (\ln x)^2) = O(x^{\frac{1}{2}} \ln x). \end{aligned} \tag{21.18}$$

Part 1: By partial summation ?? with $u = \Lambda_1$, $U = \psi_1$, and $v = \frac{1}{\ln x}$,

$$\begin{aligned} \pi(x) &= \sum_{n \leq x} \gamma(n) \\ &= \sum_{n \leq x} \Lambda_1(n) \frac{1}{\ln n} \\ &= \frac{\psi_1(x)}{\ln x} + \int_2^x \psi_1(t) \frac{dt}{t(\ln t)^2} \\ &= \frac{\psi(x)}{\ln x} + O(x^{\frac{1}{2}}) + \int_2^x \psi(t) \frac{dt}{t(\ln t)^2} + \int_2^x O(t^{-\frac{1}{2}}) dt \quad \text{by (21.18)} \\ &= \frac{\psi(x)}{\ln x} + \int_2^x \psi(t) \frac{dt}{t(\ln t)^2} + O(x^{\frac{1}{2}}). \end{aligned}$$

Part 2: By partial summation,

$$\begin{aligned}\psi_1(x) &= \sum_{n \leq x} \gamma(n) \ln(n) \\ &= \pi(x) \ln x - \int_2^x \frac{\pi(t)}{t} dt.\end{aligned}$$

Combining with (21.18) gives the result. □

Putting (21.17) into Lemma 21.4.2,

$$\begin{aligned}\pi(x) &= \frac{x}{\ln x} + O\left(\frac{xe^{-C\sqrt{\ln x}}}{\ln x}\right) + \int_2^x \left(\frac{1}{(\ln y)^2} + O\left(\frac{e^{-C\sqrt{\ln y}}}{(\ln y)^2}\right)\right) dy + O(x^{\frac{1}{2}}) \\ &= \text{li}(x) + O(xe^{-C\sqrt{\ln x}}).\end{aligned}$$

by (21.1)

5 The Riemann hypothesis

The following conjecture is worth one million dollars:

Conjecture 21.5.1 (Riemann hypothesis). *All nontrivial zeros s of $\zeta(s)$ satisfy $\Re s = \frac{1}{2}$.*

Note that for no $\varepsilon > 0$ has it been proved that all zeros satisfy $\Re s < 1 - \varepsilon$. Our zero-free region, sadly, has a boundary approaching real part 1 as $t \rightarrow \infty$.

One reason that the Riemann hypothesis is important is that it gives a strong error bound in the prime number theorem (as well as many other theorems of analytic number theory).

Theorem 21.5.2. *Suppose $\frac{1}{2} \leq \theta < 1$. The following are equivalent.*

1. $\zeta(s)$ has no zeros with $\Re s > \theta$.
2. $\pi(x) = \text{li}(x) + O(x^\theta \ln x)$.
3. $\pi(x) = \text{li}(x) + O(x^{\theta+\varepsilon})$ for every $\varepsilon > 0$, where the constant depends on ε .

In particular, the Riemann hypothesis is equivalent to $\pi(x) = \text{li}(x) + O(x^{\frac{1}{2}} \ln x)$.

Proof. (1) \implies (2): Suppose $\zeta(s)$ has no zeros with $\Re s > \theta$. Then using the estimate in (21.15), we have

$$\begin{aligned}\sum_{|\Im(\rho)| < T} \frac{x^\rho}{\rho} &\leq \max_{\rho} |x^\rho| \sum_{|\Im(\rho)| < T} \frac{1}{|\rho|} \\ &\leq x^\theta (\ln T)^2.\end{aligned}$$

Now take $T = x$ to find that

$$\begin{aligned} |\psi(x) - x| &= O\left(x^\theta(\ln T)^2 + \frac{x(\ln x)^2}{T}\right) \\ &= O(x^\theta(\ln x)^2). \end{aligned}$$

Then using Lemma 21.4.2 and (21.1),

$$\begin{aligned} \pi(x) &= \frac{\psi(x)}{\ln x} + \int_2^x \psi(y) \frac{dy}{y(\ln y)^2} + O(y^{\frac{1}{2}}) \\ &= \text{li}(x) + O\left(\frac{x^\theta(\ln x)^2}{\ln x}\right) + \int_2^x O\left(\frac{x^{\frac{1}{2}-1}(\ln x)^2}{(\ln x)^2}\right) dx \\ &= \text{li}(x) + O(x^\theta \ln x). \end{aligned}$$

(2) \implies (3): Item 2 is stronger than item 3.

(3) \implies (1): Going the other way in Lemma 21.4.2,

$$\begin{aligned} \psi(x) &= \pi(x) \ln x - \int_2^x \frac{\pi(y)}{y} dy + O(x^{\frac{1}{2}} \ln x) \\ &= \left(\frac{x}{\ln x} + \int_2^x \frac{dy}{(\ln y)^2} + O(x^{\theta+\varepsilon})\right) \ln x - \int_2^x \left(\frac{1}{\ln x} + \frac{1}{y} \int_2^y \frac{dt}{(\ln t)^2} + \frac{O(y^{\theta+\varepsilon})}{y}\right) dy + O(x^{\frac{1}{2}} \ln x) \\ &= x + O(x^{\theta+\varepsilon'}) - \underbrace{\int_2^x \frac{dy}{\ln y} + \int_2^x \frac{dy}{(\ln y)^2} \ln x - \int_2^x \left(\int_2^y \frac{dt}{(\ln t)^2} \cdot \frac{1}{y}\right) dy}_0 \end{aligned}$$

for any $\varepsilon' > \varepsilon$. Note the integrals above sum to 0 by integration by parts ($u = \ln y$, $dv = \frac{dy}{(\ln y)^2}$).

By partial summation, for $\sigma > 1$,

$$\begin{aligned} -\frac{\zeta'}{\zeta}(s) &= \sum_n \Lambda(n) n^{-s} \\ &= -\int_1^\infty \psi(n) s n^{-s-1} ds \\ &= \frac{s}{s-1} + s \int_1^\infty \underbrace{(\psi(x) - x)}_{O(x^{\theta+\varepsilon'})} x^{-s-1} dx. \end{aligned}$$

The last integral converges whenever $\sigma > \theta + \varepsilon'$, so $\frac{\zeta'}{\zeta}$ has analytic continuation to $\sigma > \theta$. This means ζ has no zeros for $\sigma > \theta$. \square

Chapter 22

L -functions and Dirichlet's theorem

1 Outline

Our goal in this chapter is to study the asymptotics of

$$\pi(x, a \bmod N) = |\{p \leq x : p \text{ prime, } p \equiv a \pmod{N}\}|$$

where a is relatively prime to N . We define $\psi(x, a \bmod N) = \sum_{n \leq x, n \equiv a \pmod{N}} \Lambda(n)$.

To study the distribution of primes in the arithmetic progression $n \equiv a \pmod{N}$, we study the asymptotics of $\psi(x, a \bmod N)$. However, this does not come from a Dirichlet series that we can easily estimate and that has nice multiplicative properties, like $\psi(x)$ comes from $\zeta(x) = \prod_p \frac{1}{1-p^{-s}}$ (after logarithmic differentiation and extracting coefficients).

The solution is to write $\psi(x, a \bmod N)$ in terms of Dirichlet series whose coefficients are multiplicative. For example, when considering primes $p \equiv 1 \pmod{4}$, we consider

$$\begin{aligned} L(s, \chi_1) &= \frac{1}{1^s} + \frac{1}{3^s} + \frac{1}{5^s} + \frac{1}{7^s} + \frac{1}{9^s} \cdots = \prod_p \frac{1}{1-p^{-s}}. \\ L(s, \chi_2) &= \frac{1}{1^s} - \frac{1}{3^s} + \frac{1}{5^s} - \frac{1}{7^s} + \frac{1}{9^s} \cdots = \prod_{p \equiv 1 \pmod{4}} \frac{1}{1-p^{-s}} \prod_{p \equiv 3 \pmod{4}} \frac{1}{1+p^{-s}} \end{aligned}$$

The multiplicative structure is from the fact that the coefficients come from group homomorphisms $(\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{C}$, i.e. Dirichlet characters (see Definition 6.1.8).

Logarithmic differentiation gives

$$\begin{aligned} -\frac{L'}{L}(s, \chi_1) &= \frac{\Lambda(1)}{1^s} + \frac{\Lambda(3)}{3^s} + \frac{\Lambda(5)}{5^s} + \frac{\Lambda(7)}{7^s} + \frac{\Lambda(9)}{9^s} \cdots \\ -\frac{L'}{L}(s, \chi_2) &= \frac{\Lambda(1)}{1^s} - \frac{\Lambda(3)}{3^s} + \frac{\Lambda(5)}{5^s} - \frac{\Lambda(7)}{7^s} + \frac{\Lambda(9)}{9^s} \cdots \\ \frac{1}{2} \left(-\frac{L'}{L}(s, \chi_1) - \frac{L'}{L}(s, \chi_2) \right) &= \frac{\Lambda(1)}{1^s} + \frac{\Lambda(5)}{5^s} + \frac{\Lambda(9)}{9^s} \cdots \end{aligned}$$

Taking the partial sum of coefficients of the last Dirichlet series gives the desired result. In general, we can always estimate $\psi(x, a \bmod N)$ using an average of these L -functions.

The main steps in the proof are the same, except with ζ replaced by L and an extra recombination step at the end using character theory. The main steps are the following.

1. Functional equation and analytic continuation for L , Theorem 22.2.5.
2. Product development, Theorem 22.2.6.
3. Estimates on $\frac{L'}{L}$ and asymptotics on number of zeros $N(T, \chi)$, Lemma 22.3.1.
4. Zero-free region for L , Theorem 22.3.3.
5. von Mangoldt's formula 22.4.1.

If we only cared about bounds for a fixed modulus N , then that's all there is to it.

However, to obtain error bounds independent of N , we need a zero free region independent of N (Theorem 22.3.3). While in Theorem 21.3.3 we had the luxury of restricting to large $|t|$, here we have to work with small $|t|$, and our resulting region may miss an “exceptional” zero. We show there is at most 1 exception (Theorem 22.4.2) and prove a version of the Prime Number Theorem for arithmetic progressions (Theorem 22.4.4). Later we prove a stronger but ineffective bound on the “exceptional zero” (Theorem 22.5.4) and obtain improved asymptotics (Theorem 22.5.1).

2 L -functions

Definition 22.2.1: Let χ be a Dirichlet character. Define the L function

$$L(s, \chi) := \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}, \quad \Re s > 1.$$

By multiplicativity of χ , L has a product expansion

$$L(s, \chi) = \prod_p \frac{1}{1 - \chi(p)p^{-s}}.$$

Only the factors with $p \nmid N$ contribute. Note that if χ is of level N and $\chi = \chi_1 \chi_2$ with χ_1 primitive of level N_1 , then

$$L(s, \chi) = L(s, \chi_1) \prod_{p|N, p \nmid N_1} (1 - \chi(p)p^{-s}). \quad (22.1)$$

Thus for convenience we can often just prove results about primitive characters.

By logarithmic differentiation we have

$$\frac{L'}{L}(s, \chi) = - \sum_p \frac{(\ln p)\chi(p)p^{-s}}{1 - p^{-s}} = - \sum_{n=1}^{\infty} \frac{\chi(n)\Lambda(n)}{n^s}.$$

Theorem 22.2.2 (Generalized Poisson summation). *Let g be a function $\mathbb{Z}/N\mathbb{Z} \rightarrow \mathbb{R}$, and suppose f is a C^2 function satisfying*

$$|f(x)|, |\hat{f}(x)| \leq C(1 + |x|)^{-1-\delta}$$

for some $C, \delta > 0$. Then

$$\sum_{m \in \mathbb{Z}} f\left(\frac{m}{N}\right) g(m) = \sum_{n \in \mathbb{Z}} \hat{f}(n) \hat{g}(n).$$

In particular, if χ is a primitive multiplicative character modulo N , then

$$\sum_{m \in \mathbb{Z}} \chi(m) f\left(\frac{m}{N}\right) = G(\chi, \chi_1^+) \sum_{n \in \mathbb{Z}} \bar{\chi}(-n) \hat{f}(n).$$

where $\chi_j^+(k) := e^{\frac{2\pi ijk}{N}}$.

Here $\hat{f}(n)$ denotes the Fourier transform

$$\hat{f}(y) = \int_{-\infty}^{\infty} f(x) e^{-2\pi ixy} dx$$

and $\hat{g}(n)$ denotes the finite Fourier transform

$$\hat{g}(n) = \sum_{m \pmod{N}} g(m) e^{-\frac{2\pi im}{N}}.$$

Proof. Consider the function

$$F(x) = \sum_{m \in \mathbb{Z}} f(x + m).$$

Note this sum converges absolutely to a continuous function by the given conditions. Since $F(x)$ has period 1 and is continuous, we can expand it in Fourier series:

$$F(x) = \sum_{n=0}^{\infty} a_n e^{2\pi inx},$$

$$a_n = \int_0^1 F(x) e^{-2\pi inx} dx = \int_0^1 \sum_{m \in \mathbb{Z}} f(x + m) e^{-2\pi inx} dx = \int_{-\infty}^{\infty} f(x) e^{-2\pi inx} dx = \hat{f}(n).$$

Plugging in $x = \frac{a}{N}$ gives

$$F\left(\frac{a}{N}\right) = \sum_{n \in \mathbb{Z}} \hat{f}(n) e^{2\pi in\left(\frac{a}{N}\right)}.$$

Now we calculate

$$\begin{aligned} \sum_{m \in \mathbb{Z}} f\left(\frac{m}{N}\right) g(m) &= \sum_{a \pmod{N}} g(a) F\left(\frac{a}{N}\right) \\ &= \sum_{a \pmod{N}} g(a) \sum_{n \in \mathbb{Z}} \hat{f}(n) e^{2\pi in\left(\frac{a}{N}\right)} \\ &= \sum_{n \in \mathbb{Z}} \hat{f}(n) \sum_{a \pmod{N}} g(a) e^{2\pi in\left(\frac{a}{N}\right)} \\ &= \sum_{n \in \mathbb{Z}} \hat{f}(n) \hat{g}(n). \end{aligned}$$

For the second part, note that

$$\begin{aligned} \sum_{m \in \mathbb{Z}} \chi(m) f\left(\frac{m}{N}\right) &= \sum_{n \in \mathbb{Z}} \hat{\chi}(n) \hat{f}(m) \\ &= \sum_{n \in \mathbb{Z}} G(\chi, \chi_1^+) \overline{\chi(n)} \hat{f}(n). \end{aligned} \quad \square$$

We apply Poisson summation to derive a transformation law for generalized theta functions.

Definition 22.2.3: Let χ be a multiplicative character modulo N . Define

$$\begin{aligned} \theta_\chi(u) &= \sum_{n \in \mathbb{Z}} \chi(n) e^{-\pi n^2 u} \\ \vartheta_\chi(u) &= \sum_{n \in \mathbb{Z}} \chi(n) n e^{-\pi n^2 u}. \end{aligned}$$

Note we need to work with $\vartheta_\chi(u)$ when χ is odd, since in this case $\theta_\chi(u) = 0$ and we cannot express $L(s, \chi)$ in terms of θ_χ .

Proposition 22.2.4 (Transformation law for θ_χ): Suppose χ is primitive. Then

$$\begin{aligned} \theta_\chi(u) &= \frac{G(\chi, \chi_1^+)}{N\sqrt{u}} \theta_{\bar{\chi}}\left(\frac{1}{N^2 u}\right) \\ \vartheta_\chi(u) &= -\frac{G(\chi, \chi^+)i}{N^2 u^{\frac{3}{2}}} \vartheta_{\bar{\chi}}\left(\frac{1}{N^2 u}\right). \end{aligned}$$

Proof. Note the Fourier transform of $e^{-\pi x^2}$ is itself; moreover, if $f(x) = g(ax)$ then $\hat{f}(y) = \hat{g}\left(\frac{y}{a}\right)$. Hence

$$\mathcal{F}(e^{-\pi u(Nx)^2}) = \frac{1}{N\sqrt{u}} e^{-\frac{\pi y^2}{uN^2}}.$$

By the Poisson summation formula 22.2.2,

$$\begin{aligned} \theta_\chi(u) &= \sum_{n \in \mathbb{Z}} \chi(n) e^{-\pi n^2 u} \\ &= \frac{G(\chi, \chi_1^+)}{N\sqrt{u}} \sum_{n \in \mathbb{Z}} \bar{\chi}(-n) e^{-\frac{\pi n^2}{uN^2}} \\ &= \frac{G(\chi, \chi_1^+)}{N\sqrt{u}} \theta_{\bar{\chi}}\left(\frac{1}{N^2 u}\right). \end{aligned}$$

For the second part, note first that $\hat{f}'(y) = 2\pi i y \hat{f}(y)$. Hence

$$\mathcal{F}(N x e^{-\pi u(Nx)^2}) = \left(-\frac{1}{2\pi u N}\right) \mathcal{F}\left(\frac{d}{dx}(x e^{-\pi u(Nx)^2})\right) = -\frac{1}{2\pi u N} \cdot 2\pi i y \frac{1}{N\sqrt{u}} e^{-\frac{\pi y^2}{uN^2}} = -\frac{i}{N^2 u^{\frac{3}{2}}} e^{-\frac{\pi y^2}{uN^2}}.$$

Then by Poisson summation,

$$\begin{aligned}
 \vartheta_\chi(u) &= \sum_{n \in \mathbb{Z}} \chi(n) n e^{-\pi n^2 u} \\
 &= -\frac{G(\chi, \chi_1^+) i}{N^2 u^{\frac{3}{2}}} \sum_{n \in \mathbb{Z}} \bar{\chi}(-n) n e^{-\frac{\pi n^2}{u N^2}} \\
 &= -\frac{G(\chi, \chi_1^+) i}{N^2 u^{\frac{3}{2}}} \vartheta_{\bar{\chi}}\left(\frac{1}{N^2 u}\right). \quad \square
 \end{aligned}$$

From this we get the functional equation for the L -function. The proof is similar to that of Theorem 21.2.2.

Theorem 22.2.5 (Analytic continuation and functional equation for L -functions). *Let χ be any character modulo N . Then $L(s, \chi)$ has a meromorphic continuation to \mathbb{C} . If χ is principal then $L(s, \chi)$ has a single pole at 1, and if χ is nonprincipal then $L(s, \chi)$ is entire.*

Now suppose χ is primitive. Defining

$$\xi(s, \chi) := \left(\frac{\pi}{N}\right)^{-\frac{s+a}{2}} \Gamma\left(\frac{s+a}{2}\right) L(s, \chi),$$

where

$$a = \begin{cases} 0, & \text{if } \chi(-1) = 1 \\ 1, & \text{if } \chi(-1) = -1, \end{cases}$$

we have

$$\xi(s, \chi) := \frac{G(\chi, \chi_1^+)}{i^a \sqrt{q}} \xi(1-s, \bar{\chi}).$$

Moreover, for any χ , $L(s, \chi)$ has zeros at $-2\mathbb{N} + a$ (the trivial zeros) and all other zeros are in the critical strip $0 \leq \Re s \leq 1$.

Note that for χ nonprincipal, partial cancellation in the Dirichlet series removes the pole at $s = 1$.

Proof. Note that it suffices to prove all statements for χ primitive, in light of (22.1). If χ is principal, the result follows from the result for ζ , so suppose χ is nonprincipal. Use partial summation ?? to find that for $s > 1$,

$$L(s, \chi) = \int_1^\infty S(x) s x^{-s-1} dx \quad (22.2)$$

where $S(x) = \sum_{n \leq x} \chi(n)$. (We use the fact that $\lim_{N \rightarrow \infty} S(N) N^{-s} = 0$ when $s > 1$.) Since $\chi(1) + \cdots + \chi(N) = 0$ by Corollary 6.1.7, $\chi(1) + \cdots + \chi(n) \leq N$. Then for $\Re s > 0$, the above integral converges absolutely, extending $L(s, \chi)$ holomorphically to $\Re s > 0$.

Case 1: Suppose $\chi(-1) = 1$; then $\chi(-n) = \chi(n)$. We calculate

$$\int_0^\infty \theta_\chi(u) u^{\frac{s}{2}} \frac{du}{u}$$

in two different ways.¹ When $0 < \Re s < 1$,

$$\begin{aligned}
 \int_0^\infty \theta_\chi(u) u^{\frac{s}{2}} \frac{du}{u} &= \int_0^\infty \sum_{n \in \mathbb{Z}} \chi(n) e^{-\pi n^2 u} u^{\frac{s}{2}} \frac{du}{u} \\
 &= 2 \sum_{n=1}^\infty \int_0^\infty \chi(n) e^{-\pi n^2 u} u^{\frac{s}{2}} \frac{du}{u} && \chi(-n) = \chi(n), \chi(0) = 0 \\
 &= 2 \sum_{n=1}^\infty \int_0^\infty \chi(n) e^{-u} \left(\frac{u}{\pi n^2} \right)^{\frac{s}{2}} \frac{du}{u} && u \leftarrow \frac{u}{\pi n^2} \\
 &= 2\pi^{-\frac{s}{2}} \left(\sum_{n=1}^\infty \frac{\chi(n)}{n^s} \right) \left(\int_0^\infty e^{-u} u^{\frac{s}{2}} \frac{du}{u} \right) \\
 &= 2\pi^{-\frac{s}{2}} L(s, \chi) \Gamma\left(\frac{s}{2}\right).
 \end{aligned}$$

Now using the transformation law 22.2.4,

$$\begin{aligned}
 \int_0^\infty \theta_\chi(u) u^{\frac{s}{2}} \frac{du}{u} &= \int_0^\infty \frac{G(\chi, \chi_1^+)}{N\sqrt{u}} \theta_{\bar{\chi}}\left(\frac{1}{N^2 u}\right) u^{\frac{s}{2}} \frac{du}{u} \\
 &= \frac{G(\chi, \chi_1^+)}{N} \int_0^\infty \theta_{\bar{\chi}}\left(\frac{1}{N^2 u}\right) u^{\frac{s}{2} - \frac{1}{2}} \frac{du}{u} \\
 &= \frac{2G(\chi, \chi_1^+)}{N} \sum_{n=1}^\infty \int_0^\infty \bar{\chi}(n) e^{-\frac{\pi n^2}{u N^2}} u^{\frac{s}{2} - \frac{1}{2}} \frac{du}{u} \\
 &= \frac{2G(\chi, \chi_1^+)}{N} \sum_{n=1}^\infty \int_0^\infty \bar{\chi}(n) e^{-u} \left(\frac{\pi n^2}{u N^2} \right)^{\frac{s}{2} - \frac{1}{2}} \frac{du}{u} && u \leftarrow \frac{\pi n^2}{u N^2} \\
 &= \frac{2G(\chi, \chi_1^+) \pi^{\frac{s}{2} - \frac{1}{2}}}{N^s} \sum_{n=1}^\infty \frac{\bar{\chi}(n)}{n^{(1-s)}} \int_0^\infty e^{-u} u^{\frac{1-s}{2}} \frac{du}{u} \\
 &= \frac{2G(\chi, \chi_1^+) \pi^{\frac{s}{2} - \frac{1}{2}}}{N^s} L(1-s, \bar{\chi}) \Gamma\left(\frac{1-s}{2}\right).
 \end{aligned}$$

Equating these two calculations gives the result.

Case 2: Suppose $\chi(-1) = -1$. We work with ϑ_χ instead of θ_χ . To compensate for the extra factor of n in ϑ_χ , we need an extra factor of $u^{\frac{1}{2}}$. We calculate

$$\int_0^\infty \vartheta_\chi(u) u^{\frac{s+1}{2}} \frac{du}{u}$$

¹Unlike in Theorem 21.2.2, there is no “-1” since $\chi(0) = 0$.

in two different ways. First,

$$\begin{aligned}
 \int_0^\infty \theta_\chi(u) u^{\frac{s+1}{2}} \frac{du}{u} &= \int_0^\infty \sum_{n \in \mathbb{Z}} \chi(n) n e^{-\pi n^2 u} u^{\frac{s+1}{2}} \frac{du}{u} \\
 &= 2 \sum_{n=1}^\infty \int_0^\infty \chi(n) n e^{-\pi n^2 u} u^{\frac{s+1}{2}} \frac{du}{u} && -n\chi(-n) = n\chi(n), \chi(0) = 0 \\
 &= 2 \sum_{n=1}^\infty \chi(n) n \int_0^\infty e^{-u} \left(\frac{u}{\pi n^2} \right)^{\frac{s+1}{2}} \frac{du}{u} && u \leftarrow \frac{u}{\pi n^2} \\
 &= 2\pi^{-\frac{s+1}{2}} \sum_{n=1}^\infty \frac{\chi(n)}{n^s} \int_0^\infty e^{-u} u^{\frac{s+1}{2}} \frac{du}{u} \\
 &= 2\pi^{-\frac{s+1}{2}} L(s, \chi) \Gamma\left(\frac{s+1}{2}\right).
 \end{aligned}$$

Now using the transformation law 22.2.4,

$$\begin{aligned}
 \int_0^\infty \theta_\chi(u) u^{\frac{s+1}{2}} \frac{du}{u} &= \int_0^\infty -\frac{G(\chi, \chi^+) i y}{N^2 u} \theta_{\bar{\chi}}\left(\frac{1}{N^2 u}\right) u^{\frac{s+1}{2}} \frac{du}{u} \\
 &= -\frac{G(\chi, \chi^+) i}{N^2} \int_0^\infty \theta_{\bar{\chi}}\left(\frac{1}{N^2 u}\right) u^{\frac{s}{2}-1} \frac{du}{u} \\
 &= -\frac{2G(\chi, \chi^+) i}{N^2} \sum_{n=1}^\infty n \bar{\chi}(n) \int_0^\infty e^{-\frac{\pi n^2}{u N^2}} u^{\frac{s}{2}-1} \frac{du}{u} \\
 &= -\frac{2G(\chi, \chi^+) i}{N^2} \sum_{n=1}^\infty \int_0^\infty \bar{\chi}(n) n e^{-u} \left(\frac{\pi n^2}{u N^2} \right)^{\frac{s}{2}-1} \frac{du}{u} && u \leftarrow \frac{\pi n^2}{u N^2} \\
 &= -\frac{2G(\chi, \chi^+) i \pi^{\frac{s}{2}-1}}{N^2 N^{n-2}} \sum_{n=1}^\infty \frac{\bar{\chi}(n)}{n^{1-s}} \int_0^\infty e^{-u} u^{1-\frac{s}{2}} \frac{du}{u} \\
 &= -\frac{2G(\chi, \chi^+) i \pi^{\frac{s}{2}-1}}{N^s} L(1-s, \bar{\chi}) \Gamma\left(1-\frac{s}{2}\right). \quad \square
 \end{aligned}$$

Again matching the two calculations gives the result.

From Proposition 19.7.2(5), Γ has no zeros, so we find that $L(s, \chi)$ is defined whenever $L(s, \bar{\chi})$ is defined; this L is entire. The description of the zeros of L follow from the functional equation and the fact that Γ has poles at $-\mathbb{N}_0$.

Theorem 22.2.6 (Product development of $\xi(s, \chi)$). *Suppose χ is primitive of level $N > 1$. The function $\xi(s, \chi)$ is entire of order 1 and has the product expansion*

$$\xi(s, \chi) = \xi(0, \chi) e^{Bs} \prod_{\rho \text{ zero of } \xi(s, \chi)} \left(1 - \frac{s}{\rho}\right) e^{\frac{s}{\rho}}.$$

Then $\frac{L'}{L}(s, \chi)$ has the partial-fraction expansion

$$\frac{L'}{L}(s, \chi) = B + \frac{1}{2} \ln\left(\frac{N}{\pi}\right) - \frac{1}{2} \frac{\Gamma'}{\Gamma}\left(\frac{s+a}{2}\right) + \sum_{\rho \text{ nontrivial zero of } \zeta} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right).$$

From now on, we only talk about nontrivial zeros of ζ .

Proof. We proceed as in Theorem 21.2.5. The argument is the same, the only major differences being that $\xi(s, \chi)$ has no poles at $s = 0, 1$, and the slight difference in definition of $\zeta(s, \chi)$ in terms of $L(s, \chi)$, versus the definition of $\xi(s)$ in terms of $\zeta(s)$. (Namely, we have $s + a$ instead of s , and an extra $N^{-\frac{s+a}{2}}$. For completeness we give the proof.

To show it has order 1 we need two inequalities.

Step 1: There is no constant C so that $\xi(s, \chi) \lesssim e^{C|s|}$: Indeed, for real s and any constant C' we have

$$\begin{aligned} \xi(s) &= \left(\frac{\pi}{N}\right)^{-\frac{s+a}{2}} \Gamma\left(\frac{s+a}{2}\right) L(s, \chi) \\ &\gtrsim s^{-\frac{1}{2}} \left(\frac{(s+a)N}{2e\pi}\right)^{\frac{s+a}{2}} \gtrsim e^{C's}. \end{aligned}$$

Step 2: There is a constant C so that $\xi(s, \chi) \lesssim e^{C|s|\ln|s|}$: $e^{|s|\ln|s|} \geq 1$ for all s so it suffices to prove this for sufficiently large s . By the integral and sum formulas for Γ and ξ , and the fact that $|x^s| = |x^{\Re s}|$, we have

$$|\xi(\sigma + ti, \chi)| \leq \left(\frac{\pi}{N}\right)^{-\frac{\sigma+a}{2}} \Gamma\left(\frac{\sigma+a}{2}\right) L(\sigma, \chi), \quad \sigma > 1.$$

By symmetry of ξ it suffices to consider $\sigma \geq \frac{1}{2}$. (We have $\xi(s, \chi) = \frac{G(\chi, \chi^+)}{i^a \sqrt{q}} \xi(1-s, \bar{\chi})$, and the multiplier has absolute value 1.) Consider 2 cases.

1. $\sigma > 2$: Then $\pi^{-\frac{\sigma+a}{2}} < 1$ and $L(\sigma, \chi) < \zeta(2)$ so we have by Stirling's approximation 19.7.4 that

$$|\xi(\sigma + ti, \chi)| \lesssim \left| N^{\frac{\sigma+a}{2}} \Gamma\left(\frac{\sigma + ti + a}{2}\right) \right| = N^{\frac{\sigma+a}{2}} e^{|\ln \Gamma(\sigma+a)|} = N^{\frac{\sigma+a}{2}} e^{\left(\frac{\sigma+a-1}{2}\right) \ln \frac{\sigma+a}{2} - \frac{\sigma+a}{2} + O(1)}$$

from which the result follows.

2. $\frac{1}{2} \leq \sigma \leq 2$: For s bounded away from 1, from (22.2),

$$L(s, \chi) = O(|s|).$$

This time $\Gamma\left(\frac{\sigma+a}{2}\right) = O(1)$ so

$$|L(s, \chi)| \leq \left| \left(\frac{\pi}{N}\right)^{-\frac{\sigma+a}{2}} L(s, \chi) \Gamma\left(\frac{\sigma+a}{2}\right) \right| = O(|s|) \lesssim e^{C|s|\ln|s|}.$$

This shows $\xi(s)$ has order 1.

Step 3: By the product development 19.6.3, noting the the zeros of $\xi(s, \chi)$ are the nontrivial zeros of $L(s, \chi)$, we get

$$\xi(s, \chi) = \xi(0, \chi)e^{Bs} \prod_{\rho \text{ zero of } L(s, \chi)} \left(1 - \frac{s}{\rho}\right) e^{\frac{s}{\rho}}.$$

Logarithmic differentiation gives

$$\frac{\xi'}{\xi}(s, \chi) = B + \sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{\rho}\right).$$

Since $L(s, \chi) = \left(\frac{\pi}{N}\right)^{\frac{s+a}{2}} \Gamma\left(\frac{s+a}{2}\right)^{-1} \xi(s, \chi)$, we get

$$\frac{L'}{L}(s, \chi) = \frac{1}{2} \ln\left(\frac{\pi}{N}\right) - \frac{1}{2} \frac{\Gamma'}{\Gamma}\left(\frac{s+a}{2}\right) + B + \sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{\rho}\right). \quad \square$$

3 Zeros of L

Lemma 22.3.1. *Define $\mathcal{L} = \ln N(|t| + 2)$. Let χ be a primitive character of level N . For $s = \sigma + it$ with $\sigma \in [-1, 2]$, we have*

$$\begin{aligned} \frac{L'}{L}(s, \chi) &= \sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{\rho}\right) + O(\mathcal{L}) \\ &= \sum_{|\Im(s-\rho)| < 1} \frac{1}{s - \rho} + O(\mathcal{L}). \end{aligned}$$

Moreover, there are $O(\ln |Nt|)$ zeros ρ with $|\Im(s - \rho)| < 1$, i.e. the number of zeros with imaginary part in $[t, t + 1]$ is $O(\ln Nt)$, as $t \rightarrow \infty$.

Note this gives $N(T) = O(T \ln(NT))$.

We follow the proof of Theorem 21.3.1.

Proof. The case $N = 1$ follows from there so we assume $N > 1$.

Step 1: Theorem 22.2.6 gives us

$$\frac{L'}{L}(s, \chi) = \underbrace{B + \frac{1}{2} \ln\left(\frac{N}{\pi}\right)}_{O(1+\ln N)} - \frac{1}{2} \underbrace{\frac{\Gamma'}{\Gamma}\left(\frac{s+a}{2}\right)}_{(A)} + \underbrace{\sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{\rho}\right)}_{(B)}. \quad (22.3)$$

From Stirling's approximation 19.7.4, (A) equals

$$\ln \left| \frac{\sigma + a}{2} + \frac{t}{2}i \right| + O(1) = O(\mathcal{L}). \quad (22.4)$$

Now suppose $s = 2 + it$. Note that

$$\left| \frac{L'}{L}(s, \chi) \right| = \left| \sum_{n=1}^{\infty} \chi(n) \Lambda(n) n^{-2-it} \right| \leq \left| \sum_{n=1}^{\infty} (\ln n) n^{-2} \right| < \infty,$$

so the LHS of (22.3) is $O(1)$. Hence (22.3) becomes

$$O(\mathcal{L}) = \sum_{\rho} \left(\frac{1}{s - \rho} + \frac{1}{\rho} \right). \tag{22.5}$$

Now finish the same way as in Theorem 1 to conclude the first step.

Step 2: Now we consider general $s = \sigma + it$, by comparing it to $2 + it$. We have by (22.3) and (22.4) that

$$\frac{L'}{L}(s, \chi) - \underbrace{\frac{L'}{L}(2 + it)}_{O(1)} = O(1) + \sum_{|\Im(s-\rho)| < 1} \frac{1}{s - \rho} + \underbrace{\sum_{|\Im(s-\rho)| < 1} \frac{1}{2 + it - \rho}}_{O(\mathcal{L})} + \underbrace{\sum_{|\Im(s-\rho)| \geq 1} \frac{(2 - \sigma) + it}{(s - \rho)(2 + it - \rho)}}_{O(\mathcal{L})}.$$

Finish as in Theorem 21.3.1, the only difference being that $\ln |t|$ is replaced by $\ln |Nt|$. \square

Theorem 22.3.2 (von Mangoldt). (*) As $T \rightarrow \infty$,

$$N(T, \chi) = \frac{T}{\pi} \ln \left(\frac{NT}{2\pi} \right) - \frac{T}{\pi} + O(\ln NT).$$

where the constant is independent of N .

Proof. The proof is similar to Theorem 1. We'll only need the weaker estimate $N(T, \chi) = O(T \ln NT)$ so we omit the proof. \square

Theorem 22.3.3 (Zero-free region for L). *There exists a constant $c > 0$, independent of χ and N , such that the following holds for all primitive χ of level N .*

1. If χ is nonreal, and $s = \sigma + it$ is a zero of $L(s, \chi)$, then

$$\sigma < 1 - \frac{c}{\mathcal{L}}. \tag{22.6}$$

2. If χ is real, then with at most 1 exception (counting multiplicity), all zeros satisfy (22.6). If it exists, the exceptional zero is real.

Unlike in Theorem 21.3.3, we have to worry about small $|t|$. Fortunately, $L(s, \chi)$ has no pole at $s = 1$ to screw us up. Things are not so easy, however.

Proof. We may assume $N \geq 2$.

As in Theorem 21.3.3, we have $0 \leq 3 + 4 \cos \theta + \cos 2\theta$, so

$$0 \leq 3 + 4\Re(\chi(n)n^{-it}) + \Re(\chi(n)^2n^{-2it}).$$

Multiplying by $\Lambda(n)n^{-\sigma}$ and summing, we get

$$0 \leq 3 \left(-\frac{L'}{L}(\sigma, \chi_0) \right) + 4\Re \left(-\frac{L'}{L}(\sigma + ti, \chi) \right) + \Re \left(-\frac{L'}{L}(\sigma + 2ti, \chi^2) \right), \quad \sigma > 1. \quad (22.7)$$

Suppose $1 < \sigma < 2$ and $\rho = (1 - \delta) + ti$ is zero. First we have

$$-\frac{L'}{L}(\sigma, \chi_0) = -\frac{\zeta'}{\zeta}(\sigma, \chi_0) - \sum_{p|N} \frac{(\ln p)p^{-s}}{1 - p^{-s}} = \frac{1}{\sigma - 1} + O(\ln N). \quad (22.8)$$

Next, we use the partial fraction decomposition 22.2.6. By Theorem 22.3.1 we have

$$\Re \left(-\frac{L'}{L}(s, \chi) \right) \leq O(\mathcal{L}) - \sum_{\rho} \Re \left(\frac{1}{s - \rho} \right). \quad (22.9)$$

1. Suppose χ^2 is not principal, i.e. χ is not real. Now (22.9) gives

$$\Re \left(-\frac{L'}{L}(\sigma + ti, \chi) \right) \leq O(\mathcal{L}) - \frac{1}{\sigma + \delta - 1}. \quad (22.10)$$

Also by Theorem 22.3.1

$$\Re \left(-\frac{L'}{L}(\sigma + 2ti, \chi^2) \right) \leq O(\mathcal{L}(2t)) = O(\mathcal{L}). \quad (22.11)$$

The remainder of this case follows the lines of Theorem 21.3.3.

2. If χ^2 is principal, then we have

$$\begin{aligned} -\frac{L'}{L}(\sigma + 2ti, \chi^2) &= -\frac{\zeta'}{\zeta}(\sigma + 2it) + \sum_{p|N} \ln p \cdot \underbrace{\frac{p^{-(\sigma+2ti)}}{1 - p^{-(\sigma+2ti)}}}_{O(1) \text{ when } \sigma \geq 1} \\ \Re \left(-\frac{L'}{L}(\sigma + 2ti, \chi^2) \right) &\leq O(\ln(|t| + 2)) + \Re \left(\frac{1}{(\sigma + 2ti) - 1} \right) + O(\ln N), \end{aligned} \quad (22.12)$$

the last inequality following from Lemma 21.3.1.

Putting (22.8), (22.9), and (22.12) into (22.11) give

$$0 \leq \left(\frac{3}{\sigma - 1} + O(\mathcal{L}) \right) + \left(-4 \sum_{\rho} \Re \left(\frac{1}{\sigma + ti - \rho} \right) + O(\mathcal{L}) \right) + \left(\Re \left(\frac{1}{\sigma + 2ti - 1} \right) + O(\mathcal{L}) \right) \quad (22.13)$$

Fix $C' > 0$; when $s = \sigma + it$ and $|t| \geq \frac{C'}{\ln N}$ then $\frac{1}{\sigma+2ti-1} = O(\ln N)$ so (22.11) holds and we proceed as in item 1.

Hence we consider $t < \frac{C'}{\ln N}$. We use a different approach. Note

$$-\frac{L'}{L}(\sigma, \chi_0) \geq \frac{L'}{L}(\sigma, \chi) \quad \text{when } \sigma \geq 1$$

because the coefficients their coefficients are $\Lambda(n) \geq -\chi(n)\Lambda(n)$ (and they are real)². Putting in (22.8) and (22.9) give

$$\frac{1}{\sigma-1} \geq \sum_{\rho} \Re\left(\frac{1}{\sigma-\rho}\right) + O(\ln N). \tag{22.14}$$

Let $\sigma = 1 + \frac{2\delta}{\ln N}$; we estimate the sum in terms of the real parts of $\sigma - \rho$. For any zero ρ we have

$$\begin{aligned} |\Im \rho| &\leq \frac{\delta}{\ln N} = \frac{1}{2} \frac{2\delta}{\ln N} \leq \Re(\sigma - \rho) \\ |\sigma - \rho|^2 &= [\Re(\sigma - \rho)]^2 + [\Im(\sigma - \rho)]^2 \end{aligned} \tag{22.15}$$

$$\leq \left(\frac{1}{4} + 1\right) \Re(\sigma - \rho)^2 = \frac{5}{4} \Re(\sigma - \rho)^2. \tag{22.16}$$

Hence (22.14) gives, for some constant A ,

$$\begin{aligned} \left(A + \frac{1}{2\delta}\right) \ln N &= \frac{1}{1-\sigma} + A \ln N \geq \sum_{|\Im(\rho)| < \frac{\delta}{\ln N}} \Re\left(\frac{1}{\sigma-\rho}\right) \\ &= \sum_{|\Im(\rho)| < \frac{\delta}{\ln N}} \frac{\Re(\sigma - \rho)}{|\sigma - \rho|^2} \\ &\geq \sum_{|\Im(\rho)| < \frac{\delta}{\ln N}} \frac{4}{5} \sum_{\rho} \frac{1}{\rho \left(1 + \frac{2\delta}{\ln N} - \Re(\rho)\right)} \quad \text{by (22.15)}. \end{aligned}$$

If $\Re(\rho) > 1 - \frac{c}{\ln N}$ then it contributes $\frac{4}{5} \frac{\ln N}{2\delta+c}$ to the RHS sum. If there are two zeros (counting multiplicity), then

$$\frac{8}{5} \frac{1}{2\delta+c} \leq A + \frac{1}{2\delta}.$$

This would be a contradiction if

$$c < \frac{2\delta(3-10A\delta)}{5(2\delta A+1)}.$$

Now choose δ small enough and c so that it works for case 1 and satisfies the above inequality.

Finally, note $\zeta(\bar{s}, \chi) = \overline{\zeta(s, \chi)}$ for real characters, so if s is an (exceptional) zero so is \bar{s} . Since there is at most one exceptional zero, it can only be real. \square

²Alternatively, put in $t = 0$ in (22.11).

4 Prime number theorem in arithmetic progressions

Theorem 22.4.1 (von Mangoldt’s formula). *For integer $x > 2$, $x \geq T$, and χ primitive of level $N > 1$,*

$$\psi(x, \chi) = - \sum_{|\Im(\rho)| < T} \frac{x^\rho}{\rho} + O\left(\frac{x[(\ln x)^2 + (\ln NT)^2]}{T}\right).$$

If χ has associated primitive character χ_1 , then for $x \geq 1$,

$$|\psi(x, \chi) - \psi(x, \chi_1)| = O(\ln N \ln x).$$

Note that unlike in Theorem 21.4.1, we have $\psi(x, \chi) \approx 0$ as opposed to $\psi(x) \approx x$. Remember this is expected because the average of values for a nontrivial character is 0, so there is cancellation in the sum. Moreover, there is no pole at $s = 1$ for L as there was in ζ , so the application of Cauchy’s Theorem in Step 2 will not give the x term.

Proof. Step 1: We estimate $\psi(x)$ using Theorem 20.4.2. Suppose x is an integer; the theorem gives

$$\begin{aligned} \left| \psi(x, \chi) - \left(\int_{c-iT}^{c+iT} x^s \left(-\frac{L'}{L}(s, \chi) \right) \frac{ds}{s} \right) \right| &\leq \Lambda(x) + \sum_{n \geq 1, n \neq x} \left(\frac{x}{n} \right)^c \chi(n) \Lambda(n) \frac{1}{T \left| \ln \left(\frac{x}{n} \right) \right|} \\ &\leq \ln(x) + \sum_{n \geq 1, n \neq x}^{\infty} \left(\frac{x}{n} \right)^c \frac{\ln(n)}{T \left| \ln \left(\frac{x}{n} \right) \right|}. \end{aligned}$$

The difference is $O\left(\frac{x(\ln x)^2}{T}\right)$ exactly as in (21.13).

Step 2: We move the line of integration to $\Re s = -1$. Assuming that T is not the imaginary part of any root, by Cauchy’s theorem

$$\begin{aligned} \int_{c-iT}^{c+iT} \frac{x^s}{s} \frac{L'}{L}(s, \chi) ds + \underbrace{\int_{c+iT}^{-1+iT} \frac{x^s}{s} \frac{L'}{L}(s, \chi) ds}_{I_{h,1}} + \underbrace{\int_{-1+iT}^{-1-iT} \frac{x^s}{s} \frac{L'}{L}(s, \chi) ds}_{I_v} + \underbrace{\int_{-1-iT}^{c-iT} \frac{x^s}{s} \frac{L'}{L}(s, \chi) ds}_{I_{h,2}} \\ = \frac{L'}{L}(0, \chi) - \sum_{|\Im \rho| < T} \frac{x^\rho}{\rho}. \end{aligned} \tag{22.17}$$

so

$$\int_{c-iT}^{c+iT} \frac{x^s}{s} \left(-\frac{L'}{L}(s) \right) ds = I_{h,1} + I_{h,2} + I_v + \frac{L'}{L}(0, \chi) - \sum_{\Im \rho < T} \frac{x^\rho}{\rho}. \tag{22.18}$$

We estimate each summand.

1. For the horizontal integrals, we use the estimate 22.3.1 to get

$$\begin{aligned} \left| \frac{\zeta'}{\zeta}(s) \right| &= \left| \sum_{|\Im(s-\rho)| < 1} \frac{1}{s-\rho} \right| + O(\ln NT), \quad s = \sigma + Ti \\ &\leq \sum_{|\Im(s-\rho)| < 1} \frac{1}{\Im(s-\rho)} + O(\ln NT). \end{aligned}$$

We would like to bound $\Im(s - \rho)$ away from 0. To do this, note that for $|T| > 2$ large there are $O(\ln NT)$ roots in with $\Im\rho \in \pm[T, T + 1]$ by Lemma 22.3.1. Hence by tweaking T slightly we can assume $|\Im(s - \rho)| > \frac{c}{\ln|NT|}$. Also by Lemma 22.3.1 there are at most $O(\ln NT)$ terms in the sum, so the sum is $O((\ln NT)^2)$. Integrating gives

$$\begin{aligned} \left| \int_{c \pm Ti}^{-1 \pm Ti} \frac{x^s}{s} \frac{L'}{L}(s, \chi) ds \right| &= O((\ln NT)^2) O\left(\frac{1}{T}\right) \int_c^{-1} |x^s| ds \\ &= O\left(\frac{x(\ln NT)^2}{T}\right). \end{aligned}$$

2. For the vertical integral, we use the same estimate, this time noting that $|s - \rho| > 1$ for every nontrivial zero ρ , since $\Re\rho > 0$. This gives that $\frac{\zeta'}{\zeta}(s) = O(\ln NT)$ and

$$\begin{aligned} \int_{-1+Ti}^{-1-Ti} \frac{x^s}{s} \frac{L'}{L}(s, \chi) ds &= O(\ln NT) \int_{-1-Ti}^{-1+Ti} \frac{x^{-1}}{|s|} ds \\ &= O\left(\frac{\ln(NT) \ln(T)}{x}\right) = O\left(\frac{x(\ln NT)^2}{T}\right). \end{aligned}$$

3. Note by Lemma 22.3.1 that $\frac{L'}{L}(0, \chi) = O(\mathcal{L}) = O(\ln(N + 1))$.

Step 1 and (22.18) together with the above estimates give the first part of the theorem.

For the second part, note that

$$\begin{aligned} \psi(x, \chi_1) - \psi(x, \chi) &= \sum_{1 \leq n \leq x} (\chi_1(n) - \chi(n)) \Lambda(n) n^{-s} \\ &\leq \sum_{1 \leq n \leq x, n=p^r, p|N} \Lambda(n) \\ &\leq \sum_{p|N} \left\lfloor \frac{\ln x}{\ln p} \right\rfloor \ln p \\ &\leq \sum_{p|N} \ln x \ln p = \ln x \ln N. \quad \square \end{aligned}$$

Theorem 22.4.2. *There is a constant $c > 0$ such that for any distinct real χ_1 and χ_2 to moduli N_1 and N_2 , at most one of $L(s, \chi_1)$ and $L(s, \chi_2)$ has a zero $\beta > 1 - \frac{c}{\ln(N_1 N_2)}$.*

Corollary 22.4.3. *There is a constant $c > 0$ such that the following holds: Fix a level N . There is at most 1 character χ of level N such that $L(s, \chi)$ has a zero with $\sigma \geq 1 - \frac{c}{\mathcal{L}}$.*

Proof of Theorem 22.4.2. The product $\chi_1 \chi_2$ is a character with modulus $N_1 N_2$. By Theorem 22.3.1, $-\frac{L'}{L}(\sigma, \chi) < O(\ln N_1 N_2)$ for $1 < \sigma < 2$. Let

$$F(s) = \zeta(s) L(s, \chi_1) L(s, \chi_2) L(s, \chi_1 \chi_2).$$

Then by logarithmic differentiation,

$$\begin{aligned}
 -\frac{F'}{F}(s) &= -\frac{\zeta'}{\zeta}(s) - \frac{L'}{L}(s, \chi_1) - \frac{L'}{L}(s, \chi_2) - \frac{L'}{L}(s, \chi_1\chi_2) \\
 &= \sum_{n=1}^{\infty} (1 + \chi_1(n) + \chi_2(n) + \chi_1(n)\chi_2(n))\Lambda(n)n^{-s} \\
 &= \sum_{n=1}^{\infty} (1 + \chi_1(n))(1 + \chi_2(n))\Lambda(n)n^{-s} \geq 0
 \end{aligned}
 \tag{22.19}$$

$$\tag{22.20}$$

since the coefficients are nonnegative.

Suppose β_1, β_2 are exceptional zeros of $L(s, \chi_1), L(s, \chi_2)$; then putting Lemma 21.3.1 into (22.20) gives

$$O(\ln N_1 N_2) + \frac{1}{\sigma - 1} - \frac{1}{\sigma - \beta_1} - \frac{1}{\sigma - \beta_2} \geq 0.$$

Let $\delta = \min(1 - \beta_1, 1 - \beta_2)$. Take $\sigma = 1 + 2\delta$ to get $\frac{1}{6\delta} \leq O(\ln N_1 N_2)$, i.e. $\delta \gtrsim \ln N_1 N_2$ with constant independent of N_1, N_2 , i.e. there is an appropriate choice of constant so that χ_1, χ_2 are not both exceptional for level $N_1 N_2$. \square

Proof of Corollary 5.2. Fix a primitive character χ of level N . Suppose χ' is of level N' , whose corresponding primitive characters has level N' . Then the theorem gives c such that at most one of $L(s, \chi')$ and $L(s, \chi)$ has a zero $\beta > 1 - \frac{c}{\ln N'N} \geq 1 - \frac{c}{\ln N}$. \square

Theorem 22.4.4 (Prime number theorem in arithmetical progressions). *Let $C > 0$ and suppose $x > e^{C(\ln N)^2}$. If there is no exceptional zero for level N , there exists $C' > 0$ such that*

$$\pi(x, a \bmod N) = (1 + O(e^{-C'\sqrt{\ln x}})) \frac{\text{li}(x)}{\varphi(N)}.$$

If there is an exceptional zero β of level N with associated character χ ,

$$\pi(x, a \bmod N) = \frac{1}{\varphi(N)} (\text{li}(x) - \chi(a) \text{li}(x^\beta) + O(xe^{-C'\sqrt{\ln x}})).$$

Proof. We have by column orthogonality 6.1.6 that

$$\psi(\chi, a \bmod N) = \sum_{n \leq x, n \equiv a \pmod{N}} \chi(n)\Lambda(N) = \sum_{n \leq x} \frac{1}{\varphi(N)} \sum_{\chi \in (\mathbb{Z}/N\mathbb{Z})^\times} \bar{\chi}(a)\chi(n)\Lambda(n) = \frac{1}{\varphi(N)} \sum_{\chi} \bar{\chi}(a)\psi(x, \chi).$$

$$\tag{22.21}$$

Letting χ_1 be the primitive character associated to χ , by Theorem 22.4.1 we have

$$\psi(x, \chi) = \begin{cases} -\sum_{\rho \text{ zero of } \psi(x, \chi_1)} \frac{x^\rho}{\rho} + O\left(\frac{x[(\ln x)^2 + (\ln NT)^2]}{T} + \ln N \ln x\right), & \chi \text{ nontrivial} \\ \psi(x) + O(\ln N \ln x), & \chi \text{ trivial.} \end{cases}$$

$$\tag{22.22}$$

We estimate $\sum_{\rho \text{ nonexceptional zero of } \psi(x, \chi_1)} \frac{x^\rho}{\rho}$ in two steps.³ Assume $T \geq 2$.

³Here “nonexceptional” means with respect to level N .

1. By Theorem 22.3.3, there is a constant c such that for all $|\Im(\rho)| < T$,

$$|x^\rho| = x^{\Re \rho} \leq x^{1 - \frac{c}{\ln NT}} = xe^{-\frac{c \ln x}{\ln NT}}$$

2. Note the zero free region in Theorem 22.3.3 means there is a constant d_0 , independent of N, χ , so that for all nonexceptional roots ρ , $|\rho| \geq d_0$. Hence using $N(T) = O(T \ln NT)$ (Lemma 21.3.1 or Theorem 22.3.2),

$$\begin{aligned} \sum_{|\Im(\rho)| < T} \frac{1}{|\rho|} &\leq \sum_{|\Im(\rho)| < T} \frac{1}{\max(\Im(\rho), d_0)} \\ &\leq \int_0^T \frac{dN(t)}{\max(t, d_0)} && \text{(Riemann-Stieltjes integral)} \\ &= \frac{N(T)}{\max(T, d_0)} + \int_{d_0}^T \frac{N(t)}{t^2} dt && \text{integration by parts} \\ &= O(\ln NT) + \int_{d_0}^T O\left(\frac{\ln Nt}{t}\right) dt \\ &= O(\ln NT) + O((\ln NT)^2) = O((\ln NT)^2). \end{aligned}$$

Putting these two estimates together,

$$\begin{aligned} \left| \sum_{|\Im(\rho)| < T, \rho \text{ nonexceptional}} \frac{x^\rho}{\rho} \right| &\leq \max_{|\Im(\rho)| < T} (|x^\rho|) \sum_{|\Im(\rho)| < T} \frac{1}{|\rho|} \\ &\leq O\left(e^{-\frac{c \ln x}{\ln NT}} (\ln NT)^2\right). \end{aligned}$$

Combining with Theorem 22.4.1, setting $T = e^{\sqrt{\ln x}}$, and using $N < e^{C\sqrt{\ln x}}$ we get

$$\begin{aligned} |\psi(x, \chi) - x| &= O\left(xe^{-\frac{c \ln x}{\ln NT}} (\ln NT)^2 + \frac{x[(\ln x)^2 + (\ln NT)^2]}{T} + \frac{x(\ln T)^2}{T}\right) - \frac{x^\beta}{\beta} - \frac{x^{1-\beta}}{1-\beta} \\ &= O\left(xe^{-\frac{c\sqrt{\ln x}}{C+1}} (C+1)^2 \ln x + xe^{-\sqrt{\ln x}}((\ln x)^2 + (C+1)^2 \ln x) + C(\ln x)^{\frac{3}{2}}\right) - \frac{x^\beta}{\beta} \\ &= O(xe^{-C_1\sqrt{\ln x}}) - \frac{x^\beta}{\beta} \end{aligned} \tag{22.23}$$

for some $C_1 > 0$ independent of N, χ , where the implied constant is independent of N, χ .

For the trivial character, (22.22) and (21.17) give

$$\psi(x, \chi) = x + O(xe^{-C_2\sqrt{\ln x}} + \ln x \ln T) = x + O(xe^{-C_2\sqrt{\ln x}}) \tag{22.24}$$

Using (22.21), (22.23), and (22.24), we get

$$\psi(\chi, a \bmod N) = \frac{1}{\varphi(N)} \left(x - \frac{\chi(a)x^\beta}{\beta} + O(xe^{-C_3\sqrt{\ln x}}) \right)$$

where the grayed-out portion appears only if there is an exceptional zero. (Note this can happen for at most 1 character by Lemma 22.4.2.) It remains to transfer the asymptotics of ψ to that for π .

The same argument as in Lemma 21.4.2 shows that

$$\pi(x, a \bmod N) = \frac{\psi(x, a \bmod N)}{\ln x} + \int_2^x \psi(y) \frac{dy}{y(\ln y)^2} + O(x^{\frac{1}{2}}),$$

giving the estimate for π . □

5 Siegel zero

In this section we obtain bounds on the exceptional zero to get a better error bound for prime number theorem on arithmetic progressions. We proceed in 2 steps.

1. Show that $L'(\beta, \chi)$ is small for β close to 1.
2. Bound $L(1, \chi)$ away from 0.

From this, we get that $L(\beta, \chi)$ cannot be 0 for β too close to 1.

Then we will be able to show the following improved form of Theorem 22.4.4.

Theorem 22.5.1 (Siegel-Walfisz). *Given any C there exists a constant C' depending only on C so that*

$$\pi(x, a \bmod N) = \frac{\text{li}(x)}{\varphi(N)} + O(xe^{-C'(\ln x)^{\frac{1}{2}}})$$

whenever

$$N \leq (\ln x)^C.$$

Unfortunately, this bound is *ineffective*; the proof does not give a way to compute a suitable value of C' .

Of course, if the Riemann hypothesis were true then it would solve all our problems.

Theorem 22.5.2. *If the Extended Riemann hypothesis holds (all nontrivial zeros of $L(s, \chi)$ satisfy $\Re s = \frac{1}{2}$), then*

$$\pi(x, a \bmod N) = \frac{\text{li}(x)}{\varphi(N)} + O(x^{\frac{1}{2}}(\ln x)^2)$$

for $x > N^2$, where the constant is independent of N .

5.1 $L'(\beta, \chi)$ is not too large

Lemma 22.5.3. *There exists an absolute constant C such that*

$$|L'(\sigma, \chi)| < C(\ln N)^2$$

for any nontrivial Dirichlet character χ modulo N and any σ with $1 - \frac{1}{\ln N} \leq \sigma \leq 1$.

Proof. Because $L(\sigma, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^\sigma}$, by Proposition 20.2.4 we can simply differentiate term-by-term to get

$$L'(\sigma, \chi) = - \sum_{n=1}^{\infty} \frac{\chi(n) \ln n}{n^\sigma}.$$

Now we bound this sum by breaking it up into two parts.

First note that for $n \leq N$, we have

$$1 - \sigma \leq \frac{1}{\ln N} \leq \frac{1}{\ln n}.$$

Hence

$$\frac{1}{n^\sigma} = \frac{1}{n} n^{1-\sigma} \leq \frac{1}{n} n^{\frac{1}{\ln n}} = \frac{e}{n}. \tag{22.25}$$

Step 1: We bound the sum from $n = 1$ to N . By (22.25),

$$\left| \sum_{n=1}^N \frac{\chi(n) \ln n}{n^\sigma} \right| \leq \sum_{n=1}^N \frac{e \ln n}{n} < C_1 (\ln N)^2 \tag{22.26}$$

for some C_1 . The last step follows from estimating using the integral $\int_1^N \frac{\ln x}{x} dx = \frac{1}{2} (\ln N)^2$.

Step 2: Now we consider the sum from $N + 1$ to ∞ . Let $U(n) := \sum_{m=L+1}^n \chi(m)$ and $v(n) = \frac{\ln n}{n^\sigma}$. By partial summation ??, we have

$$\sum_{n=N+1}^{\infty} \frac{\chi(n) \ln n}{n^\sigma} = \lim_{L \rightarrow \infty} \left[-U(L)v(L) + \sum_{n=N+1}^L U(n-1)(v(n) - v(n-1)) \right].$$

Since $v(n)$ decreases to 0 and $|U(n)| \leq N$ (as $\sum_{n=k}^{k+N-1} \chi(n) = 0$ for any k), the first term goes to 0 and we get the bound

$$\left| \sum_{n=N+1}^{\infty} \frac{\chi(n) \ln n}{n^\sigma} \right| \leq Nv(N) = N \frac{\ln N}{N^\sigma} \leq N (\ln N) \frac{e}{N} = e \ln N. \tag{22.27}$$

where in the last step we used (22.25).

Adding (22.26) and (22.27) together gives the desired bound. □

5.2 $L(1, \chi)$ is not too small

Theorem 22.5.4 (Siegel's inequality). *For each $\varepsilon > 0$ there exists $C_\varepsilon > 0$ such that*

$$L(1, \chi) > C_\varepsilon N^{-\varepsilon}$$

for all real Dirichlet characters χ modulo N .

Thus there exists $C'_\varepsilon > 0$ such that any real zero β of $L(s, \chi)$ satisfies $1 - \beta > C'_\varepsilon N^{-\varepsilon}$.

First we prove the following lemma.

Lemma 22.5.5. *Let χ_1 and χ_2 be real primitive characters with modulus N_1 and N_2 , let*

$$F(s) := \zeta(s)L(s, \chi_1)L(s, \chi_2)L(s, \chi_1\chi_2),$$

and let

$$\lambda = L(1, \chi_1)L(1, \chi_2)L(1, \chi_1\chi_2).$$

Then the following inequality holds:

$$F(s) > \frac{1}{2} - \frac{C\lambda}{1-s}(N_1N_2)^{8(1-s)}, \quad \frac{7}{8} < s < 1.$$

Note the technique of getting information about a L -function of a *single* character by looking at $F(s)$ —a function defined using *two* characters—is a lot like what we did in showing Corollary using Theorem 22.4.2. We’ll comment more later on why we looked at $F(s)$.⁴

Proof. The main idea is to expand $F(s)$ in power series and bound its coefficients (equivalently, bound the derivatives of $F(s)$) using the inequality from Cauchy’s formula, Corollary 19.4.6.

We have

$$\begin{aligned} \ln F(s) &= \ln \zeta(s) + \ln L(s, \chi_1) + \ln L(s, \chi_2) + \ln L(s, \chi_1\chi_2) \\ &= \sum_p \left(\ln \frac{1}{1-p^{-s}} + \ln \frac{1}{1-\chi_1(p)p^{-s}} + \ln \frac{1}{1-\chi_2(p)p^{-s}} + \ln \frac{1}{1-\chi_1(p)\chi_2(p)p^{-s}} \right) \\ &= \sum_p \sum_{m=1}^{\infty} \left(\frac{1}{m}p^{-ms} + \frac{1}{m}\chi_1(p^m)p^{-ms} + \frac{1}{m}\chi_2(p^m)p^{-ms} + \frac{1}{m}\chi_1(p^m)\chi_2(p^m)p^{-ms} \right) \\ &= \sum_p \sum_{m=1}^{\infty} \frac{1}{m}(1+\chi_1(p^m))(1+\chi_2(p^m))p^{-ms}. \end{aligned}$$

This means $\ln F(s)$ is a Dirichlet series with all coefficients positive. Because the power series of e^x has positive coefficients, this means that $F(s)$ also has all coefficients positive. Suppose $F(s) = \sum_{n=1}^{\infty} \frac{f(n)}{n^s}$.

Now we expand $F(s)$ in Taylor series at $s = 2$. (We can’t do it at $s = 1$ because $F(s)$ has a pole there.) We have

$$F(s) = \sum_{m=0}^{\infty} a_m(2-s)^m, \quad a_m = (-1)^m \frac{F^{(m)}(2)}{m!}.$$

⁴A deeper reason why we often look at $F(s)$ is that it is the zeta function of a *biquadratic field*. Thus we can prove nice facts about $F(s)$ by combining algebraic and analytic theory. We’ll give proofs that don’t require this knowledge.

We calculate the coefficients using 20.2.4 and get

$$a_m = \sum_{n=1}^{\infty} \frac{f(n)(\ln n)^m}{n^2} \geq 0.$$

In particular, for $m = 1$ we have $a_m \geq 1$ since $f(1) \geq 1$.

Because we know $F(s)$ has a pole of residue $\lambda = L(1, \chi_1)L(1, \chi_2)L(1, \chi_1\chi_2)$, we consider the function

$$F(s) - \frac{\lambda}{s-1} = F(s) - \frac{\lambda}{1-(2-s)} = \sum_{m=0}^{\infty} (a_m - \lambda)(2-s)^m.$$

Let Ω be the circle of radius $\frac{3}{2}$ (not its interior) centered at 2. Then for any χ of modulus N , $|L(s, \chi)| \leq C_1 N$ for some C_1 , for all s in a bounded region away from 0 because by (22.2)

$$|L(s, \chi)| = \left| \int_1^{\infty} S(x) s x^{-s-1} dx \right| \leq N \int_1^{\infty} |s x^{-s-1}| dx, \quad S(x) = \sum_{n \leq x} \chi(n).$$

Therefore,

$$|F(s)| \leq (C_1 N_1)(C_1 N_2)(C_1 N_1 N_2) = C_2 (N_1 N_2)^2, \quad C_2 = C_1^4 \quad (22.28)$$

and for $s \in \Omega$,

$$\left| \left(\frac{\lambda}{s-1} \right) \right| \leq 2L(1, \chi_1)L(1, \chi_2)L(1, \chi_1\chi_2) \leq 2C_2 (N_1 N_2)^2. \quad (22.29)$$

Now we use the inequality from Cauchy's formula, Corollary 19.4.6, to get

$$|a_m - \lambda| \leq \frac{1}{\left(\frac{3}{2}\right)^m} \max_{z \in \Omega} F(z) \leq C_3 N_1^2 N_2^2 \left(\frac{2}{3}\right)^m.$$

To bound $F(s) - \frac{\lambda}{s-1} = \sum_{m=0}^{\infty} (a_m - \lambda)(2-s)^m$ when $\frac{7}{8} < s < 1$, we first bound the sum from some M (to be determined) to ∞ .

Firstly,

$$\begin{aligned} \sum_{m=M}^{\infty} |a_m - \lambda|(2-s)^m &\leq \sum_{m=M}^{\infty} C_3 N_1^2 N_2^2 \left| \frac{2}{3}(2-s) \right|^m \\ &\leq \sum_{m=M}^{\infty} C_3 N_1^2 N_2^2 \left(\frac{3}{4}\right)^m, & \frac{7}{8} < s < 1 \\ &\leq C_4 N_1^2 N_2^2 \left(\frac{3}{4}\right)^M \\ &\leq C_4 N_1^2 N_2^2 e^{-M/4}, & e^{-1/4} \approx 0.78. \end{aligned}$$

We choose M so that $C_4 N_1^2 N_2^2 e^{-M/4} \in [\frac{1}{2}e^{-\frac{1}{4}}, \frac{1}{2}]$. Note the lower bound rearranges to $M \leq 8 \ln N_1 N_2 + C_5$. Then because the coefficients a_m are all nonnegative, we can drop some of them in the inequality to get

$$\begin{aligned}
 F(s) - \frac{\lambda}{s-1} &\geq 1 - \lambda \sum_{m=0}^{M-1} (2-s)^m - C_4 N_1^2 N_2^2 e^{-M/4} \\
 &> 1 - \frac{\lambda}{1-s} [(2-s)^M - 1] - \frac{1}{2}, & C_4 N_1^2 N_2^2 e^{-M/4} &\leq \frac{1}{2} \\
 \implies F(s) &> \frac{1}{2} - \frac{\lambda}{1-s} (2-s)^M \\
 &\geq \frac{1}{2} - \frac{\lambda}{1-s} e^{M(1-s)}, & e^x &\leq 1+x \\
 &> \frac{1}{2} - \frac{C_6 \lambda}{1-s} (N_1 N_2)^{8(1-s)}, & M &\leq 8 \ln N_1 N_2 + C_5.
 \end{aligned}$$

This finishes the proof of the lemma. □

Proof of Theorem 22.5.4. Fix $\varepsilon > 0$. We want to choose χ_1 so that $0 \geq F(s)$. Consider two cases.

1. For some χ , $L(s, \chi)$ has a real zero in the range $(1 - \frac{1}{16}\varepsilon, 1)$. Then choose χ_1 to be this character and β_1 to be this zero. We then have $F(\beta_1) = 0$.
2. Else, let χ_1 be any primitive character and $\beta_1 \in (1 - \frac{1}{16}\varepsilon, 1)$. Note the following:
 - In this case there are no zeros for any L-function in $(1 - \frac{1}{16}\varepsilon, 1)$, so they all have the same sign as their value at 1. The value at 1 is nonnegative (in fact, positive) because the product expansion gives that the L-function is positive for $\sigma > 1$.
 - $\zeta(s) < 0$ for $0 < s < 1$, and

Thus $F(\beta_1) < 0$.

In either case $F(\beta_1) \leq 0$, and the choice of β_1 depends only on ε . From Lemma 22.5.5, we now get the inequality

$$\begin{aligned}
 0 &\leq \frac{1}{2} - \frac{C\lambda}{1-\beta_1} (N_1 N_2)^{8(1-\beta_1)}. \\
 \lambda &> C_{\varepsilon,1} (N_1 N_2)^{-8(1-\beta_1)}
 \end{aligned}$$

for some $C_{\varepsilon,1}$ depending only on ε . Now we also have an upper bound for λ :

$$\begin{aligned}
 \lambda &= L(1, \chi_1)L(1, \chi_2)L(1, \chi_1\chi_2) \\
 &< (C_1 \ln N_1)L(1, \chi_2)(C_1 \ln N_1 N_2).
 \end{aligned}$$

Now suppose that $N_2 \geq N_1$. Combining the two inequalities and noting that $\ln N_1$ is a constant depending only on ε and is less than $\ln N_2$, we have

$$\begin{aligned} L(1, \chi_2) &> C_{\varepsilon,2} N_2^{-8(1-\beta_1)} (\ln N_2)^{-1} \\ &> C_{\varepsilon,2} N_2^{-\frac{\varepsilon}{2}} (\ln N_2)^{-1} \\ &> C_{\varepsilon,3} N^{-\varepsilon}. \end{aligned}$$

By choosing the constant to be smaller, we may ensure that this bound also works for $N_2 < N_1$.

Finally, combining Lemma 22.5.3 and the bound $L(1, \chi) > C_\varepsilon N^{-\varepsilon}$ immediately gives the fact that any real zero of $L(s, \chi)$ must satisfy $\beta < 1 - C'_\varepsilon N^{-\varepsilon}$. \square

Note that it was essential to work with $F(s)$ rather than $G(s) = \zeta(s)L(s, \chi)$: Something like Lemma 22.5.5 would go through, but if we used $G(s)$ then $G(s)$ may have a zero close to $s = 1$ so we don't know the region where $G(s)$ is nonpositive, and we may have to take $s = \beta_1$ arbitrarily close to 1. This kills the proof because of the term $\frac{1}{1-s}$. When we work with $F(s)$, the case where there is a zero close to 1 is dealt with nicely.

5.3 Proof of Siegel-Walfisz

Proof of Theorem 22.5.1. Suppose there is an exceptional zero β . By Siegel's inequality 22.5.4, for any $\varepsilon > 0$ we have

$$\beta - 1 < -C_\varepsilon N^{-\varepsilon}.$$

The prime number theorem in arithmetic progressions 22.4.4 gives

$$\pi(x, a \bmod N) = \frac{1}{\varphi(N)} (\text{li}(x) - \chi(a) \text{li}(x^\beta) + O(xe^{-C'\sqrt{\ln x}})).$$

We show that $\text{li}(x^\beta)$ gets absorbed into the O term. Indeed, we have

$$\begin{aligned} x^{-C_\varepsilon N^{-\varepsilon}} &\leq e^{-C'\sqrt{\ln x}} \\ \iff (\ln x) C_\varepsilon N^{-\varepsilon} &\geq C'\sqrt{\ln x} \\ \iff \sqrt{\ln x} &\geq \frac{C'}{C_\varepsilon} N^\varepsilon \\ \iff \left(\frac{C_\varepsilon}{C'}\right)^{\frac{1}{\varepsilon}} (\ln x)^{\frac{1}{2\varepsilon}} &\geq N. \end{aligned}$$

Now given $N \leq (\ln x)^C$, choose $\varepsilon = \frac{1}{2C}$. For large enough C' , the equivalences above give $x^{-C_\varepsilon N^{-\varepsilon}} \leq e^{-C'\sqrt{\ln x}}$. Therefore,

$$\text{li}(x^\beta) = O\left(x \frac{x^{\beta-1}}{\beta \ln x}\right) = O(x \cdot x^{-C_\varepsilon N^{-\varepsilon}}) = O(xe^{-C'\sqrt{\ln x}})$$

for some $C' > 0$, as needed. \square

Chapter 23

Special values of L -functions

0 Introduction

0.1 Points of interest

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} + \dots \quad (23.1)$$

$$\frac{\ln(\sqrt{2} + 1)}{\sqrt{2}} = 1 - \frac{1}{3} - \frac{1}{5} - \frac{1}{7} + \frac{1}{9} + \dots \quad (23.2)$$

$$\begin{aligned} \frac{2 \ln\left(\frac{1+\sqrt{5}}{2}\right)}{\sqrt{5}} &= 1 - \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \frac{1}{6} + \dots \quad (23.3) \\ &= \sum_{n=1}^{\infty} \frac{1}{5n+1} - \frac{1}{5n+2} - \frac{1}{5n+3} + \frac{1}{5n+4} \end{aligned}$$

See Mazur's article in PCTM [GBGL10]¹ for an introduction to algebraic number theory that highlights these formulas.

In this article we explore a formula that relates *analytic* with *algebraic* quantities: values of the zeta function with the class number, regulator, and discriminant of a number field. The main theorem is the following.

¹Available online at http://www.math.polytechnique.fr/~chenevier/MAT552/barry_mazur.pdf

Theorem 23.0.1 (Analytic class number formula). *Let K be a number field. Then*

$$\operatorname{Res}_{s=1} \zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} h_K \operatorname{Reg}_K}{\sqrt{|\Delta_K|} w_K} \quad (23.4)$$

where h_K is the class number, Reg_K is the regulator, w_K is the number of roots of unity in K , Δ_K is the discriminant, and $r_1, 2r_2$ are the number of real and complex embeddings of K .

0.2 Road map

Our plan is the following. First, we grok what's going on by looking at the class number formula for quadratic fields, which already show all the different types of behavior. To prove the theorem for general K we need to get reacquainted with the embedding $\sigma : \mathcal{O}_K \rightarrow \mathbb{R}^r \times \mathbb{C}^s$. Second, we'll massage our formula into nicer forms for quadratic fields and for cyclotomic fields, and see what algebraic information we can milk out from the class number formula.

How does this formula come about?² $\zeta_K(s)$ is a sum over ideals of \mathcal{O}_K where the ideals are weighted depending on their norm; we are “counting ideals” with appropriate weight. The growth of ζ_K near $s = 1$ depends on estimates for the number of ideals with norm $< r$. An expression for the *analytic* quantity $\operatorname{Res}_{s=1} \zeta_K(s)$ then comes from combining the following *geometric* and *algebraic* information.

1. Geometric: We count algebraic integers in K . Geometrically (using the embedding σ), they form a lattice in $\mathbb{R}^r \times \mathbb{C}^s$. We can estimate the number of points in a given region around the origin. This depends on the embedding, on the volume of a fundamental parallelootope of the lattice. This is how we get the quantity³

$$\frac{2^{r_1} (2\pi)^{r_2}}{\sqrt{|\Delta_K|}}.$$

2. Algebraic: Note $\zeta_K(s)$ is a sum over *ideals*, not over *numbers* in \mathcal{O}_K . So we need to multiply by a factor that tells us what we're off by in considering numbers rather than ideals; we can do this because of the multiplicative structure of \mathcal{O}_K . This will depend on the units (the fundamental units and the roots of unity) and the class number. This is how we get the factor

$$\frac{h_K \operatorname{Reg}_K}{w_K}.$$

²We follow the discussion in <http://math.stackexchange.com/questions/292104/how-to-derive-the-class-number-formula>.

³The grouping of the 2π is actually a little misleading: volume calculation gives the π^{r_2} (think circles); think of the 2^{r_1} as grouped with the $\sqrt{|\Delta_K|}$.

0.3 First explorations

Make sure you remember the following.

1. How to show $\text{Res}_{s=1} \zeta(s) = 1$. (We used a “summation by parts” argument; see (21.3). This was the argument that analytically continued $\zeta(s)$ from $\Re s > 1$ to $\Re s > 0$.)
2. The embeddings used to prove the finiteness of the class group and find the rank of the unit group. The definition of the regulator.

Problem 23.0.2: Recall that the zeta function for \mathbb{Q} can be defined as

$$\zeta(s) = \prod_{\text{prime } p} \left(1 - \frac{1}{p^s}\right) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

for $s > 1$. How do we modify this definition to define the zeta function for an arbitrary finite extension K/\mathbb{Q} ?

The identity above relied on unique factorization. In general we only have unique factorization of *ideals*, and we measure the size of an ideal with the *norm*. So we define

$$\zeta_K(s) = \sum_{\mathfrak{a} \in I_K} \frac{1}{\mathfrak{N}\mathfrak{a}^s}$$

and find by unique factorization of ideals that it equals⁴ $\prod_{\mathfrak{p} \subset \mathcal{O}_K \text{ prime}} \left(1 - \frac{1}{\mathfrak{N}\mathfrak{p}^s}\right)$.

First we explore the class number formula for quadratic number fields.

⁴We always restrict to nonzero ideals

Problem 23.0.3: 1. Consider the zeta function for $K = \mathbb{Q}(i)$. When we expand out $\zeta_K(s) = \sum_n \frac{a_n}{n^s}$, what do the coefficients a_n represent?

2. Evaluate $\text{Res}_{s=1} \zeta_K(s)$ (recall how we evaluated $\text{Res}_{s=1} \zeta(s)$), using some area calculations.
3. Can you write $\zeta_K(s)$ in terms of L -functions you are familiar with? Derive (23.1).
4. Do the same for $\mathbb{Z}[\sqrt{-2}]$, $\mathbb{Z}\left[\frac{1+\sqrt{-3}}{2}\right]$ and $\mathbb{Z}[\sqrt{-5}]$. What's different about the last case?
5. Do the same for $\mathbb{Z}[\sqrt{2}]$ and $\mathbb{Z}\left[\frac{1+\sqrt{5}}{2}\right]$. (The argument is more complicated, but make a guess.) Derive (23.2) and (23.3).
6. Conjecture the general class number formula (we already told you, but don't look back). Try to generalize your arguments for quadratic fields to prove it.
7. We saw that in (23.1) through (23.3) that we can evaluate in closed form series of the form $\sum \pm \frac{1}{n}$ (where the \pm are periodic). Can you do this in general?

Given a character $\chi : \mathbb{Z}/N \rightarrow \{-1, 1\}$, evaluate in closed form

$$\sum_{n=1}^{\infty} \frac{\chi(n)}{n}$$

in 2 ways: algebraically, as in (23.1) through (23.3), and analytically (without mentioning algebraic quantities). Try to simplify your formula as much as possible. What happens when you equate these two?

Hint for the analytic expression: (a) power series are nice, (b) $\sum \frac{1}{n}$ reminds us of logarithms. However we seem to have nasty sums such as $\sum_{n \equiv m \pmod{p}} \frac{1}{n^s}$. What's the technique?

For a leisurely account of the class number formula for quadratic fields (assuming few prerequisites) see the PROMYS notes [Wei].⁵ We will discuss 1–5 in Section 1, 6 in Section 1.1, and 7 in Section 2.

⁵Available at <http://math.uchicago.edu/~chonoles/expository-notes/promys/promys2012-analyticclassnumberformulanotes.pdf>

1 The analytic class number formula

1.1 Quadratic fields

This section will be mostly to build intuition for the general case; we will give a self-contained proof of the general case in the next section.

1. We have $\zeta_K(s) = \sum_{a \in I_K} \frac{1}{\mathfrak{N}a^s} = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$ where

$$a_n = \text{the number of ideals of norm } n.$$

Now we specialize to the case $K = \mathbb{Q}(i)$. Every ideal is principal, of the form $(a + bi)$, and its norm is $a^2 + b^2$. So does a_n count the number of solutions to $a^2 + b^2 = n$? Almost: multiplying by a root of unity gives the same ideal. $(a + bi)$, $(i(a + bi))$, $(i^2(a + bi))$, $(i^3(a + bi))$ represent the same ideal, so a_n is $\frac{1}{4}$ the number of solutions.

We can think of this another way: each ideal is of the form $(a + bi)$ for a unique pair $(a, b) \in \mathbb{Z}^2$ with $a > 0$ and $b \geq 0$ (“in the first quadrant”). Hence a_n counts the number of solutions

$$a_n = |\{(a, b) \in \mathbb{Z}^2, a > 0, b \geq 0 : a^2 + b^2 = n\}|.$$

2. We don't have a nice expression for a_n that we can work with analytically, but we can work with $\sum_{k=1}^n a_k$: it counts the number of solutions

$$\sum_{k=1}^n a_k = |\{(a, b) \in \mathbb{Z}^2, a > 0, b \geq 0 : a^2 + b^2 \leq n\}|,$$

i.e., the points inside the circle of radius \sqrt{n} in the first quadrant.

This suggests that we use *summation by parts* on ζ_K : (this is valid as $(n+1)^{-s} \sum_{k=1}^n a_k$ converges to 0 by the estimate below, (23.5))

$$\begin{aligned} \zeta_K(s) &= \sum_n a_n n^{-s} \\ &= \sum_{n=1}^{\infty} \left(\sum_{k=1}^n a_k \right) (n^{-s} - (n+1)^{-s}) \\ &= \sum_{n=1}^{\infty} \left(\sum_{k=1}^n a_k \right) s \int_n^{n+1} x^{-s-1} dx. \end{aligned}$$

Now we need to estimate $\sum_{k=1}^n a_k$. It is approximately the area of the circle of radius \sqrt{n} in the first quadrant, so

$$\sum_{k=1}^n a_k = \frac{\pi}{4}n + e_n \text{ where } e_n = O(\sqrt{n}). \quad (23.5)$$

(The standard argument is to consider a square for each lattice point; these squares exactly cover the sector except possibly for a strip near the boundary of area proportional to the circumference, $\sim \sqrt{n}$.) Putting this in, we have for $s \geq 1$,

$$\begin{aligned}
 \zeta_K(s) &= s \sum_{n=1}^{\infty} \left(\frac{\pi}{4}n + e_n \right) \int_n^{n+1} x^{-s-1} dx & (23.6) \\
 &\text{writing } n = (n-x) + x, \\
 &= s \sum_{n=1}^{\infty} \frac{\pi}{4} \left(\int_n^{n+1} x^{-s} dx + \int_n^{n+1} x^{-s-1}(n-x) dx + e_n \int_n^{n+1} x^{-s-1} dx \right) dx \\
 &= \frac{\pi}{4} s \int_1^{\infty} x^{-s} dx + \frac{\pi}{4} s \sum_{n=1}^{\infty} \left(\int_n^{n+1} x^{-s-1}(x-n) dx + e_n \int_n^{n+1} x^{-s-1} dx \right) \\
 &= \frac{\pi}{4} \frac{s}{s-1} + \frac{\pi}{4} s \underbrace{\sum_{n=1}^{\infty} \left(\int_n^{n+1} x^{-s-1}(x-n) dx + e_n \int_n^{n+1} x^{-s-1} dx \right)}_{O(n^{-s-\frac{1}{2}})}
 \end{aligned}$$

since $e_n = O(n^{\frac{1}{2}})$. The last sum converges for $\Re s > \frac{1}{2}$ to an analytic function, so we obtain an analytic continuation for ζ_K to $\Re s > \frac{1}{2}$. (We have $\Re s > \frac{1}{2}$ rather than $\Re s > 0$ as for ζ because of the error term.) Breaking up $\frac{s}{s-1} = 1 + \frac{1}{s-1}$ and reading off the coefficient of $\frac{1}{s-1}$, we see

$$\boxed{\text{Res}_{s=1} \zeta_K(s) = \frac{\pi}{4}}.$$

Summarizing, the π comes from an area calculation, and the 4 comes from the fact there are 4 roots of unity in $\mathbb{Q}(i)$.

3. We'd like to write $\zeta_K(s) = \prod_{\mathfrak{p}} \left(1 - \frac{1}{\mathfrak{p}^s} \right)$ in terms of primes over p . In a quadratic field, a prime p in \mathbb{Z} either

- (a) is inert in K , i.e., is a prime of norm p^2 in K
- (b) splits into 2 primes of norm p , or
- (c) ramifies.

For a field of discriminant D , these cases happen when $\left(\frac{D}{p}\right) = 1$, $\left(\frac{D}{p}\right) = -1$, and $p \mid D$, respectively. We get

$$\begin{aligned}
 \zeta_K(s) &= \prod_{p \text{ inert}} \frac{1}{1-p^{-2s}} \prod_{p \text{ splits}} \left(\frac{1}{1-p^{-s}} \right)^2 \prod_{p \text{ ramifies}} \left(\frac{1}{1-p^{-s}} \right) \\
 &= \prod_p \frac{1}{1-p^{-s}} \prod_{p \text{ inert}} \left(\frac{1}{1+p^{-s}} \right) \prod_{p \text{ splits}} \left(\frac{1}{1-p^{-s}} \right) \\
 &= \zeta(s) \prod_{p \nmid D} \left(\frac{1}{1 - \left(\frac{D}{p}\right) p^{-s}} \right)
 \end{aligned}$$

$$\zeta_K(s) = \zeta(s)L(s, \chi), \quad \chi = \left(\frac{D}{p}\right). \tag{23.7}$$

(The L function encodes the splitting law for primes in K .)⁶ Consider $K = \mathbb{Q}(i)$. Here

$$\chi(n) = \begin{cases} 1, & n \equiv 1 \pmod{4} \\ -1, & n \equiv 3 \pmod{4}. \end{cases} \text{ Now we know } \text{Res}_1(\zeta_K) = 1, \text{ so we must have}$$

$$\boxed{\frac{\pi}{4} = L(s, \chi) = \sum_n \frac{\chi(n)}{n^s} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots}$$

Incidentally, note that the identity (23.7) gives (via expanding the Dirichlet series) the formula

$$|\{(a, b) \in \mathbb{Z}^2 : a^2 + b^2 = n\}| = 4(d_1(n) - d_3(n))$$

where $d_j(n)$ is the number of divisors of n that are $\equiv j \pmod{4}$.

4. For $\sqrt{-2}$, what changes in the above calculations?
 - (a) We now count instead the solutions to $x^2 + 2y^2 \leq n$. Alternatively, keeping the shape the same, we're counting the lattice points inside the circle where the lattice is generated by 1 and $\sqrt{2}i$ (each rectangle has area $\sqrt{2}$; in general it has area $\sqrt{\frac{|\Delta_K|}{4}}$). The number of points is approximately $\sqrt{2}\pi n$.
 - (b) There are only 2 roots of unity, so we would count points in the whole upper part of the circle/ellipse, and divide by 2 instead of 4.

⁶We note that χ is a homomorphism modulo D . I show this using a case analysis (not sure if there is a cleaner way...). In the following we use the notation (a if P) to mean a if P is true and 1 otherwise. First note that D is squarefree except possibly for 4 (and either $2^0, 2^2$, or $2^3 \parallel D$); by multiplicativity of $\left(\frac{\bullet}{p}\right)$ and quadratic reciprocity,

$$\begin{aligned} \left(\frac{D}{p}\right) &= \left(\left(\frac{-1}{p}\right) \text{ if } D < 0\right) \left(\frac{2}{p}\right)^{v_p(D)} \left(\prod_{q|D, q \neq 2} \left(\frac{p}{q}\right) (-1)^{\frac{p-1}{2} \frac{q-1}{2}}\right) \\ &= \left((-1)^{\frac{p-1}{2}} \text{ if } D < 0\right) \left((-1)^{\frac{p^2-1}{8}} \text{ if } 8 \mid D\right) \cdot \prod_{q|D, q \neq 2} \left(\frac{p}{q}\right) \cdot (-1)^{\frac{p-1}{2} |\{p=4k+1 \mid D\}|} \end{aligned}$$

so

- (a) if $8 \mid D$, then χ is a primitive character modulo $8 \prod_{p|D, p \neq 2} p = D$,
- (b) if $4 \mid D$, so that D is divisible by an odd number of $4k + 3$ primes (so that $(-1)^{\frac{p-1}{2} |\{p=4k+1 \mid D\}|} = (-1)^{\frac{p-1}{2}}$) and $D > 0$, or D is divisible by an even number of $4k + 3$ primes and $D < 0$, then χ is a primitive character modulo $4 \prod_{p|D, p \neq 2} p = D$,
- (c) if $4 \nmid D$, so that D is divisible by an even number of $4k + 3$ primes and $D > 0$, or D is divisible by an odd number of $4k + 3$ primes and $D < 0$, then χ is a primitive character modulo $\prod_{p|D, p \neq 2} p = D$. To see this in the second case, note we have the factors $(-1)^{\frac{p-1}{2}} (-1)^{\frac{p-1}{2}} = 1$.

Similarly for $\sqrt{-3}$ we get $\frac{\sqrt{3}}{2}\pi n$ and we divide by 6. We get

$$\begin{aligned}\operatorname{Res}_{\mathbb{Q}(\sqrt{2}i)} \zeta(s) &= \frac{\pi}{\sqrt{22}} \\ &= 1 - \frac{1}{3} - \frac{1}{5} + \frac{1}{7} + \frac{1}{9} - \frac{1}{11} - \frac{1}{13} + \frac{1}{15} + \dots \\ \operatorname{Res}_{\mathbb{Q}(\sqrt{3}i)} \zeta(s) &= \frac{\pi}{\frac{\sqrt{3}}{2} \cdot 6} \\ &= 1 - \frac{1}{5} + \frac{1}{7} - \frac{1}{11} + \frac{1}{13} - \frac{1}{17} + \dots\end{aligned}$$

For $\sqrt{-5}$, the problem is that the class number is > 1 : a prime can now split into nonprincipal ideals, so we can't simply count the number of solutions to $x^2 + 5y^2 = 1$. We'll do this rigorously later, but it makes sense that since there are 2 ideal classes, we'll only capture half the sum in this way, and we'll have to multiply by 2.

In general, if K is a imaginary quadratic field with discriminant D_K , then

$$L(1, \chi) = \operatorname{Res}_{s=1} \zeta_K(s) = \frac{2\pi h_K}{\sqrt{|D_K|} w_K}. \quad (23.8)$$

where $\chi(j) = \left(\frac{D_K}{p}\right)$, for any prime $p \equiv j \pmod{D_K}$. [Here $D_K < 0$.]

5. Recall the embedding we use in order to apply geometry to $K = \mathbb{Q}(\sqrt{d})$ with d positive: $x + y\sqrt{d} \mapsto (x + y\sqrt{d}, x - y\sqrt{d}) =: (x', y')$. The area of the fundamental parallelogram of this lattice L is $\sqrt{|\Delta_K|}$.

The norm is different now; it is $x'y' = x^2 - d^2y$. Again we have a factor h_K . We now count lattice points under the hyperbola $x'y' < n$ in the first and second quadrant. However, now associates can differ not just by a root of unity (-1), but also by a unit, which must be some power ε^n of the fundamental unit ε (which is sent to the lattice point $(\varepsilon, \frac{1}{\varepsilon})$). Drawing a picture, we see that each $x + y\sqrt{d}$ with norm at most n has an associate in the region bounded by $x'y' = n$ and the lines joining the origin and $(1, 1)$, and the origin and $(\varepsilon, \frac{1}{\varepsilon})$, or the rotation of this region by 90° . The area of these slices can be calculated to be $2 \ln \varepsilon$. Thus we expect around $\frac{2 \ln \varepsilon}{\sqrt{|\Delta_K|}}$ lattice points. We obtain the following:

If d is a real quadratic field and ε a fundamental unit, then (note $w_K = 2$)

$$L(1, \chi) = \text{Res}_{s=1} \zeta_K(s) = \frac{2h_K \ln \varepsilon}{\sqrt{|D_K|}} \quad (23.9)$$

where $\chi(j) = \left(\frac{D_K}{p}\right)$, for any prime $p \equiv j \pmod{D_K}$. [Here $D_K > 0$.]

In the case of $\sqrt{5}$, the discriminant is 5, the character is $\chi(n) = \left(\frac{n}{5}\right)$, and the fundamental unit is $\frac{1+\sqrt{5}}{2}$, so

$$1 - \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \frac{1}{6} - \cdots = \text{Res}_{s=1} \zeta_{\mathbb{Q}(\sqrt{5})}(s) = \frac{2 \ln \left(\frac{1+\sqrt{5}}{2}\right)}{\sqrt{5}}.$$

1.2 Proof in general

We now prove Theorem 23.0.1 in 4 steps. Note that both (23.8) and (23.9) are special cases of (23.4): in the first case there is 1 pair of complex embeddings, hence the factor of π ; in the second there are 2 real embeddings, hence the factor of 2. The group of units of a real quadratic field is 1-dimensional, so the regulator is simply the logarithm of the fundamental unit.

In general the area of a fundamental parallelopete is $\sqrt{|\Delta_K|} 2^{-r_2}$ (Proposition 3.3.1).

1.2.1 Reduce to sum over principal ideals

We reduce $\zeta_K(s)$ to a sum over numbers in \mathcal{O}_K , not ideals. We separate the sum by ideal classes:

$$\begin{aligned} \zeta_K(s) &= \sum_{\mathfrak{a}} \frac{1}{\mathfrak{N}\mathfrak{a}^s} \\ &= \sum_{[I] \in \text{Cl}_K} \sum_{\mathfrak{a} \in [I]} \frac{1}{\mathfrak{N}\mathfrak{a}^s} \\ &\quad \text{where the sum } [I] \in \text{Cl}_K \text{ is over ideal classes.} \\ &\quad \text{Now for each } [I] \text{ choose } \mathfrak{b}_I \in [I]^{-1} \\ &= \sum_{[I] \in \text{Cl}_K} \sum_{\mathfrak{a} \in [I]} \frac{1}{\mathfrak{N}(\mathfrak{a}\mathfrak{b}_I)^s} \mathfrak{N}(\mathfrak{b}_I)^s \\ &= \sum_{[I] \in \text{Cl}_K} \sum_{(\alpha), \alpha \in \mathfrak{b}_I} \frac{1}{\mathfrak{N}(\alpha)^s} \mathfrak{N}(\mathfrak{b}_I)^s \end{aligned} \quad (23.10)$$

since $\mathfrak{a} \mapsto \mathfrak{a}\mathfrak{b}_I$ is a bijection from the ideals in $[I]$ to principal ideals that are divisible by \mathfrak{b} , i.e., principal ideals in the form (α) , $\alpha \in \mathfrak{b}$. If we manage to show each of the inner sums is $\frac{2^{r_1}(2\pi)^{r_2} \text{Reg}_K}{\sqrt{|\Delta_K|} w_K}$, then we'll be done.

1.2.2 Reduce to sum over numbers

The sums are now over principal ideals, but we'd like them to be over elements. The obstruction is that many elements will give the same ideal, so we'd like a canonical way of choosing a generator of (α) . We'd like a region such that every set of associated elements has exactly one element in that region, a kind of *fundamental domain*.

Any two generators will differ by a product $\zeta \varepsilon_1^{a_1} \cdots \varepsilon_{r+s-1}^{a_{r+s-1}}$ where ζ is a root of unity and $\varepsilon_1, \dots, \varepsilon_{r+s-1}$ are the fundamental units.

If we have a vector space, and we say 2 elements are equivalent if they differ by an element of a lattice, we know what a fundamental domain would look like—a parallelotope. We have here a product rather than a \mathbb{Z} -linear combination, so we'll have to embed and then take logs.

Recall the embedding and the log map from Proposition 5.2.2:

$$L : \mathcal{O}_K^\times \xrightarrow{\sigma=(\sigma_1, \dots, \sigma_{r_1+r_2})} \mathbb{R}^r \times \mathbb{C}^s \xrightarrow{\ell=(\ln|\cdot|, \dots, \ln|\cdot|, \ln 2|\cdot|, \dots, \ln 2|\cdot|)} \mathbb{R}^{r_1+r_2}.$$

(The kernel of this map is the roots of unity.) Recall that $L(U_K)$ is the hyperplane H with $x_1 + \cdots + x_{r_1+r_2} = 0$, i.e. the hyperplane with $\mathbf{x} \cdot (1, \dots, 1) = 0$. Given a principal ideal we choose a generator α such that the projection of $L(\alpha)$ onto this subspace is in the fundamental parallelotope made by the $L(\varepsilon_1), \dots, L(\varepsilon_{r+s-1})$, i.e. we require

$$L(\alpha) \in \underbrace{([0, 1]L(\varepsilon_1) + \cdots + [0, 1]L(\varepsilon_{r_1+r_2-1}))}_{=:P} + \mathbb{R}(1, \dots, 1) \tag{23.11}$$

(For a set S of scalars and a vector v we let $Sv = \{sv : s \in S\}$.) Taking the inverse image under L and noting the kernel is the roots of unity (Proposition 5.2.2), we have the following.

- For any $\alpha \in \mathcal{O}_K$, there are exactly w_K associates β of α such that⁷

$$L(\beta) \in P + \mathbb{R}(1, \dots, 1).$$

Combining with (23.10),

$$\zeta_K(s) = \frac{1}{w_K} \sum_{[I] \in \text{Cl}_K} \mathfrak{N}(\mathfrak{b}_I)^s \sum_{\alpha \in L^{-1}(P + \mathbb{R}(1, \dots, 1)) \cap \mathfrak{b}_I} \frac{1}{\mathfrak{N}(\alpha)^s}. \tag{23.12}$$

1.2.3 Reduce to a volume calculation

Now we show how the sums in (23.12) reduce to a volume calculation. We claim the following.

⁷I.e., α is unique up to multiplication by a root of unity. To make α unique, we can require that α have argument in $[0, \frac{2\pi}{w_K})$.

Lemma 23.1.1. *Let L be a lattice in \mathbb{R}^n , and let $\Delta(L)$ denote the volume of its fundamental parallelotope.⁸ Let V be a bounded, measurable set containing the origin that the points $B(\partial V, c)$ at a distance of at most c from the boundary satisfy*

$$\text{Vol}[B(\partial V, c)] \lesssim_c t^{n-1}.$$

(There aren't too many points near the surface.)⁹ Then

1. The number of points inside tV satisfies

$$|tV \cap L| = \frac{\text{Vol}(V)}{\Delta(L)} t^m + O(t^{m-1})$$

2. Suppose $[0, 1]V = V$. Let $N(x) = (\inf \{c > 0 : x \in cV\})^m$ (the norm associated to V , made homogeneous of degree m). The function

$$f(s) = \sum_{x \in L} \frac{1}{N(x)^s}$$

is meromorphic for $\Re s \geq 1 - \frac{1}{m}$ with $\text{Res}_{s=1} f(s) = \frac{\text{Vol}(V)}{\Delta(L)}$.

Proof. 1. Take a fundamental parallelotope and place one at each point in $tV \cap L$; now bound the difference between this area and the area of tV , cf. (23.5).

2. This is exactly as in (23.6). Noting that the number of $x \in L$ with $N(x) \leq n$ is $\frac{\text{Vol}(V)}{\Delta(L)} n + e_n$ for some $e_n = O(n^{\frac{m-1}{m}})$,

$$\begin{aligned} f(s) &= s \sum_{n=1}^{\infty} \left(\frac{\text{Vol}(V)}{\Delta(L)} n + e_n \right) \int_n^{n+1} x^{-s-1} dx \\ &\quad \text{writing } n = (n-x) + x, \\ &= s \sum_{n=1}^{\infty} \frac{\text{Vol}(V)}{\Delta(L)} \left(\int_n^{n+1} x^{-s} dx + \int_n^{n+1} x^{-s-1}(x-n) dx + e_n \int_n^{n+1} x^{-s-1} dx \right) dx \\ &= s \frac{\text{Vol}(V)}{\Delta(L)} \int_1^{\infty} x^{-s} dx + \frac{\text{Vol}(V)}{\Delta(L)} s \sum_{n=1}^{\infty} \left(\int_n^{n+1} x^{-s-1}(x-n) dx + e_n \int_n^{n+1} x^{-s-1} dx \right) \\ &= \frac{\text{Vol}(V)}{\Delta(L)} \frac{s}{s-1} + \frac{\text{Vol}(V)}{\Delta(L)} s \underbrace{\sum_{n=1}^{\infty} \left(\int_n^{n+1} x^{-s-1}(x-n) dx + e_n \int_n^{n+1} x^{-s-1} dx \right)}_{O(n^{-s-\frac{1}{m}})} \end{aligned}$$

since $e_n = O(n^{\frac{m-1}{m}})$. The last sum converges for $\Re s > 1 - \frac{1}{m}$ to an analytic function, so we obtain an analytic continuation for ζ_K to $\Re s > 1 - \frac{1}{m}$. (We have $\Re s > 1 - \frac{1}{m}$ rather than $\Re s > 0$ for ζ because of the error term.) Reading off the coefficient of $\frac{1}{s-1}$, we see

$$\text{Res}_{s=1} f(s) = \frac{\text{Vol}(V)}{\Delta(L)}. \quad \square$$

⁸Distinguish this from Δ_K ; the relationship between Δ_K and $\Delta(L)$ is given by $\Delta(L) = \sqrt{|\Delta_K|} 2^{-r_2}$; see Proposition 3.3.1.

⁹See also VI.§2 in Lang [?]. His Theorem 2 is phrased in terms of $(n-1)$ -Lipschitz parametrizable boundaries.

1.2.4 Calculate the volume

We will apply Lemma 23.1.1 to

$$V = \ell^{-1}(P - [0, \infty)(1, \dots, 1)) = [0, 1]\ell^{-1}(P), \quad L = \mathfrak{b}_I$$

where P is as in (23.11). The norm on V is exactly the norm we want: $N(\mathbf{x}) = |x_1 \cdots x_{r_1+r_2}|$ on $[0, \infty)V \subseteq \mathbb{R}^{r_1} \times \mathbb{C}^{r_2}$.




Figure 23.1: The light blue and gray regions comprise $[0, 1]\ell^{-1}(P)$. The light blue region is $[0, 1]e^P$. (In this example, P is 1-dimensional.)

We calculate the volume of V : We want to evaluate (note $x_i \in \mathbb{R}$ or \mathbb{C})

$$\begin{aligned} \text{Vol}(V) &= \int \cdots \int_{[0,1]\ell^{-1}(P) \subseteq \mathbb{R}^{r_1} \times \mathbb{C}^{r_2}} dx_1 \cdots dx_{r_1+r_2} \\ &= \int \cdots \int_{[0,1] \cdot \{-1,1\}, \dots, \{-1,1\}, D, \dots, D} \cdot e^{\text{diag}(1, \dots, 1, \frac{1}{2}, \dots, \frac{1}{2})P} \subseteq \mathbb{R}^{r_1} \times \mathbb{C}^{r_2} dx_1 \cdots dx_{r_1+r_2} \\ &= \int \cdots \int_{[0,1]e^{\text{diag}(1, \dots, 1, \frac{1}{2}, \dots, \frac{1}{2})P} \subseteq \mathbb{R}^{r_1+r_2}} 2^{r_1} (2\pi)^{r_2} x_{r_1+1} \cdots x_{r_1+r_2} dx_1 \cdots dx_{r_1+r_2} \end{aligned}$$

where

- $\text{diag}(1, \dots, 1, 2, \dots, 2)$ is the map sending

$$(x_1, \dots, x_{r_1}, x_{r_1+1}, \dots, x_{r_1+r_2}) \mapsto (x_1, \dots, x_{r_1}, 2x_{r_1+1}, \dots, 2x_{r_1+r_2}),$$

- $D = \{z \in \mathbb{C} : |z| = 1\}$.
- multiplication and exponentiation is coordinatewise.

When we take the inverse image $\ell^{-1}(P)$, we not only get $[0, 1]e^{\text{diag}(1, \dots, 1, \frac{1}{2}, \dots, \frac{1}{2})P}$; the real coordinates also have the freedom of being positive or negative and the complex coordinates also have the freedom of varying by $|z| = 1$. We know the area of P , so we change coordinates to make the integral over P . Change coordinates via the map ℓ (so $x_i =$

$\begin{cases} e^{x'_i}, & 1 \leq i \leq r_1 \\ e^{x'_i/2}, & r_1 < i \leq r_1 + r_2 \end{cases}$ which has determinant of Jacobian $2^{-r_2} e^{x'_1} \dots e^{x'_r} e^{x'_{r+1}/2} \dots e^{x'_{r_1+r_2}/2}$
to get

$$\begin{aligned} & \int \dots \int_{P-[0,\infty)(1,\dots,1)} 2^{r_1} (2\pi)^{r_2} e^{x'_{r+1}/2} \dots e^{x'_s/2} (2^{-r_2} e^{x'_1} \dots e^{x'_{r_1+r_2}/2} dx'_1 \dots dx'_{r_1+r_2}) \\ &= 2^{r_1} \pi^{r_2} \int \dots \int_{P-[0,\infty)(1,\dots,1)} e^{x'_1} \dots e^{x'_{r_1+r_2}} dx'_1 \dots dx'_s \\ & \text{setting } y = \frac{1}{\sqrt{r_1+r_2}}(x'_1 + \dots + x'_{r_1+r_2}); \text{ the integral } \int_P \text{ in the hyperplane of } P \\ &= 2^{r_1} \pi^{r_2} \int_P \int_{-\infty}^0 e^{y\sqrt{r_1+r_2}} dy d\mathbf{x} \\ &= 2^{r_1} \pi^{r_2} \int_P \frac{1}{\sqrt{r_1+r_2}} d\mathbf{x} \\ &= 2^{r_1} \pi^{r_2} \text{Vol}(P) \frac{1}{\sqrt{r_1+r_2}} \\ &= 2^{r_1} \pi^{r_2} \text{Reg}_K \end{aligned}$$

where the last line follows from Proposition 23.4.1. Now each sum in (23.12) has, by Lemma 23.1.1, residue equal to

$$\begin{aligned} \text{Res}_{s=1} \sum_{\alpha \in \mathbb{R}\ell^{-1}(P) \cap \mathfrak{b}_I} \frac{1}{\mathfrak{N}(\alpha)^s} &= \frac{\text{Vol}(V)}{\Delta(\sigma(\mathfrak{b}_I))} \\ &= \frac{2^{r_1} \pi^{r_2} \text{Reg}_K}{\sqrt{|\Delta_K|} 2^{-r_2} \mathfrak{N}(\mathfrak{b}_I)} \text{ by Prop. 3.3.1} \\ \implies \text{Res}_{s=1} \zeta_K(s) &= \frac{1}{w_K} \sum_{[I] \in \text{Cl}_K} \mathfrak{N}\mathfrak{b}_I \left(\text{Res}_{s=1} \sum_{\alpha \in \mathbb{R}\ell^{-1}(P) \cap \mathfrak{b}_I} \frac{1}{\mathfrak{N}(\alpha)^s} \right) \\ &= \frac{1}{w_K} \sum_{[I] \in \text{Cl}_K} \mathfrak{N}\mathfrak{b}_I \frac{2^{r_1} \pi^{r_2} \text{Reg}_K}{\sqrt{|\Delta_K|} 2^{-r_2} \mathfrak{N}(\mathfrak{b}_I)} \\ &= \frac{2^{r_1} (2\pi)^{r_2} \text{Reg}_K}{w_K \sqrt{|\Delta_K|}}. \end{aligned}$$

This finishes the proof of Theorem 23.0.1.

2 Special values of L -functions

2.1 Evaluating $L(1, \chi)$ for quadratic characters

We were able to find closed forms for the sums (23.1)–(23.3). The same method works in general.

We now evaluate $L(1, \chi)$ for quadratic characters in 2 ways.

1. First, we found that we have an algebraic expression for it. If $\chi(p) = \left(\frac{D}{p}\right)$, then letting K be the field with discriminant D , we can use the formula for $\text{Res}_{s=1} \zeta_K(s)$, either (23.8) or (23.9).¹⁰
2. Second, we find an closed form for it analytically (Theorem 2.1).

Equating these two will give us a way to calculate the class number analytically. There are 4 cases, depending on whether D is positive or negative, and whether χ is odd or even. We won't write these all out, but note when $D = \pm p$, in 2 of the cases, the identity will give us information on quadratic residues and nonresidues.

Theorem 23.2.1 (Formulas for L -functions). *Let χ be a character modulo m . Let $\tau(\chi)$ denote the Gauss sum $\sum_{j \in (\mathbb{Z}/m)} \chi(j)\chi^+(j)$ where $\chi^+(j) = \zeta^j$ is the additive character. We have*

$$L(1, \chi) = -\frac{\tau(\chi)}{m} \sum_{k \in (\mathbb{Z}/m)^\times} \overline{\chi(k)} \ln \sin \frac{\pi k}{m}, \quad \chi \text{ even} \quad (23.13)$$

$$L(1, \chi) = \frac{\pi i \tau(\chi)}{m^2} \sum_{k \in (\mathbb{Z}/m)^\times} \overline{\chi(k)} k, \quad \chi \text{ odd.} \quad (23.14)$$

(In [BS66], this is Theorem 3, p. 336 and Theorem 1, p. 344.)

Proof. We write $L(1, \chi)$ as a sum over residues modulo n , and then write the nasty sum $\sum_{n \equiv m} \frac{1}{n^s}$ (a partial zeta function) in terms of nice sums using characters. We have

$$\begin{aligned} L(1, \chi) &= \sum_{n=1}^{\infty} \chi(n) \frac{1}{n} = \sum_{k \in \mathbb{Z}/m} \chi(k) \sum_{\substack{n \equiv k \\ (\text{mod } D)}} \frac{1}{n} \\ &= \sum_{k \in \mathbb{Z}/m} \chi(k) \sum_{j \in \mathbb{Z}/m} \frac{1}{m} \sum_n \zeta^{(n-k)j} \frac{1}{n} \quad \text{orthogonality of characters} \\ &= \frac{1}{m} \sum_{k \in \mathbb{Z}/m} \sum_{j \in \mathbb{Z}/m} \chi(k) \zeta^{-jk} \sum_{n=1}^{\infty} \frac{\zeta^{nj}}{n}. \end{aligned}$$

The inner sum is just the Taylor series of a logarithm. We have¹¹

$$\sum_{n=1}^{\infty} \frac{\zeta^{nj}}{n} = -\ln(1 - \zeta^j).$$

¹⁰Every quadratic character is in this form. To use a sledgehammer, this follows by class field theory since quadratic characters are in bijection with the nontrivial elements of Galois groups of quadratic extensions. From first principles, note that χ is a continuous homomorphism on $\hat{\mathbb{Z}} = \prod_p \mathbb{Z}_p$, and for $p \neq 2$, $1 + p\mathbb{Z}_p \subseteq \mathbb{Z}_p^2$; for $p = 2$, $1 + 8\mathbb{Z}_2 \subseteq \mathbb{Z}_2^2$, so χ must be 1 on these subgroups. Thus χ is a primitive character modulo D for some $D \mid 8 \prod_{p \neq 2} p$, and corresponds to the field with discriminant D or $-D$.

¹¹We use the fact that if $\sum_{n=1}^{\infty} e^{i\theta n} a_n$ converges, then $\lim_{r \rightarrow 1^-} r^n e^{i\theta n} a_n$ equals that sum.

Letting $\chi_j^+(n) = \zeta^{nj}$, we find the sum equals (using Proposition 6.2.2)

$$\begin{aligned} L(1, \chi) &= -\frac{1}{m} \sum_{j \in \mathbb{Z}/m} \sum_{k \in \mathbb{Z}/m} \chi(k) \zeta^{-jk} \ln(1 - \zeta^j) \\ &= -\frac{1}{m} \sum_{j \in \mathbb{Z}/m} G(\chi, \chi_{-j}^+) \ln(1 - \zeta^j) \\ &= -\frac{\tau(\chi)}{m} \sum_{j \in (\mathbb{Z}/m)^\times} \overline{\chi(-j)} \ln(1 - \zeta^j) \end{aligned}$$

where the Gauss sum is 0 for $j \not\equiv 0 \pmod{m}$. (Proof: let j' be smallest nonzero so that $jj' \equiv 0 \pmod{m}$. We can group the terms in the sum as $\sum_{k=0}^{j'-1} (\chi(k) + \chi(k+j') + \cdots + \chi(k+p-j')) \zeta^{-jk}$. Now note $\chi(k) + \chi(k+j') + \cdots + \chi(k+p-j') = \chi(k)(\chi(1) + \chi(1+j') + \cdots + \chi(1+p-j'))$, and because χ is primitive, it is nontrivial on the subgroup $(1+j'\mathbb{Z})/m\mathbb{Z}$, so $\chi(1) + \chi(1+j') + \cdots + \chi(1+p-j') = 0$.) We want to express this sum in terms of \ln of real values. We use a standard trick: match up j and $-j$ in the sum. We consider two cases.

1. χ is even, i.e., $\chi(j) = \chi(-j)$. Then the sum equals

$$\begin{aligned} -\frac{\tau(\chi)}{m} \frac{1}{2} \sum_{j \in (\mathbb{Z}/m)^\times} \overline{\chi(-j)} (\ln(1 - \zeta^j) + \ln(1 - \zeta^{-j})) &= -\frac{\tau(\chi)}{m} \frac{1}{2} \sum_{j \in (\mathbb{Z}/m)^\times} \overline{\chi(j)} \ln((\zeta^{\frac{j}{2}} - \zeta^{-\frac{j}{2}})^2) \\ &= -\frac{\tau(\chi)}{m} \sum_{j \in (\mathbb{Z}/m)^\times} \overline{\chi(j)} \ln \left(2 \sin \frac{\pi j}{m} \right) \\ &= -\frac{\tau(\chi)}{m} \sum_{j \in (\mathbb{Z}/m)^\times} \overline{\chi(j)} \ln \left(\sin \frac{\pi j}{m} \right) \end{aligned}$$

where the 2 came out as a $\ln 2$ and we use $\sum_{j \in (\mathbb{Z}/m)^\times} \overline{\chi(j)} = 0$.

2. χ is odd, i.e., $\chi(-j) = -\chi(j)$. Then the sum equals

$$\begin{aligned} -\tau(\chi) \frac{1}{2} \sum_{j \in (\mathbb{Z}/m)^\times} \overline{\chi(-j)} (\ln(1 - \zeta^j) - \ln(1 - \zeta^{-j})) &= -\frac{\tau(\chi)}{m} \frac{1}{2} \sum_{j \in (\mathbb{Z}/m)^\times} \overline{-\chi(j)} (\ln(\zeta^j)) \\ &= \frac{\tau(\chi)}{m} \frac{1}{2} \sum_{j \in (\mathbb{Z}/m)^\times} \overline{\chi(j)} \left(\frac{2\pi i j}{m} \right) \\ &= \frac{\pi i \tau(\chi)}{m^2} \sum_{j \in (\mathbb{Z}/m)^\times} \overline{\chi(j)} j. \quad \square \end{aligned}$$

2.2 The class number counts quadratic residues/nonresidues

Note that Theorem 2.1 can give information on quadratic residues modulo p when $\chi(k) = \left(\frac{k}{p}\right)$. When does this happen? We note

$$\chi_{\mathbb{Q}(\sqrt{p})}(q) = \left(\frac{p}{q}\right) = \left(\frac{q}{p}\right) (-1)^{\frac{p-1}{2} \frac{q-1}{2}} = \left(\frac{q}{p}\right) \text{ when } p \equiv 1 \pmod{4}. \quad (23.15)$$

$$\chi_{\mathbb{Q}(\sqrt{-p})}(q) = \left(\frac{-p}{q}\right) = (-1)^{\frac{q-1}{2}} \left(\frac{p}{q}\right) = (-1)^{\frac{q-1}{2}} \left(\frac{q}{p}\right) (-1)^{\frac{p-1}{2} \frac{q-1}{2}} \quad (23.16)$$

$$= \left(\frac{q}{p}\right) (-1)^{\frac{p+1}{2} \frac{q-1}{2}} \text{ when } p \equiv 3 \pmod{4}. \quad (23.17)$$

Hence when $p \equiv 1 \pmod{4}$ we consider $\mathbb{Q}(\sqrt{p})$, and when $p \equiv 3 \pmod{4}$ we consider $\mathbb{Q}(\sqrt{-p})$.

2.2.1 $p \equiv 1 \pmod{4}$

In the case $p \equiv 1 \pmod{4}$, we find an identity involving the fundamental unit, the class number, and quadratic residues/nonresidues. There is no explicit formula for the fundamental unit, but we can find an explicit formula that gives a *power* of the fundamental unit. The class number appears as the exponent.

Theorem 23.2.2. *Let K be a real quadratic field with discriminant D and character χ . Then*

$$\eta := \prod_{a \perp D, 0 < a < \frac{D}{2}} \left(\sin \frac{\pi a}{D} \right)^{-\chi(a)}$$

is a unit and if $\varepsilon > 1$ is the fundamental unit,

$$\varepsilon^h = \eta.$$

In particular, if $p \equiv 1 \pmod{4}$, then $\prod_{\left(\frac{b}{p}\right)=-1} \sin \frac{\pi b}{p} > \prod_{\left(\frac{a}{p}\right)=1} \sin \frac{\pi a}{p}$.

For $p \equiv 1 \pmod{4}$, think of the last statement as saying that quadratic residues mod p cluster at the beginning of $(0, p/2)$ (where $\sin \frac{\pi x}{p}$ is small) and the quadratic residues mod p cluster at the end.

Proof. Note χ is even. Let $h = h_K$. We equate (23.9) with (23.13) to get

$$\begin{aligned} \frac{2h \ln \varepsilon}{\sqrt{D}} &= L(1, \chi) = -\frac{\tau(\chi)}{D} \sum_{k \in (\mathbb{Z}/D)^\times} \overline{\chi(k)} \ln \sin \frac{\pi k}{D}. \\ &= -\frac{1}{\sqrt{D}} \sum_{k \in (\mathbb{Z}/D)^\times} \overline{\chi(k)} \ln \sin \frac{\pi k}{D}. \end{aligned}$$

where, because χ is even, by Theorem 23.4.2, $\tau(\chi) = \sqrt{D}$. We get (matching up $\chi(x) = \chi(-x)$)

$$\begin{aligned}
 (\ln \varepsilon)h &= \sum_{x \perp D, 0 < x < \frac{D}{2}} \ln \sin \frac{\pi x}{D} (-\chi(x)) \\
 \varepsilon^h &= \prod_{a \perp D, 0 < a < \frac{D}{2}} \left(\sin \frac{\pi a}{D} \right)^{-\chi(a)}. \quad \square
 \end{aligned}$$

2.2.2 $p \equiv 3 \pmod{4}$

In the imaginary quadratic case, we have an explicit formula for the class number that depends just on calculating quadratic residues, and not on anything analytic (ln, sin,...).

Theorem 23.2.3. *Let $p \equiv 3 \pmod{4}$, $p \neq 3$ and let R, N be the number of quadratic residues and nonresidues in $(0, \frac{p}{2})$. Then the class number of $\mathbb{Q}(\sqrt{-p})$ is*

$$h = \begin{cases} \frac{1}{3}(R - N), & p \equiv 3 \pmod{8} \\ R - N, & p \equiv 7 \pmod{8}. \end{cases}$$

In particular, $R > N$.

Proof. Let $K = \mathbb{Q}(\sqrt{-p})$. Note χ is odd, because it equals $\left(\frac{\cdot}{p}\right)$ by (23.17), and $p \equiv 3 \pmod{4}$. Let $h = h_K$. We equate (23.8) with (23.14) to get

$$\begin{aligned}
 \frac{2\pi h}{\sqrt{p}w_K} &= L(1, \chi) = \frac{\pi i \tau(\chi)}{p^2} \sum_{k \in (\mathbb{Z}/p)^\times} \overline{\chi(k)} k \\
 &= \frac{-\pi}{p\sqrt{p}} \sum_{k \in (\mathbb{Z}/p)^\times} \chi(k) k, \\
 \implies h &= -\frac{1}{p} \sum_{k \in (\mathbb{Z}/p)^\times} \chi(k) k,
 \end{aligned}$$

where, because χ is odd, $\tau(\chi) = i\sqrt{p}$ by Theorem 23.4.2, and we note $w_K = 2$ for $p \neq 3$.

Now we proceed in two steps.

1. Because χ is odd, we can pair up $\chi(k)$ and $\chi(p - k) = -\chi(k)$ to get

$$\begin{aligned}
 h &= -\frac{1}{p} \sum_{0 < k < \frac{p}{2}} \chi(k) k + (-\chi(k))(p - k) \\
 &= -\frac{2}{p} \sum_{0 < k < \frac{p}{2}} \chi(k) k + \sum_{0 < k < \frac{p}{2}} \chi(k). \tag{23.18}
 \end{aligned}$$

2. We use a trick to get rid of the sum of $\chi(k)k$. We get another expression for h by first flipping some of the k into $m - k$, so that we hit exactly the multiples of 2, and then take out a $\chi(2)$:

$$\begin{aligned}
 h &= -\frac{1}{D} \sum_{k \in (\mathbb{Z}/p)^\times} \chi(k)k \\
 &= -\frac{1}{D} \left(\sum_{\substack{k \in (\mathbb{Z}/p)^\times \\ k \text{ even}}} \chi(k)k + \sum_{\substack{k \in (\mathbb{Z}/p)^\times \\ k \text{ odd}}} \chi(k)k \right) \\
 &= -\frac{1}{D} \left(\sum_{\substack{k \in (\mathbb{Z}/p)^\times \\ k \text{ even}}} \chi(k)k + \sum_{\substack{k \in (\mathbb{Z}/p)^\times \\ k \text{ odd}}} -\chi(m - k)k \right) \\
 &= -\frac{1}{D} \left(\sum_{\substack{k \in (\mathbb{Z}/p)^\times \\ k \text{ even}}} \chi(k)k + \sum_{\substack{k \in (\mathbb{Z}/p)^\times \\ k \text{ even}}} -\chi(k)(m - k) \right) \\
 &= \sum_{0 < k < \frac{p}{2}} \chi(2k)((2k) + (2k - m)) \\
 \implies h\chi(2) &= -\frac{1}{p} \sum_{0 < k < \frac{p}{2}} \chi(k)(4k - m) \\
 &= -\frac{4}{p} \sum_{0 < k < \frac{p}{2}} \chi(k)k + \sum_{0 < k < \frac{p}{2}} \chi(k). \tag{23.19}
 \end{aligned}$$

Now multiply (23.18) by 2 and subtract (23.19) to get

$$h(2 - \chi(2)) = \sum_{0 < k < \frac{p}{2}} \chi(k).$$

Noting $\chi(k) = \left(\frac{k}{p}\right)$, $\left(\frac{2}{p}\right) = (-1)^{\frac{p^2-1}{8}}$ gives the result. \square

Note that there does not seem to be an “elementary” proof of $R > N$ for $p \equiv 3 \pmod{8}$. See <http://mathoverflow.net/questions/25707/intuition-for-a-formula-that-exp> for a discussion.

3 For further study

- See [BS66, §5.5] or [Was82, Chapter 4] for the application of L -series to class numbers of cyclotomic fields.
- The conjectural analogue of the analytic class number formula for elliptic curves is the Birch-and-Swinnerton Dyer conjecture.

Conjecture 23.3.1 (Strong Birch and Swinnerton-Dyer conjecture). *We have*

$$\lim_{s \rightarrow 1} (s - 1)^{\text{rank } E} L(E, s) = \frac{\Omega_E \text{Reg} E(\mathbb{Q}) |\text{III}(E/\mathbb{Q})| \prod_p c_p}{|E(\mathbb{Q})_{\text{tors}}|^2}.$$

where $c_p = [E(\mathbb{Q}_p) : E_0(\mathbb{Q}_p)]$ is the Tamagawa number of E/\mathbb{Q}_p . Here

- $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}} = \langle P_1, \dots, P_r \rangle$ comes with a quadratic form (canonical height)
- $\text{Reg} E(\mathbb{Q}) = \det([P_i, P_j])_{i,j=1,\dots,r}$ where $[P, Q] = \hat{h}(P + Q) - \hat{P} - \hat{Q}$. The regulator measures the volume of a fundamental domain of the associated lattice.
- $\Omega_E = \int_{E(\mathbb{R})} \frac{dx}{|2y+a_1+a_3|}$ where a_i are the coefficients of a globally minimal Weierstrass equation.

4 Appendix

Proposition 23.4.1: Let P be the fundamental parallelogram of $L(U_K)$, and let Vol be the $(r_1 + r_2 - 1)$ -dimensional area. Then

$$\text{Reg}_K = \frac{\text{Vol}(P)}{\sqrt{r_1 + r_2}}.$$

Proof. The regulator is defined by deleting any column of the matrix with rows $L(\varepsilon_i)$, i.e., it is calculated by taking a fundamental parallelotope, projecting it to any hyperplane $x_i = 0$, and then taking the area. (The proof will show these projections have the same area.)

Let $t = r_1 + r_2$. Let $L(\varepsilon_i) = (a_{i,1}, \dots, a_{i,t-1}, -a_{i,1} - \dots - a_{i,t-1})$. (Recall that the coordinates sum to 0.) To find the volume within the hyperplane, we note that $(\frac{1}{\sqrt{t}}, \dots, \frac{1}{\sqrt{t}})$ is a unit vector perpendicular to the hyperplane, and evaluate

$$\begin{vmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,t-1} & -a_{1,1} - a_{1,2} - \cdots - a_{1,t-1} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,t-1} & -a_{2,1} - a_{2,2} - \cdots - a_{2,t-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{t-1,1} & a_{t-1,2} & \cdots & a_{t-1,t-1} & -a_{t-1,1} - a_{t-1,2} - \cdots - a_{t-1,t-1} \\ \frac{1}{\sqrt{t}} & \frac{1}{\sqrt{t}} & \cdots & \frac{1}{\sqrt{t}} & \frac{1}{\sqrt{t}} \end{vmatrix}.$$

Take the cofactor expansion along the bottom row. The i th term, for $i < t$, is

$$(-1)^{i-1} \frac{1}{\sqrt{t}} \begin{vmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,t-1} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,t-1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{t-1,1} & a_{t-1,2} & \cdots & a_{t-1,t-1} \end{vmatrix}$$

but with the i th column deleted and the column $(-a_{i,1} - \cdots - a_{i,t-1})_{1 \leq i \leq t-1}$ added as the last column. Add the first $t - 2$ columns to this column (which doesn't change the determinant); permute the columns with the permutation $t - 1 \rightarrow i \rightarrow i + 1 \rightarrow \cdots$, which has sign

$$(-1)^{t-i-1}; \text{ and take out a } -1 \text{ to get } (-1)^{t+1} \frac{1}{\sqrt{t}} \begin{vmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,t-1} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,t-1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{t-1,1} & a_{t-1,2} & \cdots & a_{t-1,t-1} \end{vmatrix}. \text{ (Incidentally}$$

this shows that deleting any column gives the same absolute value of determinant, so that the regulator is well-defined.) Adding these all up gives

$$\text{Vol}(P) = t \frac{1}{\sqrt{t}} \left\| \begin{matrix} a_{1,1} & a_{1,2} & \cdots & a_{1,t-1} \\ a_{2,1} & a_{2,2} & \cdots & a_{2,t-1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{t-1,1} & a_{t-1,2} & \cdots & a_{t-1,t-1} \end{matrix} \right\| = \sqrt{t} \text{Reg}_K.$$

□

A character (of the multiplicative group) is **quadratic** if $\chi^2 = \chi_0$ and **primitive** modulo m if χ is not a character modulo m' for any $1 \neq m' \mid m$.

Theorem 23.4.2. *Let χ be a primitive quadratic character modulo m . Then the Gauss sum $\tau(\chi)$ satisfies*

$$\tau(\chi) = \begin{cases} \sqrt{m} & \text{if } \chi \text{ is even } (\chi(-1) = 1), \\ i\sqrt{m} & \text{if } \chi \text{ is odd } (\chi(-1) = -1). \end{cases}$$

Proof. This is somewhat involved. See [BS66, §5.5, Theorem 7]. (See Theorem ??) □

Appendix A

Multiplication modulo n

1 Introduction

In Chapter ?? you learned how modular arithmetic works: you fix a number n , and then compute by taking all integers modulo n , that is, you throw away all information except the remainder when dividing by n . This system obeys all the “rules” we expect it to, and is especially well-behaved when n is a prime.

There is a lot of similarity between addition and multiplication modulo n , but it seems multiplication is less “transparent.” Compare the following.

1. We saw how we could use modular arithmetic to solve problems such as: what is the last digit of $7 \cdot 2011$? What is the last digit of 7^{2011} ?

You probably see immediately the first answer is 7. To solve the second problem we looked for a repeating pattern in the powers of 7 modulo 10: 1, 7, 9, 3, ... We have $7^4 \equiv 1 \pmod{10}$, so the “cycle” repeats after every 4 times, and $7^{2011} \equiv 7^3 \equiv 3 \pmod{10}$.

Making an analogy, note that if look at the sequence $7, 2 \cdot 7, 3 \cdot 7, \dots$ we get back to 0 after 10 steps, while in the sequence $7, 7^2, 7^3$ we get back to 1 after 4 steps. It’s easy to understand how the cycle repeats when we add repeatedly (i.e., multiply), but what about when we multiply repeatedly (i.e., take powers)?

In general, given $a \pmod{n}$, how many times do we have to multiply a by itself to get back to 1 (what is the “order” of a modulo n)? How does this depend on a and n ? Is there some general rule that determines the order?

2. We have the notion of inverses. For instance, what is the additive inverse of 7 modulo 19, i.e., what do we add to 7 to get 0 modulo 19? It is simply $-7 \equiv 12 \pmod{19}$, because

$$7 + 12 \equiv 0 \pmod{19}.$$

What is the multiplicative inverse of 7 modulo 19, i.e., what do we multiply 7 by to get 1 modulo 19? You might have to think a bit more on this one—it is $11 \pmod{19}$,

because

$$7 \cdot 11 = 77 \equiv 1 \pmod{19}.$$

In this chapter you will learn to completely determine the multiplicative structure of the integers modulo n . In Section 2, we will formalize the notion of the **order** of an element: how many times we have to multiply a number to get back to 1 modulo n . We will develop a toolbox of basic theorems: Euler's Theorem and Fermat's Little Theorem in Section 3 and Wilson's Theorem in Section 8. Euler's Theorem will give us m such that $a^m \equiv 1 \pmod{n}$.

We will use these theorems to solve a variety of problems in Section 4. For instance (keep in the theme of repeating patterns) we will explain why the period of the repeating decimals $\frac{1}{17}$ and $\frac{1}{19}$ are 16 and 18:

$$\begin{aligned} \frac{1}{17} &= .\underbrace{0588235294117647}_{16} \\ \frac{1}{19} &= .\underbrace{052631578947368421}_{18}. \end{aligned}$$

At this point we find that a lot of the stuff we've been doing works in a much more *abstract* setting, that of **groups**. In Section 5 we recast all the theorems in the language of group theory, and in this way give a way think about the similarities behind addition and multiplication modulo n . With the language of group theory, we give a complete characterization of the multiplicative structure of $\mathbb{Z}/n\mathbb{Z}$ in Sections 6 and 7.

Finally we give two applications, to primality testing (Section ??) and basic cryptography (Section ??).

2 Order of an element

We've seen that the values taken by a powers of an element modulo n form a repeating pattern, and under certain relative primality conditions, start each cycle at 1. For example,

$$\begin{aligned} 3^0 &\equiv 1 \pmod{5} \\ 3^1 &\equiv 3 \pmod{5} \\ 3^2 &\equiv 4 \pmod{5} \\ 3^3 &\equiv 2 \pmod{5} \\ 3^4 &\equiv 1 \pmod{5} \\ 3^5 &\equiv 3 \pmod{5}, \end{aligned}$$

so the powers of 3 cycle 1, 3, 4, 2, ... modulo 5. In particular, we get back to 1 in 4 steps: $3^4 \equiv 1 \pmod{5}$. Hence we call 4 the *order* of 3:

Definition 1.2.1: Let $n > 1$ and let a be an integer relatively prime to n . The **order** of a modulo n is the smallest positive integer m such that $a^m \equiv 1 \pmod{n}$. In symbols,

$$\text{ord}_n(a) = \min \{m \in \mathbb{N} : a^m \equiv 1 \pmod{n}\}.$$

For example,

$$\text{ord}_5(3) = 4.$$

Note that we place the modulus as an

Note that the order is well-defined for all a relatively prime to n : Indeed, there are only a finite number of residues modulo n , so two powers of a must be equal modulo n . So suppose $0 < m_1 < m_2$ and

$$a^{m_1} \equiv a^{m_2} \pmod{n}.$$

Since a is relatively prime to n , we can take inverses to find $a^{m_2-m_1} \equiv 1 \pmod{n}$.

Our first result is that the set of all positive integers k for which $a^k \equiv 1 \pmod{n}$ is completely determined by its smallest element, i.e. the order. In the case above, the set of all m such that $3^m \equiv 1 \pmod{5}$ is exactly the set of multiples of 4.

Proposition 1.2.2: Let $n > 1$ and $a \perp n$.

1. The set of m such that $a^m \equiv 1 \pmod{n}$ is exactly the set of multiples of $\text{ord}_n(a)$. In other words,

$$a^m \equiv 1 \pmod{n} \iff \text{ord}_n(a) \mid m.$$

2. The numbers

$$1, a, \dots, a^{\text{ord}_n(a)-1}$$

are all distinct, and every power of a is congruent to one of these.

Proof. Let $d = \text{ord}_n(a)$.

1. The reverse direction is easy: If $d \mid m$, then write $m = dk$. We have

$$a^m \equiv (a^d)^k \equiv 1^k \equiv 1 \pmod{n}.$$

Conversely, suppose that $a^m \equiv 1 \pmod{n}$. We use the same technique as [gcd?], noting that we picked $\text{ord}_n(a)$ to be the *least* positive integer with this property. Using division with remainder, write

$$m = dk + r, \quad 0 \leq r < d.$$

We have

$$a^r = a^{m-dk} = a^m a^{-dk} \equiv a^m \equiv 1 \pmod{n}.$$

Since d is the least positive integer for which $a^d \equiv 1 \pmod{n}$, and $0 \leq r < d$, we must have $r = 0$. Hence $d \mid m$.¹

¹For another way to phrase this proof, see Problem C.1.

2. For the second part, writing $m = dk + r$ as above we note that

$$a^{dk+r} = (a^d)^k a^r \equiv a^r \pmod{n}.$$

If $0 \leq r_1 < r_2 < \text{ord}_n(a)$, then $0 < r_2 - r_1 < \text{ord}_n(a)$ implies $a^{r_2-r_1} \not\equiv 1 \pmod{n}$ and hence $a^{r_1} \not\equiv a^{r_2} \pmod{n}$. \square

Now we have an abstract description of the numbers m such that $a^m \equiv 1 \pmod{n}$. We know there is some positive integer with this property, and that all others are multiples of that number. But we would like something more concrete: is there some m depending on n , so that we will always have $a^m \equiv 1 \pmod{n}$? The next section will answer that question.

3 Euler's theorem and Fermat's little theorem

Theorem 1.3.1 (Euler's theorem). *Let $n > 1$ be an integer. For any integer a relatively prime to n , $\text{ord}_n(a) \mid \varphi(n)$ and*

$$a^{\varphi(n)} \equiv 1 \pmod{n}.$$

Corollary 1.3.2 (Fermat's little theorem). *Let p be a prime. For any integer a ,*

$$a^p \equiv a \pmod{p}.$$

If $a \not\equiv 0 \pmod{p}$, then

$$a^{p-1} \equiv 1 \pmod{p}.$$

Let G be the set of invertible residues modulo n . We present two proofs.

Proof 1. Let m_a denote the function $G \rightarrow G$ defined by

$$m_a(g) = ag.$$

Note that this is an invertible function as its inverse is

$$m_a^{-1}(g) = a^{-1}g.$$

Hence it is a bijection $G \rightarrow G$. This means that the elements $ag, g \in G$ are an reordering of the elements of G . Hence

$$\prod_{g \in G} ag \equiv \prod_{g \in G} g \pmod{n}.$$

Dividing both sides by $\prod_{g \in G} g$ and noting $|G| = \varphi(n)$ gives

$$a^{\varphi(n)} \equiv 1 \pmod{n}. \quad \square$$

Proof 2. The main idea is that there are $\varphi(n)$ possible invertible residues modulo n , and so the number of elements in the set $H := \{a^m \bmod n : m \in \mathbb{N}\}$ must be a divisor of $\varphi(n)$. To show this we show that “translates” of this set cover all $\varphi(n)$ nonzero residues without overlap. The fact that H has nice multiplicative structure will be essential.

First note the following facts.

1. $1 \in H$. (This is because $1 = a^0$.)
2. If $h \in H$ then $h^{-1} \in H$. (If $h \equiv a^m \pmod{n}$ then $h^{-1} \equiv a^{-m} \pmod{n}$.)
3. If $h_1, h_2 \in H$ then $h_1 h_2 \in H$. (If $h_j \equiv a^{m_j} \pmod{n}$ then $h_1 h_2 \equiv a^{m_1+m_2} \pmod{n}$.)

Given two nonzero residues b, c modulo p , we write $b \sim c$ if $\frac{b}{c} \in H$. We claim that \sim is an equivalence relation. We check the following.

1. $b \sim b$: This holds by item 1 above, since $\frac{b}{b} = 1$.
2. If $b \sim c$ then $c \sim b$: This holds by item 2 above, since $\frac{b}{c} = \frac{c}{b}$.
3. If $b \sim c$ and $c \sim d$ then $b \sim d$: This holds by item 3 above since $\frac{b}{d} = \frac{b}{c} \cdot \frac{c}{d}$.

Thus G is split into equivalence classes. If C is an equivalence class and c is any element in C , then we have

$$C = \{d : d \sim c\} = \left\{d : \frac{d}{c} \in H\right\} = \{ch : h \in H\}.$$

Since multiplication by c is invertible, C has $|H|$ elements. (It is the RHS that suggests the sets C are “translates” of H .)

Thus, letting $[G : H]$ denote the number of equivalence classes, we have

$$|G| = [G : H]|H|.$$

Hence $|H|$ divides $|G| = \varphi(n)$. But by Proposition 1.2.2(2), $|H| = \text{ord}_n(a)$. Since $\text{ord}_n(a) \mid \varphi(n)$, by Proposition 1.2.2(1), we get

$$a^{\varphi(n)} \equiv 1 \pmod{n}. \quad \square$$

Although the first proof is shorter, the first reveals hints at some important ideas with broad generalizations, which we will discuss in Section 5.

Proof of Corollary 1.3.2. Since $\varphi(p) = p - 1$ and the invertible residues modulo p are exactly the nonzero residues, we get

$$a^{p-1} \equiv 1 \pmod{p}$$

for $a \not\equiv 0 \pmod{p}$. Multiplying by a gives the first statement for $a \not\equiv 0 \pmod{p}$. If $a \equiv 0 \pmod{p}$ the first statement obviously holds. \square

Remark 1.3.3: The converse of Fermat’s little theorem is not true: if $a^p \equiv a \pmod{p}$ for all a , then p is not necessarily prime. For example, $2^{11 \cdot 31} \equiv 2 \pmod{11 \cdot 31}$, but $11 \cdot 31$ is not a prime. Indeed, there are certain numbers n such that for all integers a , we have $a^n \equiv a \pmod{n}$ with n not a prime. Such numbers are called *Carmichael numbers*, and the first few are given by $n = 561, 1105, 1729, 2465$.

4 Examples

4.1 Using Euler's theorem

Without further ado, we give some applications of Fermat's little theorem and Euler's theorem. The first, most popular application is in finding large powers modulo a certain number. While before, we had to evaluate a, a^2, a^3, \dots until we got back to 1, our work is now shorter.

Example 1.4.1: Find $3^{1006} \pmod{2012}$.

The prime factorization of 2012 is $2^2 \cdot 503$, so $\varphi(2012) = 2 \cdot 502 = 1004$. As 3 is relatively prime to 2012, by Euler's Theorem

$$3^{1006} \equiv 3^2 \equiv 9 \pmod{2012}.$$

“Find big power modulo n problem”

“Tower of exponents problem”

Remark about “thinking backwards”

Example 1.4.2: Show that for all primes $p \geq 7$, the number $\underbrace{11 \cdots 1}_{p-1}$ is divisible by p .

Solution. The key to this problem is writing an algebraic expression for $\underbrace{11 \cdots 1}_{p-1}$. By the geometric series formula,

$$\underbrace{11 \cdots 1}_{p-1} = 1 + 10 + \cdots + 10^{p-2} = \frac{10^{p-1} - 1}{9}.$$

Because $p \nmid 10$, by Fermat's little theorem 1.3.2 we have

$$10^{p-1} \equiv 1 \pmod{p} \implies p \mid 10^{p-1} - 1.$$

Because $\gcd(9, p) = 1$, we have $(10^{p-1} - 1)/9 \equiv 0 \pmod{p}$ as desired.

4.2 Computing the order

The following proposition gives practical ways to compute the order of an element.

Proposition 1.4.3: Let $n > 1$, let a be an integer relatively prime to n , and set $d = \text{ord}_n(a)$.

1. (Power of the base)

$$\text{ord}_n(a^k) = \frac{d}{\gcd(d, k)}.$$

2. (Multiplying the base) Let $d = \text{ord}_n(a), c = \text{ord}_n(b)$. If $\gcd(d, c) = 1$, then $\text{ord}_n(ab) = dc$.

3. (Multiplying the modulus) Let the prime factorization of n be $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$. Let $d_i = \text{ord}_{p_i^{\alpha_i}}(a)$. Then

$$d = \text{lcm}(d_1, d_2, \dots, d_k).$$

Warning: It is not necessarily true that $\text{ord}_n(ab) = \text{lcm}(\text{ord}_n(a), \text{ord}_n(b))$.

Proof. 1. Set $m = \text{gcd}(d, k)$. Write $d = md_1, k = mk_1$, where $\text{gcd}(d_1, k_1) = 1$. Set $t = \text{ord}_n(a^k)$. Then we have

$$(a^k)^{d_1} = a^{mk_1 d_1} = (a^d)^{k_1} \equiv 1 \pmod{n}.$$

Hence $t \leq d_1$. On the other hand, $a^{kt} = (a^k)^t \equiv 1 \pmod{n}$. Then we have $d \mid kt$, hence $d_1 \mid k_1 t$. As d_1, k_1 are relatively prime, we have $d_1 \mid t$, hence $d_1 \leq t$. It follows that $t = d_1$ as desired.

2. Set $e = \text{ord}_n(ab)$. Then we have $(ab)^e \equiv 1 \pmod{n}$. Hence $(a^{ce})(b^{ce}) = a^{ce}(b^c)^e \equiv a^{ce} \equiv 1 \pmod{n}$. Hence $d \mid ce$. As $\text{gcd}(d, c) = 1$, we have $d \mid e$. Analogously, we have $(a^{de})(b^{de}) = (a^d)^e b^{de} \equiv b^{de} \equiv 1 \pmod{n}$. Hence $c \mid de$, so $c \mid e$. As $\text{gcd}(d, c) = 1$, we have $dc \mid e$. However, we have $(ab)^{dc} = (a^d)^c (b^c)^e \equiv 1 \cdot 1 = 1 \pmod{n}$. Hence $dc = e$ as desired.
3. Set $e = \text{ord}_n(ab)$. Then we have $(ab)^e \equiv 1 \pmod{n}$. Hence $(a^{ce})(b^{ce}) = a^{ce}(b^c)^e \equiv a^{ce} \equiv 1 \pmod{n}$. Hence $d \mid ce$. As $\text{gcd}(d, c) = 1$, we have $d \mid e$. Analogously, we have $(a^{de})(b^{de}) = (a^d)^e b^{de} \equiv b^{de} \equiv 1 \pmod{n}$. Hence $c \mid de$, so $c \mid e$. As $\text{gcd}(d, c) = 1$, we have $dc \mid e$. However, we have $(ab)^{dc} = (a^d)^c (b^c)^e \equiv 1 \cdot 1 = 1 \pmod{n}$. Hence $dc = e$ as desired. □

Problem 1.4.4: Let $a > 1$ and n be positive integers. Show that n divides $\varphi(a^n - 1)$.

Proof. We have that the order of a modulo $a^n - 1$ is n . But we have $\text{ord}_{a^n - 1}(a) \mid \varphi(a^n - 1)$, hence $n \mid \varphi(a^n - 1)$ as desired. □

Note that trying to use the formula for $\varphi(m)$ in terms of the prime factorization of m doesn't work for this problem.

Problem 1.4.5: Determine all positive integers n such that n divides $2^n - 1$.

Proof. We shall show that $n = 1$ is the only solution. Suppose that $n \mid 2^n - 1$ for $n > 1$. Then n must be odd. Let p be the least prime divisor of n ; then $2^n \equiv 1 \pmod{p}$. Write $d = \text{ord}_p(2)$. Then $d > 1$ and $d \mid n$. Hence $p \leq d$ since p is the least prime divisor (and hence least divisor greater than 1) of n . But by Fermat's little theorem, $2^{p-1} \equiv 1 \pmod{p}$. Hence $d \mid p - 1$ - that is, $d \leq p - 1$. Hence $p \leq p - 1$. Contradiction; hence no such n exist. □

Problem 1.4.6: Let a be a positive integer, and let $p, q > 2$ be primes with $a^p \equiv 1 \pmod{q}$. Prove that either $q \mid a - 1$ or $q = 1 + 2kp$ for some positive integer k .

Proof. Obviously we have $\gcd(a, q) = 1$. Write $d = \text{ord}_q(a)$. Then we have $d \mid p$, so $d = 1$ or $d = p$. If $d = 1$, then $a \equiv 1 \pmod{q}$, so $q \mid a - 1$. If $d = p$, then we have $p \mid \varphi(q) = q - 1$, so $q = 1 + np$ for some integer n . But as $q > 2$ is a prime, q must be odd. As p is odd, we must have n even for q to be odd. Writing $n = 2k$, we find $q = 1 + 2kp$ as desired. \square

Problem 1.4.7: Let p, q be primes with $a^{p-1} + a^{p-2} + \cdots + a + 1 \equiv 0 \pmod{q}$. Prove that either $q = p$ or $q \equiv 1 \pmod{p}$.

Proof. If $p = 2$ then either $q = 2 = p$ or q is odd and $q \equiv 1 \pmod{p}$. The case $p > 2$, $q = 2$ is impossible since the left-hand expression is odd. Now we have the case where p and q are odd. Then we have $a^p - 1 = (a - 1)(a^{p-1} + a^{p-2} + \cdots + a + 1)$, hence $a^p \equiv 1 \pmod{q}$. Thus either $q \equiv 1 \pmod{p}$ or $q \mid a - 1$. If $q \mid a - 1$, then we have $a \equiv 1 \pmod{q}$, hence $1^{p-1} + 1^{p-2} + \cdots + 1 + 1 = p \equiv 0 \pmod{q}$, from which it follows that $p = q$. \square

Problem 1.4.8: Let n be an odd positive integer. Prove that if $n \mid 3^n + 1$ then $n = 1$.

Proof. Obviously n is not divisible by 3. Suppose that $n > 1$ and let p be the least prime divisor of n ; then $p \geq 5$. Write $d = \text{ord}_p(3)$. As $3^n \equiv -1 \pmod{n}$, we have $3^{2n} \equiv 1 \pmod{p}$, so $d \mid 2n$. As $3^{p-1} \equiv 1 \pmod{p}$, we have also $d \mid p - 1$. If d is odd, then we have $d \mid n$. As p is the least divisor of n greater than 1, we must have thus $d = 1$. Hence $3 \equiv 1 \pmod{p}$, implying $p = 2$. But $p \geq 5$; contradiction. Hence d must be even. Write $d = 2k$; then $k \mid n$, and if $k > 1$ then we have $1 < k < d < p$, contradicting the fact that p is the minimal divisor of n greater than 1. Hence $d = 2$ and $3^2 \equiv 1 \pmod{p}$, so $p = 2$. Contradiction; hence $n = 1$. \square

Problem 1.4.9: Let $\gcd(a, b) = 1$ with b odd. Show that $\gcd(n^a + 1, n^b - 1) \leq 2$ for any natural number n .

Proof. Write $l = \gcd(n^a + 1, n^b - 1)$, and suppose that $l > 1$. Write $d = \text{ord}_l(n)$. Then we have $n^b \equiv 1 \pmod{l}$, so $d \mid b$. Hence d is odd. But then as $n^a \equiv -1 \pmod{l}$, we have $d \mid 2a$. Hence $d \mid a$. If $d > 1$ then we have $d \mid a, b$, so $\gcd(a, b) > 1$. Contradiction; hence $d = 1$. Thus $n^a \equiv 1 \pmod{l}$, hence $1 \equiv 1 \pmod{l}$. Hence $l = 1$ or $l = 2$ as desired. \square

Problem 1.4.10 (IMO 1990/3): Determine all positive integers n such that n^2 divides $2^n + 1$.

Proof. Clearly $n = 1$ is a solution. Suppose that $n > 1$; then n is odd. Let p be the least prime divisor of n , and write $d = \text{ord}_p(2)$. As $2^{2n} \equiv 1 \pmod{p}$ we have $d \mid 2n$. As $2^{p-1} \equiv 1 \pmod{p}$ we have $d \mid p - 1$. If $d > 2$, then let q be a prime greater than 2 dividing d . Then $q \mid 2n$ and $q \mid p - 1$, contradicting the fact that p is the minimal prime dividing n . But we have $d > 1$, hence $d = 2$ so $p = 3$.

Write $n = 3^s m$, with $s, m \geq 1$ and $3 \nmid m$. Suppose that $s > 1$. Then we have

$$3^{2s} \mid (2^{3^s} + 1) (2^{3^s(m-1)} - 2^{3^s(m-2)} + \cdots - 2^{3^s} + 1).$$

But since $2^{3^s} \equiv -1 \pmod{3}$, we have $2^{3^s(m-1)} - 2^{3^s(m-2)} + \dots - 2^{3^s} + 1 \equiv 1 + 1 + \dots + 1 = m \not\equiv 0 \pmod{3}$. Hence $3^{2s} \mid 2^{3^s} + 1$.

We claim that for all s , we have $3^{s+2} \nmid 2^{3^s} + 1$. We may write

$$2^{3^s} + 1 = \left(3^{3^s} - \binom{3^s}{1} 3^{3^s-1} + \binom{3^s}{2} 3^{3^s-2} - \dots - \binom{3^s}{3^s-2} 3^2 + \binom{3^s}{3^s-1} 3^1 - 1 \right) + 1.$$

But then we have 3^{s+2} divides all terms in this expansion except $\binom{3^s}{3^s-1} 3^1$; hence $3^{s+2} \nmid 2^{3^s} + 1$.

As we have $3^{2s} \mid 2^{3^s} + 1$, we have thus $2s < s + 2$. Hence $s = 1$, so $n = 3m$. Suppose that $m > 1$. Let q be the least prime divisor of m ; then $q \geq 5$. Write $e = \text{ord}_q(2)$; then as we have $2^{2n} \equiv 1 \pmod{q}$, we have $e \mid 2n = 6m$. As $2^{q-1} \equiv 1 \pmod{q}$, we have also that $l \mid q - 1$. Hence we cannot have $l \mid n$, as this would contradict the fact that q is the smallest prime divisor of n . Thus as $q \geq 5$ we have $l = 3$ or $l = 6$, meaning that $q = 7$. But in this case we have $7 \mid 2^n + 1$; however, as $n = 3m$, we have $2^n + 1 = (2^3)^m + 1 \equiv 1^m + 1 = 2 \pmod{7}$. Contradiction; hence $m = 1$, so $n = 3$. Thus $n = 1, 3$ are our only solutions. \square

Example 1.4.11: Let p, q, r be distinct primes such that

$$pq \mid r^p + r^q.$$

Prove that either p or q equals 2.

Solution Suppose the relation holds but $p \neq 2, q \neq 2$. By Fermat's Little Theorem, $r^p \equiv r \pmod{p}$ and $r^q \equiv r \pmod{q}$. Then since r is relatively prime to p, q ,

$$\begin{aligned} r^p + r^q &\equiv 0 \pmod{p} \implies \\ r^{q-1} &\equiv -1 \pmod{p} \\ r^p + r^q &\equiv 0 \pmod{q} \implies \\ r^{p-1} &\equiv -1 \pmod{q} \end{aligned}$$

Since $-1 \not\equiv 1 \pmod{p, q}$, we get

$$\text{ord}_p(r) \nmid q - 1, \text{ord}_q(r) \nmid p - 1. \tag{A.1}$$

Since

$$\begin{aligned} r^{2(q-1)} &\equiv 1 \pmod{p} \\ r^{2(p-1)} &\equiv 1 \pmod{q}, \end{aligned}$$

we get

$$\text{ord}_p(r) \mid 2(q - 1), \text{ord}_q(r) \mid 2(p - 1). \tag{A.2}$$

For an integer n let $v_2(n)$ denote the highest power of 2 dividing n . Let $x = v_2(\text{ord}_p(r))$ and $y = v_2(\text{ord}_q(r))$. From relations in (A.1) and (A.2),

$$\begin{aligned} x &= v_2(2(q - 1)) = v_2(q - 1) + 1 \\ y &= v_2(2(p - 1)) = v_2(p - 1) + 1. \end{aligned} \tag{A.3}$$

By Fermat's Little Theorem, $\text{ord}_p(r) \mid p - 1$ and $\text{ord}_q(r) \mid q - 1$. Hence

$$\begin{aligned} x &\leq v_2(p - 1) \\ y &\leq v_2(q - 1) \end{aligned} \tag{A.4}$$

Putting (A.3) and (A.4) together, we get $x \leq y - 1, y \leq x - 1$, contradiction.

5 Groups

For the moment, it is helpful to “forget” where our set comes from and just work from the basic axioms that it satisfies.

Definition 1.5.1: A **group** is a set G together with a binary operation \circ , satisfying the following properties:

1. (Associative law) For any $a, b, c \in G$,

$$(a \circ b) \circ c = a \circ (b \circ c).$$

2. (Identity) There exists an element id , called the identity, such that for all a ,

$$\text{id} \circ a = a \circ \text{id} = a.$$

3. (Inverses) For any a there exists an element a' , called the inverse of a , such that

$$a \circ a' = a' \circ a = \text{id}.$$

G is called an **abelian group** if additionally it satisfies the following.

4. (Commutativity) For all $a, b \in G$, $a \circ b = b \circ a$.

We will be dealing exclusively with abelian groups.

Define order, exponent. Largest order IS the exponent (for abelian groups)

6 Primitive roots

We now know that $a^{\varphi(n)} \equiv 1 \pmod{n}$ for all a relatively prime to n , and that $\text{ord}_n(a) \mid \varphi(n)$. We can ask, does there exist a for which $\text{ord}_n(a)$ is exactly n ?

Equivalently, by the second part of Proposition 1.2.2, since there are $\varphi(n)$ possible invertible residues modulo n , this says that the powers of a achieve every possible invertible residue modulo n .

Definition 1.6.1: A **primitive root** modulo n is an integer a such that

$$\text{ord}_n(a) = \varphi(n).$$

For example, 3 is a primitive root modulo 5, as $\text{ord}_5(3) = 4$.

Theorem 1.6.2. *Primitive roots exist modulo n if and only if $n = 2, 4, p^k$, or $2p^k$ for p an odd prime.*

Moreover, if g is a primitive root modulo p^2 , then it is a primitive root modulo p^k and $2p^k$ for any k .

Proof. We will prove the “if” part of the theorem. The “only if” part will fall out from Theorem 1.7.1 in the next section.

For $n = 2$ or 4 , we see that 1 and 3 are primitive roots, respectively.

Part 1: Now suppose $n = p$ is prime. We note that by Fermat’s little theorem 1.3.2 that

$$x^{p-1} - 1 \equiv 0 \pmod{p}$$

for all nonzero residues x modulo p .

Note that if there are elements of order d_1, \dots, d_k then there is an element of order $\text{lcm}(d_1, \dots, d_k)$ (Proposition ??). Hence if d is the maximal order of an element in $(\mathbb{Z}/p\mathbb{Z})^\times$, then all orders must divide d . Hence

$$x^d - 1 \equiv 0 \pmod{p}.$$

Now we need the following lemma.

Lemma 1.6.3. *A nonzero polynomial $f(X) \in \mathbb{Z}/p\mathbb{Z}[X]$ of degree d has at most d roots.*

Proof. We induct on the degree. If $d = 0$ the assertion is clear. If $f(X)$ has a root, then

$$f(X) \equiv (X - a)g(X) \pmod{p}$$

for some $g(X) \in \mathbb{Z}/p\mathbb{Z}[X]$ of degree $d - 1$. Now $f(X) \equiv 0 \pmod{p}$ implies that one of the factors $X - a$ or $g(X)$ is 0 modulo p : this is because there are no zerodivisors modulo p . Hence the roots are a and the roots of $g(X)$; the latter total at most $d - 1$ by the induction hypothesis. \square

Now $x^d - 1 = 0$ can have at most d roots modulo p , but we know all $p - 1$ invertible residues are roots. Hence $d \geq p - 1$. But we know that the order of any element divides $p - 1$, so $d \mid p - 1$ and we get $d = p - 1$.

Part 2: Now we prove the theorem for p^k .

We first show that there is a primitive root modulo p^2 . Take a primitive root x modulo p ; suppose it is not primitive modulo p^2 . Now

$$p - 1 = \text{ord}_p(x) \mid \text{ord}_{p^2}(x) \mid p(p - 1)$$

where the right-hand divisibility is strict. Hence $\text{ord}_p(x) = p - 1$. Now note

$$\text{ord}_{p^2}(p + 1) = p,$$

since $(1+p)^k \equiv 1+kp \pmod{p^2}$ and this is 1 modulo p^2 for the first time when $k=p$. By Proposition ??(2),

$$\text{ord}_{p^2}(x(p+1)) = p(p-1) = \varphi(p^2)$$

so $x(p+1)$ is a primitive root modulo p^2 .

Now suppose $x \in \mathbb{Z}$ is a primitive root modulo p^2 . It attains every residue modulo p^2 so *a fortiori* it attains every residue modulo p , i.e. is primitive modulo p . We show by induction that

$$x^{p^{k-1}(p-1)} = p^k j + 1 \tag{A.5}$$

for some j not a multiple of p . For the case $k=1$, this is since x is a primitive root modulo p , but $x^{p-1} \not\equiv 1 \pmod{p^2}$. Suppose it proved for k ; then

$$x^{p^k(p-1)} = (p^k j + 1)^p = 1 + \binom{p}{1} p^k j + \binom{p}{2} p^{2k} j^2 + \cdots = 1 + p^{k+1}(j + pj')$$

for some j' . This shows the claim for $k+1$. Since

$$p-1 = \text{ord}_p(x) \mid \text{ord}_{p^k}(x) \mid p^{k-1}(p-1)$$

we know $\text{ord}_{p^k}(x)$ must be in the form $p^{j-1}(p-1)$ for some j . Equation (A.5) shows that $j=k$.

Part 3: Note that $\varphi(2p^k) = \varphi(p^k)$. Thus any primitive root modulo p^k is automatically a primitive root modulo $2p^k$. \square

Remark 1.6.4: Note the existence of a primitive root modulo n is equivalent to the fact that $(\mathbb{Z}/n\mathbb{Z})^\times$ is generated by one element, i.e. is the cyclic group $C_{\varphi(n)}$. Hence if there are primitive roots modulo p^k for all k , then the quotient maps

$$\cdots \rightarrow (\mathbb{Z}/p^3\mathbb{Z})^\times \rightarrow (\mathbb{Z}/p^2\mathbb{Z})^\times \rightarrow (\mathbb{Z}/p\mathbb{Z})^\times$$

correspond to maps

$$\cdots \rightarrow C_{p^2(p-1)} \rightarrow C_{p(p-1)} \rightarrow C_{p-1}.$$

Let g_k be a generator for $C_{p^{k-1}(p-1)}$; the kernel of the map $C_{p^{k-1}(p-1)} \rightarrow C_{p^{k-2}(p-1)}$ must be the cyclic group of order p generated by $g_k^{p^{k-2}(p-1)}$. Writing $x \equiv g_k^j \pmod{p^k}$, the conditions that $x \pmod{p^2}$ generates $(\mathbb{Z}/p^2\mathbb{Z})^\times$ translates into the fact that j is relatively prime to both $p(p-1)$, and hence that x is a primitive root modulo p^k .

This rationalizes the last statement of the theorem, and suggests that it should be used to prove the existence of primitive roots.

Remark 1.6.5: The proof of the first part can be generalized to the fact that all finite *fields* have a primitive root. See Proposition ??(2).

7 Multiplicative structure of $\mathbb{Z}/n\mathbb{Z}$

Theorem 1.7.1.

1. Suppose $p \neq 2$ is prime. Then

$$(\mathbb{Z}/p^n\mathbb{Z})^\times \cong C_{p^{n-1}(p-1)}.$$

2. For the case $p = 2$, for $n \geq 2$ we have

$$(\mathbb{Z}/2^n\mathbb{Z})^\times \cong C_2 \times C_{2^{n-2}}.$$

Moreover, $(\mathbb{Z}/2^n\mathbb{Z})^\times$ is generated by -1 , which has order 2, and 3, which has order 2^{n-2} . The isomorphism is given by $(-1)^a 3^b \leftrightarrow (a, b)$.

3. In general,

$$(\mathbb{Z}/p_1^{\alpha_1} \cdots p_n^{\alpha_n}\mathbb{Z})^\times \cong \prod (\mathbb{Z}/p_k^{\alpha_k}\mathbb{Z})^\times.$$

Proof. The first follows from existence of primitive roots modulo p^n .

For the second, we show by induction that for every $k \geq 1$,

$$3^{2^k} = 2^{k+2}j + 1$$

for some odd j . This is true for $k = 1$ as $3^2 = 8 + 1$. Suppose the above holds; then

$$3^{2^{k+1}} = (2^{k+2}j + 1)^2 + 1 = 2^{k+3}(j + 2^{k+1}j^2) + 1,$$

showing the induction step.

Note $\text{ord}_{2^n}(3)$ must divide $|(\mathbb{Z}/2^n\mathbb{Z})^\times| = 2^{n-1}$. The above then shows that $\text{ord}_{2^n}(3) = 2^{n-2}$. Finally, note that for $n \geq 3$, no power of 3 is equal to -1 modulo 2^n : if so, then by Theorem ??, $3^{2^{n-3}} = 3^{\frac{1}{2}\text{ord}_{2^n}(3)} \equiv -1 \pmod{2^n}$. However, $3^{2^{n-3}} \equiv 1 \pmod{2^{n-1}}$ by the above, so it is not congruent to -1 modulo 2^n .

The last follows from the Chinese Remainder Theorem. □

8 Wilson's theorem

Wilson's theorem is a multiplicative congruence of a slightly different kind.

Theorem 1.8.1 (Wilson's theorem). *A positive integer p is prime if and only if*

$$(p-1)! \equiv -1 \pmod{p}.$$

Proof. We may easily verify that the theorem is true for $p = 2, 3, 4$. Suppose that $p \geq 5$ is a prime; consider the set $S = \{2, 3, \dots, p-2\}$. We will show that for any $s \in S$ there exists some $s' \in S$ with $ss' \equiv 1 \pmod{p}$. Indeed, given such s we set $s' = s^{p-2}$. Then we have that $ss' = s^{p-1} \equiv 1 \pmod{p}$. Now if $s' \notin S$, then we have either $s' \equiv 1 \pmod{p}$ or $s' \equiv -1 \pmod{p}$. If $s' \equiv 1 \pmod{p}$, then $s \equiv 1 \pmod{p}$. This is obviously impossible. Similarly, if $s' \equiv -1 \pmod{p}$, then $s \equiv -1 \pmod{p}$. This is similarly impossible; hence we have $s' \in S$. Similarly, if we have $s, t \in S$ with $s' = t'$, then $s = t$. We may see that $ss' - tt' = (s-t)s' \equiv 0 \pmod{p}$. As $s' \not\equiv 0 \pmod{p}$, we must have $p \mid s-t$. As $|s-t| < p$, we thus have $|s-t| = 0$ as desired. Finally, $s \neq s'$; if $s = s'$, then we have $ss' = s^2 \equiv 1 \pmod{p}$, implying that $p \mid s-1$ or $p \mid s+1$. This cannot be true, as $s \not\equiv \pm 1 \pmod{p}$; it follows that $s \neq s'$.

Now we are ready to prove Wilson's theorem. As p is odd, and as $|S| = p-3$, there are an even number of elements in S . We pair these elements up into disjoint 2-element sets $\{s_1, s'_1\}, \{s_2, s'_2\}, \dots, \{s_{(p-3)/2}, s'_{(p-3)/2}\}$. These sets must contain all elements of S exactly once. Furthermore, when we take the product $s_1 s'_1 s_2 s'_2 \cdots s_{(p-3)/2} s'_{(p-3)/2}$ we will obtain 1, as the product of each pair is congruent to 1 modulo p . Hence we have

$$(p-1)! = 1 \cdot 2 \cdots p-1 \equiv 1 \cdot s_1 s'_1 s_2 s'_2 \cdots s_{(p-3)/2} s'_{(p-3)/2} \cdot p-1 \equiv 1 \cdot 1 \cdots -1 = -1 \pmod{p}$$

exactly as desired. \square

Problem 1.8.2: Let p be a prime of the form $4k+3$, and let a_1, a_2, \dots, a_{p-1} be consecutive positive integers. Prove that these numbers cannot be partitioned into two sets such that the products of the elements of the two sets are equal.

Proof. Suppose for a contradiction that there do exist sets $X = \{x_1, x_2, \dots, x_m\}, Y = \{y_1, y_2, \dots, y_n\}$ such that the product of the elements of X (denoted $P(X)$) and the product of the elements of Y (denoted $P(Y)$) are equal. If any of the a_i are divisible by p , then exactly one of the a_i may be divisible by p . In this case we have p dividing exactly one of $P(X), P(Y)$, so these products cannot be equal.

Now if $p \nmid a_i$ for $i = 1, 2, \dots, p-1$, then $a_i \equiv i \pmod{p}$. Hence

$$[P(X)]^2 = P(X)P(Y) = x_1 x_2 \cdots x_m y_1 y_2 \cdots y_n = a_1 a_2 \cdots a_{p-1} \equiv 1 \cdot 2 \cdots p-1 \pmod{p}.$$

But from this we immediately have $[P(X)]^2 \equiv (p-1)! \equiv -1 \pmod{p}$; hence we have $[P(X)]^2 + 1 \equiv 0 \pmod{p}$, so $p \mid [P(X)]^2 + 1^2$. As p is of the form $4k+3$, we have thus $p \mid P(X)$ and $p \mid 1$. Contradiction; hence these numbers cannot be so partitioned. \square

Problem 1.8.3: Let p be a prime. Prove that the congruence $x^2 \equiv -1 \pmod{p}$ has a solution if and only if $p = 2$ or p is of the form $4k+1$.

Proof. If $p = 2$, the conclusion is clear. If p is of the form $4k+3$ and there does exist such an x , then we have $p \mid x^2 + 1$, so $p \mid x, p \mid 1$. Contradiction; hence there are no solutions for $p = 4k+3$. Now if $p = 4k+1$, then set $U = (2k)!$. We claim that $U^2 \equiv -1 \pmod{p}$. We

write

$$\begin{aligned}
 U^2 &= 1 \cdot 2 \cdots (2k) \cdot (2k) \cdot (2k-1) \cdots 1 \\
 &\equiv 1 \cdot 2 \cdots (2k) \cdot (p-2k)(-1)(p-(2k-1))(-1) \cdots (p-1)(-1) \pmod{p} \\
 &\equiv 1 \cdot 2 \cdots (2k) \cdot (2k+1) \cdot (2k+2) \cdots (4k) \cdot (-1)^{2k} \\
 &\equiv (p-1)! \equiv -1 \pmod{p}
 \end{aligned}$$

Hence there does exist some $x = U$ with $x^2 \equiv -1 \pmod{p}$. □

Problem 1.8.4: Determine all positive integers p, m such that

$$(p-1)! + 1 = p^m.$$

Proof. Note that if $p \leq 5$, then we have the solutions $(p, m) = (2, 1), (3, 1), (5, 2)$. Now suppose that $p > 5$. Then Wilson's theorem gives the result that p must be a prime. We have $2 < (p-1)/2 < p-1$, hence $(p-1)^2 \mid (p-1)!$. Hence $(p-1)^2 \mid p^m - 1$, so $p-1 \mid p^{m-1} + p^{m-2} + \cdots + p + 1$. It follows from work in previous lectures that $p-1 \mid m$, hence $m \geq p-1$. Hence

$$p^m \geq p^{p-1} > 2 \cdot 2 \cdot 3 \cdot 4 \cdots (p-2) \cdot (p-1) = 2(p-1)! > (p-1)! + 1,$$

hence there are no solutions for $p > 5$. Thus the solutions given above are the only such p, m . □

Problem 1.8.5: Let p be an odd prime, and let $A = \{a_1, a_2, \dots, a_{p-1}\}, B = \{b_1, b_2, \dots, b_{p-1}\}$ be complete sets of nonzero residue classes modulo p - that is, if for some n we have $p \nmid n$, then there exist i, j with $n \equiv a_i \equiv b_j$. Show that the set $\{a_1 b_1, \dots, a_{p-1} b_{p-1}\}$ is not a complete set of nonzero residue classes.

Proof. We have

$$a_1 a_2 \cdots a_{p-1} \equiv 1 \cdot 2 \cdots p-1 = (p-1)! \equiv -1 \pmod{p}.$$

Similarly, $b_1 b_2 \cdots b_{p-1} \equiv -1 \pmod{p}$. Wilson's theorem implies that if any set S is a complete set of nonzero residue classes, then the product of all of its elements must be congruent to -1 modulo p . But we have

$$(a_1 b_1)(a_2 b_2) \cdots (a_{p-1} b_{p-1}) = (a_1 a_2 \cdots a_{p-1})(b_1 b_2 \cdots b_{p-1}) \equiv (-1) \cdot (-1) = 1 \pmod{p}.$$

As $p > 2$, we have $1 \not\equiv -1 \pmod{p}$. Hence $\{a_1 b_1, \dots, a_{p-1} b_{p-1}\}$ cannot be a complete set of nonzero residue classes modulo p . □

9 Problems

Some challenging problems on order.

(ISL 2000/N4) Find all solutions to $a^m + 1 \mid (a + 1)^n$.

(IMO 2000/5) Does there exist an integer n with 2000 prime divisors such that $n \mid 2^n + 1$?
+variant with squarefree

(IMO 1999/4) Solve: p prime, $x \leq 2p$, $x^{p-1} \mid (p-1)^x + 1$.

(ISL 1997) Let $b > 1, m \neq n$. If $b^m - 1$ and $b^n - 1$ have the same prime divisors then $b + 1$ is a power of 2. (In fact, stronger thing.)

(TST 2003/3) Find all ordered triples of primes (p, q, r) such that $p \mid q^r + 1$, $q \mid r^p + 1$, and $r \mid p^q + 1$.

(IMO 2003/6) Prove that for any prime p there is a prime number q that does not divide any of the numbers $n^p - p$ with $n \geq 1$.

(MOSP 2007/5.4) Given positive integers a and c and integer b , prove that there exists a positive integer x such that $a^x + x \equiv b \pmod{c}$.

1. Let p be a prime number. Find all natural numbers n such that p divides $\varphi(n)$ and such that n divides $a^{\frac{\varphi(n)}{p}} - 1$ for all positive integers a relatively prime to n .

Appendix B

Quadratic residues

1 Quadratic residues

Definition 2.1.1: Let $p > 2$ be a prime and a an integer. The **Legendre symbol** $\left(\frac{a}{p}\right)$ is defined as follows.

$$\left(\frac{a}{p}\right) = \begin{cases} 1, & \text{if } a \text{ is a square modulo } p \text{ and } p \nmid a \\ -1, & \text{if } a \text{ is not a square modulo } p \\ 0, & \text{if } p|a. \end{cases} \quad (\text{B.1})$$

Note $\left(\frac{a}{p}\right)$ is pronounced “ a on p .” If a is a square modulo p we also say a is a **quadratic residue** modulo p .

Note that $\left(\frac{a}{p}\right)$ depends only on the residue of a modulo p , so we may think of $\left(\frac{\bullet}{p}\right)$ as a map

$$\left(\frac{\bullet}{p}\right) : \mathbb{Z}/p\mathbb{Z} \rightarrow \{\pm 1\}.$$

The following offers a theoretical, although impractical, way to calculate $\left(\frac{a}{p}\right)$.

Lemma 2.1.2. For $p > 2$,

$$\left(\frac{a}{p}\right) \equiv a^{\frac{p-1}{2}} \pmod{p}.$$

Note this gives the actual value of $\left(\frac{a}{p}\right)$ since $-1 \not\equiv 1 \pmod{p}$.

Proof. The lemma clearly holds for $a \equiv 0 \pmod{p}$. Now suppose $a \not\equiv 0 \pmod{p}$. Note that $a^{\frac{p-1}{2}} \equiv \pm 1$, since $(a^{\frac{p-1}{2}})^2 \equiv 1 \pmod{p}$ by Fermat’s Little Theorem.

First suppose a is a square modulo p . Write $a \equiv b^2 \pmod{p}$. Then

$$a^{\frac{p-1}{2}} \equiv b^{p-1} \equiv 1 \pmod{p}$$

by Fermat’s Little Theorem.

Now suppose $a^{\frac{p-1}{2}} = 1$. Let g be a primitive root modulo p . Then we can write $a \equiv g^k \pmod{p}$ for some integer k . The hypothesis gives

$$g^{\frac{k(p-1)}{2}} \equiv a^{\frac{p-1}{2}} \equiv 1 \pmod{p}.$$

Since g is a primitive root, this implies $p-1 \mid \frac{k(p-1)}{2}$, i.e. k is even. Then $a \equiv (b^{\frac{k}{2}})^2$ is a square modulo p .¹ \square

As a corollary of the preceding lemma, we obtain the following multiplicative property.

Proposition 2.1.3: For any integers a and b and any prime $p > 2$,

$$\left(\frac{ab}{p}\right) = \left(\frac{a}{p}\right) \left(\frac{b}{p}\right).$$

In other words,

$$\left(\frac{\bullet}{p}\right) : (\mathbb{Z}/p\mathbb{Z})^\times \rightarrow \{\pm 1\}$$

is a group homomorphism.

Proof. By Lemma 2.1.2,

$$\left(\frac{ab}{p}\right) = (ab)^{\frac{p-1}{2}} = a^{\frac{p-1}{2}} b^{\frac{p-1}{2}} = \left(\frac{a}{p}\right) \left(\frac{b}{p}\right).$$

The second statement follows from the first and the fact that $\left(\frac{1}{p}\right) = 1$. \square

This means that to calculate $\left(\frac{a}{p}\right)$, we can factor a into primes

$$a = q_1^{\alpha_1} \cdots q_n^{\alpha_n}$$

and find that

$$\left(\frac{a}{p}\right) = \left(\frac{q_1}{p}\right)^{\alpha_1} \cdots \left(\frac{q_n}{p}\right)^{\alpha_n},$$

so it remains to find an easy way to evaluate $\frac{q}{p}$ where both p and q are prime. We do this in the next section.

¹ Alternatively, we can avoid the use of primitive roots as follows. In the first part we've shown that

$$\{a \mid a^{\frac{p-1}{2}} \equiv 1 \pmod{p}\} \subseteq (\mathbb{Z}/p\mathbb{Z})^{\times 2}.$$

The set on the LHS has $\frac{p-1}{2}$ elements. Indeed, $x^{p-1} - 1 = 0$ splits completely modulo p and has distinct roots, namely $1, \dots, p-1$ by Fermat's little theorem. Then $x^{\frac{p-1}{2}} - 1$, as a factor of $x^{p-1} - 1$, must have $\frac{p-1}{2}$ distinct roots.

It suffices to show the set on the RHS has at most $\frac{p-1}{2}$ elements. This is true since for every a , a^2 and $(-a)^2$ are equal. Hence there are at most $\frac{p-1}{2}$ nonzero squares modulo p , namely $1^2, \dots, \left(\frac{p-1}{2}\right)^2$.

2 Quadratic reciprocity

Quadratic reciprocity relates $\left(\frac{p}{q}\right)$ with $\left(\frac{q}{p}\right)$, i.e., it gives a relationship between whether p is a square modulo q and whether q is a square modulo p . (See example. ADD.)

Theorem 2.2.1 (Quadratic reciprocity). *Let $p \neq q$ be odd primes. Then*

$$\left(\frac{p}{q}\right) \left(\frac{q}{p}\right) = (-1)^{\frac{p-1}{4} \frac{q-1}{4}}.$$

In other words,

$$\left(\frac{q}{p}\right) = \begin{cases} -\left(\frac{p}{q}\right), & p \equiv q \equiv 3 \pmod{4} \\ \left(\frac{p}{q}\right), & \text{otherwise.} \end{cases}$$

For the prime 2, or when $p = -1$, we use the following instead.

Theorem 2.2.2 (Complement to quadratic reciprocity). *Let p be an odd prime. Then*

$$\begin{aligned} \left(\frac{-1}{p}\right) &= (-1)^{\frac{p-1}{2}} \\ \left(\frac{2}{p}\right) &= (-1)^{\frac{p^2-1}{8}}. \end{aligned}$$

In other words,

$$\begin{aligned} \left(\frac{-1}{p}\right) &= \begin{cases} 1, & p \equiv 1 \pmod{4} \\ -1, & p \equiv 3 \pmod{4}. \end{cases} \\ \left(\frac{2}{p}\right) &= \begin{cases} 1, & p \equiv \pm 1 \pmod{8} \\ -1, & p \equiv \pm 3 \pmod{8}. \end{cases} \end{aligned}$$

We know $\left(\frac{a}{p}\right) \equiv a^{\frac{p-1}{2}} \pmod{p}$ by Lemma 2.1.2. To prove quadratic reciprocity, we first find an alternate way to express $a^{\frac{p-1}{2}}$.

Lemma 2.2.3 (Gauss's lemma). *For an integer a and an odd prime p , define the **least residue** of a modulo p , denoted $LR_p(a)$, to be the element $b \in \left(-\frac{p}{2}, \frac{p}{2}\right)$ such that*

$$a \equiv b \pmod{p}.$$

(In other words, $LR_p(a)$ is the integer of smallest absolute value congruent to a modulo p .)

Let μ be the number of elements of $\{ka \mid 1 \leq k \leq \frac{p-1}{2}\}$ such that $LR_p(ka) < 0$. Then

$$a^{\frac{p-1}{2}} \equiv (-1)^\mu \pmod{p}$$

Hence,

$$\left(\frac{a}{p}\right) = (-1)^\mu.$$

Proof. We calculate the product

$$a \cdot 2a \cdots \left(\frac{p-1}{2} \cdot a \right)$$

modulo p in two ways.

First, combining powers of a we get

$$a \cdot 2a \cdots \left(\frac{p-1}{2} \cdot a \right) \equiv a^{\frac{p-1}{2}} \left(\frac{p-1}{2} \right)! \pmod{a}. \quad (\text{B.2})$$

Secondly, reducing each factor to the least residue first gives

$$\begin{aligned} a \cdot 2a \cdots \frac{p-1}{2} \cdot a &\equiv \text{LR}_p(a) \cdot \text{LR}_p(2a) \cdots \text{LR}_p\left(\frac{p-1}{2}\right) \\ &\equiv (-1)^\mu |\text{LR}_p(a)| \cdots \left| \text{LR}_p\left(\frac{p-1}{2}\right) \right| \\ &\equiv (-1)^\mu \left(\frac{p-1}{2} \right)! \pmod{p}. \end{aligned} \quad (\text{B.3})$$

In the last step we used the fact that $|\text{LR}_p(a)|, \dots, \left| \text{LR}_p\left(\frac{p-1}{2}\right) \right|$ is a permutation of $1, \dots, \frac{p-1}{2}$. To see this, note $-\text{LR}_p(m) = \text{LR}_p(-m)$, so

$$\begin{aligned} \left\{ \pm \text{LR}_p(ka) : 1 \leq k \leq \frac{p-1}{2} \right\} &= \{ \pm \text{LR}_p(ka) : 1 \leq k \leq p-1 \} \\ &= \left\{ -\frac{p-1}{2}, \dots, -1, 1, \dots, \frac{p-1}{2} \right\}. \end{aligned}$$

Hence $\{ \pm \text{LR}_p(ka) : 1 \leq k \leq \frac{p-1}{2} \}$ must contain one element from each pair $\pm 1, \dots, \pm \frac{p-1}{2}$, as needed.

Equating (B.2) and (B.3) and cancelling $\left(\frac{p-1}{2}\right)!$ gives the desired result.

The second statement follows because $1 \not\equiv -1 \pmod{p}$. □

Now we prove quadratic reciprocity.

Proof of Theorem 2.2.1. The strategy is as follows.

1. Establish a correspondence between $x \in \left(0, \frac{q}{2}\right)$ such that $\text{LR}_p(xq) < 0$, with lattice points in a certain region (B.10). Similarly establish such a correspondence with $y \in \left(0, \frac{p}{2}\right)$ such that $\text{LR}_q(yq) < 0$. By Lemma 2.2.3, $\left(\frac{q}{p}\right) \left(\frac{p}{q}\right)$ is the total number of lattice points in this region.
2. Pair up the points in the region. We will find that there is an odd point out exactly when $p \equiv q \equiv 3 \pmod{4}$.

Let

$$\begin{aligned}\mu_1 &= \left| \left\{ x \in \left(0, \frac{q}{2}\right) : \text{LR}_q(xp) \right\} \right| \\ \mu_2 &= \left| \left\{ y \in \left(0, \frac{p}{2}\right) : \text{LR}_p(yq) \right\} \right|\end{aligned}$$

By Lemma 2.2.3,

$$\left(\frac{p}{q}\right) \left(\frac{q}{p}\right) = (-1)^{\mu_1} (-1)^{\mu_2} = (-1)^{\mu_1 + \mu_2}. \quad (\text{B.4})$$

We would like to know the parity of $\mu_1 + \mu_2$.

Claim 2.2.4. *There is a bijection between integers $x \in (0, \frac{q}{2})$ satisfying $\text{LR}_q(xp) < 0$ and lattice points (x, y) satisfying*

$$0 < x < \frac{q+1}{2} \quad (\text{B.5})$$

$$0 < y < \frac{p+1}{2} \quad (\text{B.6})$$

$$-\frac{q}{2} < xp - yq < 0. \quad (\text{B.7})$$

Proof. If (x, y) satisfies the above inequalities, then inequality (B.7) gives that the least residue of xp is in $(-\frac{q}{2}, 0)$.

Conversely, given such a x , choose y so that yq is the closest multiple of q to xp . Then $\text{LR}_q(xp) = xp - yq$, so inequality (B.7) follows. Moreover, this is the only value of y that will satisfy (B.7). Then (B.6) follows since (B.7) gives

$$0 < \frac{p}{q}x < y < \frac{p}{q}x + \frac{1}{2} < \frac{p}{q} \cdot \frac{q}{2} + \frac{1}{2} = \frac{p+1}{2}.$$

□

Note (B.7) is equivalent to

$$\frac{p}{q}x < y < \frac{p}{q}x + \frac{1}{2}. \quad (\text{B.8})$$

Applying the claim with p and q switched, and x and y switched, inequality (B.7) becomes $-\frac{p}{2} < yq - xp < 0$, which rearranges to

$$\frac{p}{q}x - \frac{p}{2q} < y < \frac{p}{q}x. \quad (\text{B.9})$$

Noting that there are no points on the line $y = \frac{p}{q}x$ in the following region, we see that $\mu_1 + \mu_2$ equals the number of lattice points in the region \mathcal{R} defined by

$$\begin{aligned}0 < x < \frac{q+1}{2} \\ 0 < y < \frac{p+1}{2} \\ \frac{p}{q}x - \frac{p}{2q} < y < \frac{p}{q}x + \frac{1}{2}.\end{aligned} \quad (\text{B.10})$$

This region is symmetric around the point $(\frac{q+1}{4}, \frac{p+1}{4})$. Indeed, making the change of variables $x' = x - \frac{q+1}{4}$ and $y' = y - \frac{p+1}{4}$, we get

$$\begin{aligned} -\frac{q+1}{4} < x' < \frac{q+1}{4} \\ -\frac{p+1}{4} < y' < \frac{p+1}{4} \\ \frac{p}{q}x' - \left(\frac{p}{4q} + \frac{1}{4}\right) < y' < \frac{p}{q}x' + \left(\frac{p}{4q} + \frac{1}{4}\right). \end{aligned}$$

Hence we can pair up the lattice points in \mathcal{R} by matching (x, y) with $(\frac{q+1}{2} - x, \frac{p+1}{2} - y)$ (this corresponds to $(x', y') \leftrightarrow (-x', -y')$). The only point which would not be paired up is $(\frac{q+1}{4}, \frac{p+1}{4})$, but this is an integer if and only if $p \equiv q \equiv 3 \pmod{4}$. Thus $\mu_1 + \mu_2$ is odd iff $p \equiv q \equiv 3 \pmod{4}$. In light of (B.4), this proves the theorem. \square

Proof of Theorem 2.2.2. The fact that $\left(\frac{-1}{p}\right) = (-1)^{\frac{p-1}{2}}$ comes directly from Proposition 2.1.2.

To calculate $\left(\frac{2}{p}\right)$, we can use Lemma 2.2.3 directly. In this case μ is the number of elements in the set $\{2, 4, \dots, p-1\}$ in the interval $(\frac{p}{2}, p)$. By casework, this is even when $p \equiv \pm 1 \pmod{8}$ and odd when $p \equiv \pm 3 \pmod{8}$. \square

Problems

1. (IMO 1996/4) The positive integers a, b are such that $15a + 16b$ and $16a - 15b$ are both squares of positive integers. What is the least possible value that can be taken by the smaller of these two squares?

3 Jacobi symbol

Appendix C

Unique factorization

1 Unique factorization domains

MOTIVATION

First, we define what exactly unique factorization means. Let R be an integral domain.

Definition 3.1.1: An element $a \in R$ is **irreducible** if it is not a unit, and its only factors are units and associates. A unit is an invertible element in R , while an associate of a is a unit times a .

For the positive integers we often just say a is irreducible if $a \neq 1$, and its only factors are 1 and itself. However, if we work with the integers, then there will also be the factors -1 and $-a$, and we don't want to view these as different. For example, 5 is irreducible over the integers because its only factors are units, ± 1 , and associates, ± 5 .

Definition 3.1.2: A **unique factorization domain (UFD)** is a integral domain where factoring terminates and every nonzero, nonunit element factors uniquely into irreducible elements. That is, if

$$a = p_1 \cdots p_m = q_1 \cdots q_n,$$

and $p_1, \dots, p_m, q_1, \dots, q_n$ are irreducible elements, then $m = n$ and we can reorder the q_i 's so that p_i is an associate of q_i , for each i .

For example, we regard $6 = 2 \cdot 3 = -2 \cdot -3$ as the same factorization.

Unique factorization doesn't hold for all domains—for example, consider $\mathbb{Z}[\sqrt{-5}]$, that is, numbers of the form $a + b\sqrt{-5}$. Then

$$(1 + \sqrt{-5})(1 - \sqrt{-5}) = 6 = 2 \cdot 3$$

are two factorizations of 6 into irreducible elements.

The notion of a prime is related to that of an irreducible element. People use them as synonyms in elementary math—because they coincide for the integers—but the distinction between them will be quite important for us.

Definition 3.1.3: A prime in R is an element p , not a unit, such that if $p|ab$ then $p|a$ or $p|b$.

This tells us that if a prime p divides a , then *no matter how we factor a* , we can't avoid p dividing one element of a . The connection between primes, irreducibles, and unique factorization is given by the following.

Lemma 3.1.4. *If R is a ring where factoring terminates, and every irreducible element is prime, then R is a UFD. Conversely, in a UFD, every irreducible element is prime.*

Proof. Suppose $a = p_1 \dots p_m = q_1 \dots q_n$ are two factorizations into irreducible elements. Since p_1 is irreducible, it is prime, and hence must divide one of the q_i . Since q_i is irreducible, its only factors are units and associates, so p_1 must be associated with q_i . Then we can cancel them, leaving a unit. Repeating this process, every factor in the left factorization is paired with one in the right factorization.

For the converse, suppose p is irreducible and $p|ab$. Then $pd = ab$ for some d . Factoring a , b , and d shows that p must divide one of the factors of a or b by unique factorization. \square

(Note that primes are always irreducible, because if $p = ab$ were a proper factorization, then $p \nmid a$ and $p \nmid b$.)

The main strategy for proving unique factorization is the following.

1. Show that the ring R in question (here, $K[x]$) admits **division with remainder**, with some measure of size so that the remainder is smaller than the quotient.
2. Show that if we have division of remainder, then **greatest common divisors** exist, and moreover that they have the nice property given by Bézout's Theorem.
3. Show that this implies that all irreducible elements are prime, and hence R is a UFD.

The advantage of such an abstract approach lies in the fact that it works for a variety of different number systems. In particular, once we've shown items 2 and 3, then given any ring, we only have to show that we can have division with remainder, and it will follow that it is a UFD. This simultaneously shows unique factorization for \mathbb{Z} , $K[x]$, and even $\mathbb{Z}[i] = \{a + bi \mid a, b \in \mathbb{Z}\}$.¹

In the language of abstract algebra, the above steps are phrased as follows:

1. R is an Euclidean domain.
2. An Euclidean domain is a principal ideal domain.
3. A principal ideal domain is a unique factorization domain.

We now carry out this program.

¹The converse is not true; a UFD is not necessarily a PID or Euclidean domain. For example $\mathbb{Z}[\frac{1+\sqrt{-163}}{2}]$ is a UFD but not an Euclidean domain.

1.1 Step 1: Euclidean domains

Definition 3.1.5: An integral domain R is an **Euclidean domain** if there is a function $|\cdot| : R \rightarrow \mathbb{N}_0$ (called the norm) such that the following hold.

1. $|a| = 0$ iff $a = 0$.
2. For any nonzero $a, b \in R$ there exist $q, r \in R$ such that $b = aq + r$ and $|r| < |a|$.

Note that both the integers \mathbb{Z} and $K[x]$ are Euclidean domains. The norm on \mathbb{Z} is simply the absolute value, while the norm on $K[x]$ is the degree of the polynomial. Theorem 4.2.1 shows that $K[x]$ is an Euclidean domain.

1.2 Step 2: Euclidean domain \implies PID

We'd like to prove Bézout's Theorem for an Euclidean domain, that given a, b in R there exists a greatest common divisor g and s, t so that $as + bt = g$. Rather than thinking of this as an equation in variables s, t , we can think of it as an equation in sets (a) and (b) , where (x) denotes the set of multiples of x . For two sets S, T we define $S + T = \{s + t \mid s \in S, t \in T\}$; then it turns out what we want is

$$(a) + (b) = (g).$$

(See Lemma 3.1.8 below.)

Definition 3.1.6: An **ideal** in a ring R is a subset I such that if $a, b \in I$ then $ra, a + b \in I$ for any $r \in R$. A principal ideal is an ideal generated by one element, that is, there is a a such that $I = \{ra \mid r \in R\}$. We write $I = (a)$.

A **principal ideal domain** (PID) is an integral domain where every ideal is principal.

Theorem 3.1.7. *An Euclidean domain is a PID.*

Proof. Let R be an Euclidean domain, $I \subseteq R$ an ideal, and b be the nonzero element of smallest norm in I . Suppose $a \in I$. Then we can write $a = qb + r$ with $0 \leq r < |b|$, but since b has minimal nonzero norm, $r = 0$ and $b \mid a$. Thus $I = (b)$ is principal. \square

Lemma 3.1.8. *A PID satisfies Bézout's Theorem.*

Proof. Let R be a PID. Since every ideal in R is principal, for every a, b (not both 0) we have $(a) + (b) = (d)$ for some $d \in R$. (Note the sum of two ideals is an ideal—check this for yourself.) This says there exist $s, t \in R$ such that

$$as + bt = d.$$

From this, any divisor of a, b must divide d . Furthermore, d must divide both a and b since $a = a + 0$ and $b = 0 + b$ are both in $(a) + (b) = (d)$. In other words, d is the greatest common divisor of a, b . \square

1.3 Step 3: PID \implies UFD

Theorem 3.1.9. *A PID is a UFD.*

Proof. Suppose p is irreducible; we show p is prime. Suppose $p|ab$ but p does not divide a . Then using Bezout's Theorem and the fact that a and p are relatively prime, we get $as + pt = 1$ for some s, t . Multiply by b to get

$$abs + ptb = b.$$

Since $p|ab|abs, p|ptb$, we have $p|b$. This shows that irreducible elements are prime in \mathbb{Z} .

It remains to show factoring terminates.² Otherwise, there would be an infinite sequence of nonassociated elements $a_1, a_2, \dots \in R$ such that $a_{i+1}|a_i$. Then $(a_1) \subset (a_2) \subset \dots$. However, $\bigcup_{i \geq 1} (a_i)$ is an ideal, so it is principal, say generated by b . Then $b \in (a_i)$ for some i ; this implies that $(b) = (a_i)$. Hence $(a_i) = (a_{i+1}) = \dots$, a contradiction.

Since irreducible elements are prime and every nonzero element of R factors into irreducibles, R is a UFD. \square

Corollary 3.1.10. *\mathbb{Z} and $K[x]$ are UFDs.*

2 Example: $x^2 + y^2 = n$

Theorem 3.2.1. *Let n be a positive integer. Then the equation*

$$x^2 + y^2 = n$$

has a solution in integers iff every prime $p \equiv 3 \pmod{4}$ appears in n with even exponent.

If $n = 2^a p_1^{b_1} \cdots p_k^{b_k} q_1^{c_1} \cdots q_m^{c_m}$ where p_j and q_j are primes congruent to 1, 3 modulo 4, then the equation $x^2 + y^2 = n$ has

$$4(b_1 + 1) \cdots (b_k + 1)$$

solutions in integers.

Proof. Each solution to $x^2 + y^2 = n$ corresponds to a factoring $(x + yi)(x - yi) = n$ over the Gaussian integers $\mathbb{Z}[i]$. Thus the number of solutions is the number of z such that $z\bar{z} = n$, or 4 times the number of nonassociated $z \in \mathbb{Z}[i]$ such that $z\bar{z} = n$. (Two Gaussian numbers are associated if they differ by a unit $\pm 1, \pm i$, so $x + yi, -y + xi, -x - yi, y - xi$ are considered the same.)

Now factor $n = 2^a p_1^{b_1} \cdots p_k^{b_k} q_1^{c_1} \cdots q_m^{c_m}$ where p_j and q_j are primes congruent to 1, 3 modulo 4, respectively. From knowledge of factoring in $\mathbb{Z}[i]$ we know that

1. 2 ramifies in $\mathbb{Z}[i]$, that is, it is the product of two associated primes $1 + i, 1 - i$.

² This argument is not needed for our purposes: Both \mathbb{Z} and $K[x]$ are Euclidean domains, and factoring must terminate for them because factors always have smaller norm (absolute value and degree, respectively).

2. The $p_j \equiv 1 \pmod{4}$ split, that is, $p_j = z_j \bar{z}_j$ where z is prime in $\mathbb{Z}[i]$ and not associated to \bar{z} .
3. The $q_j \equiv 3 \pmod{4}$ remain prime.

Now if $z\bar{z} = n$ and a Gaussian prime divides z , then its conjugate must divide \bar{z} . Thus, since we have unique factorization in $\mathbb{Z}[i]$, each such z , up to multiplication by associates, corresponds to a way of splitting the prime factors of n into complex conjugate pairs. We note the following:

1. The factors q_j are their own conjugates, so z and \bar{z} must each get $q_j^{c_j/2}$. If one of the c_j is odd there is no solution. So we suppose they are all even.
2. It doesn't matter how the prime factors of 2^a are split since they are all associates.
3. There are $b_j + 1$ ways to split the factors of $q_j^{b_j}$, since we can have either $z_j^{b_j}$, or $z_j^{b_j-1} \bar{z}_j, \dots$ or $\bar{z}_j^{b_j}$ divide z . Thus there are $(b_1 + 1) \cdots (b_k + 1)$ solutions to $z\bar{z} = n$ up to associates.

□

A similar argument works for the equations $x^2 + 2y^2 = n$ and $x^2 + xy + y^2 = n$.

3 Problems

1. (The power of ideals) We rephrase some earlier results that used the “division with remainder argument” in terms of ideals.
 - (a) Let $n > 1$ and a be relatively prime to n . Show that

$$\{m : a^m \equiv 1 \pmod{n}\}$$
 is an ideal.
 - (b) Conclude Proposition [A.1.2.2](#).
2. For which n does $x^2 + 2y^2 = n$ have a solution? How many solutions are there? How about $x^2 + xy + y^2 = n$?

Appendix D

Polynomials

1 Gauss's argument

We've shown that $K[x]$ is a UFD, but the argument above does not show that $\mathbb{Z}[x]$ is a UFD, because division with remainder fails for $\mathbb{Z}[x]$. We will need a further argument. The basic idea is that a polynomial factors in $\mathbb{Z}[x]$ the same way it does in $\mathbb{Q}[x]$, except with its factors adjusted by constants so the coefficients are in \mathbb{Z} .

Let R be a UFD and let K be the field of fractions of R . That is, K consists of the numbers $\frac{a}{b}$ where $a, b \in R$ and $b \neq 0$, and we say $\frac{a}{b} = \frac{c}{d}$ iff $ad = bc$. For example, \mathbb{Q} is the field of fractions for \mathbb{Z} .

Definition 4.1.1: A nonzero polynomial $f \in R[x]$ is said to be **primitive** if all its coefficients do not have a common proper divisor; equivalently, there does not exist a prime $p \in R$ such that $p|f$.

Lemma 4.1.2. *If R is an integral domain, then so is $R[x]$.*

Proof. Take any $p, q \in R[x]$ not equal to 0. We can write

$$p = \sum_{i=0}^m a_i x^i, \quad a_m \neq 0$$
$$q = \sum_{i=0}^n b_i x^i, \quad b_n \neq 0$$

Then the leading coefficient of pq is $a_m b_n x^{m+n}$. It is nonzero because since R is an integral domain, $a_m, b_n \neq 0$ imply that $a_m b_n \neq 0$. Hence $pq \neq 0$. This shows that $R[x]$ is an integral domain. \square

Lemma 4.1.3 (Gauss's lemma). (A) *An element of R is prime in $R[x]$ iff it is a prime in R . Hence if a prime p of R divides a product fg of polynomials in $R[x]$, then $p|f$ or $p|g$.*
(B) *The product of primitive polynomials in $R[x]$ is primitive.*

Proof. If $p \in R$ is nonzero, and a prime in $R[x]$, then it is a prime in the subring R .

Conversely, let p be any prime element in R . Then $R/(p)$ is an integral domain¹ so by lemma 4.1.2, $R/(p)[x]$ is an integral domain.

Suppose $p|fg$ for $f, g \in R[x]$. Then in $R/(p)[x]$, $\overline{fg} = \overline{f}\overline{g} = 0$. Since $R/(p)[x]$ is an integral domain, either $\overline{f} = 0$ or $\overline{g} = 0$. In other words, either $p|f$ or $p|g$ in $R[x]$. Thus p is a prime in $R[x]$.

If f, g are primitive, then $p \nmid f$ and $p \nmid g$ for all primes $p \in R$. Since p is also prime in $R[x]$, $p \nmid fg$. Hence fg is not divisible by any prime in R , and it is primitive. \square

Lemma 4.1.4. *Every nonconstant polynomial $f \in K[x]$ can be written uniquely (up to multiplication by units) in the form $f = cf_0$, where $c \in K$ and f_0 is a primitive polynomial in $R[x]$.*

Proof. Each coefficient a_i of f is in the form $\frac{p_i}{q_i}$, where $p_i, q_i \in R$. We can find a nonzero $t \in R$ such that t is divisible by each denominator (for, example, take t to be the product of the denominators). Then we can write

$$tf = f_1,$$

where $f_1 \in R[x]$. Let $s \in R$ be a greatest common divisor of the coefficients of f_1 . Then we have

$$f = \frac{s}{t}f_0$$

in $K[x]$ where $f_0 \in R[x]$ and the coefficients of f_0 have no common divisor. This gives the desired representation.

Next we check uniqueness. Suppose

$$f = cf_0 = c'f'_0,$$

where $c, c' \in K$ and $f_0, f'_0 \in R[x]$ are primitive. Multiply by an element of R to “clear denominators,” to reduce to the case where $c, c' \in R$. Now take any prime $p|c$. Since p is prime in $R[x]$, $p|c'$ or $p|f'_0$. The second is impossible since f'_0 is primitive. Hence $p|c'$, and we can cancel p . Continuing in this way, we get that c and c' share the same prime factors with the same multiplicities. Hence c, c' are associates. \square

Lemma 4.1.5. *Let f_0 be a primitive polynomial and let $g \in R[x]$. If $f_0|g$ in $K[x]$ then $f_0|g$ in $R[x]$.*

¹ If I is an ideal, then R/I is the quotient ring: Two elements a, b in R are considered to be the same in R/I if they differ by an element in I . Keep in mind the example $R = \mathbb{Z}$; then $R/(p)$ is simply the integers modulo p .

Now $R/(p)$ is an integral domain, because if $ab = 0$ in $R/(p)$, then $ab \in (p)$, i.e. p divides one of a, b . But since p is prime either $p|a$ or $p|b$, which translates back into $a = 0$ or $b = 0$ in $R/(p)$.

Proof. If $f_0|g$ in $K[x]$, then we can write $g = f_0h$ where $h \in K[x]$. We need to show $h \in R[x]$. By lemma 4.1.4, we can write $h = ch_0$, where $c \in K$ and h_0 is primitive. Then $g = cf_0h_0$. By lemma 4.1.3, the product f_0h_0 of primitive polynomials is primitive. We can write $c = \frac{s}{t}$, where $s, t \in R$ have no common factors. If a prime p in R divides the denominator t then $p \nmid s$ so $p|f_0h_0$, contradicting the fact that f_0h_0 is primitive. Hence t is a unit, and $c \in R$. Then $h = ch_0 \in R[x]$, so $f_0|g$ in $R[x]$. □

Lemma 4.1.6. *Let f be a nonzero element of $R[x]$. Then f is an irreducible element of $R[x]$ iff it is an irreducible element of R or a primitive irreducible polynomial in $K[x]$.*

Proof. If $f \in R$, then the only factors of f in $R[x]$ are in R , so f is irreducible in R iff it is irreducible in $R[x]$. This proves the lemma for $f \in R$. Now suppose $f \notin R$.

If $f \in R[x]$ is a primitive polynomial irreducible in $K[x]$, then it is irreducible in $R[x]$.

If $f \in R[x]$ is not primitive, then it is reducible in $R[x]$. Thus it suffices to show if $f \in R[x]$ is reducible in $K[x]$, then it is reducible in $R[x]$. Suppose $f \in R[x]$, and $f = gh$ is a proper factorization of f in $K[x]$. We can write $g = cg_0, h = c'h_0$ where $c, c' \in K$ and g_0, h_0 are primitive. Since g_0 and h_0 are both primitive, so is g_0h_0 . Then $f = cc'(g_0h_0)$, so by uniqueness in lemma 4.1.4, cc' must be in R (and is the gcd of the coefficients of f). Thus $f = (cc')g_0h_0$ is a proper factorization of f in $R[x]$ as well, as needed. □

Theorem 4.1.7. *The ring $R[x]$ is a unique factorization domain.*

Proof. It suffices to show that every irreducible element f of $R[x]$ is a prime element, and that factoring terminates. By Lemma 4.1.6, f is either irreducible in R or a primitive irreducible polynomial in $K[x]$. In the first case f is prime in R (R is a UFD) and hence prime in $R[x]$ by Lemma 4.1.3.

In the second case, f is primitive irreducible in $K[x]$, thus a prime in $K[x]$, since $K[x]$ is a UFD. Hence $f|g$ or $f|h$ in $K[x]$. By Lemma 4.1.5, $f|g$ or $f|h$ in $R[x]$. This shows f is prime.

A polynomial $f \in R[x]$ can only be the product of at most $\deg(f)$ many polynomials p_i of positive degree in $R[x]$ because the sum of the degrees of the p_i must equal $\deg(f)$. Factor terminates for the factors of f in R because factoring terminates in the UFD R , and the primes in R dividing f are the primes dividing every coefficient of f .

Hence $R[x]$ is a UFD. □

Corollary 4.1.8. $\mathbb{Z}[x]$ is a UFD.

If R is a UFD then $R[x_1, \dots, x_n]$ is a UFD.

Proof. Since \mathbb{Z} is a UFD, so is $\mathbb{Z}[x]$. The second statement follows from Theorem 4.1.7 by induction. □

1.1 More Proofs

Theorem 4.1.9 (Chinese Remainder Theorem). *If polynomials $Q_1, \dots, Q_n \in K[x]$ are pairwise relatively prime, then the system $P \equiv R_i \pmod{Q_i}, 1 \leq i \leq n$ has a unique solution modulo $Q_1 \cdots Q_n$.*

Proof. Let $Q = Q_1 \cdots Q_n$. Note Q_i and $\frac{Q}{Q_i}$ are relatively prime. Hence by Bézout's Theorem there exist f_i and g_i so that

$$f_i Q_i + g_i \frac{Q}{Q_i} = 1.$$

Now

$$(1 - q_i f_i) R_i = R_i g_i \frac{Q}{Q_i}$$

is congruent to R_i modulo Q_i , and zero modulo Q_j for $j \neq i$. Hence

$$P = \sum_{i=1}^n (1 - q_i f_i) R_i$$

is the desired polynomial.

For uniqueness, suppose P_1 and P_2 satisfy the conditions of the problem. Then $P_1 - P_2$ is zero modulo Q_i . Since the Q_i are pairwise relatively prime, $P_1 - P_2 \equiv 0 \pmod{Q_1 \cdots Q_n}$. \square

Theorem 4.1.10 (Rational Roots Theorem). *Suppose that R is a UFD and K its fraction field. (For instance, take $R = \mathbb{Z}$ and $K = \mathbb{Q}$.) Suppose $f(x) = a_n x^n + \cdots + a_0 \in R[x]$ and $a_n \neq 0$. Then all roots of f in K are in the form*

$$\frac{\text{factor of } a_0}{\text{factor of } a_n}.$$

In particular, if $a_n = \pm 1$, then all roots of f in K are actually in R .

Proof. Write $x = \frac{r}{s}$ in simplest terms. Then multiplying through by s^n gives

$$a_n \left(\frac{r}{s}\right)^n + \cdots + a_1 \left(\frac{r}{s}\right) + a_0 = 0$$

$$a_n r^n = -s(a_{n-1} r^{n-1} + \cdots + a_1 r s^{n-2} + a_0 s^{n-1}).$$

Since s and r have no common factor, s must divide a_n . (This uses the fact that R is a UFD—how?). Rewriting as

$$a_0 s^n = -r(a_n r^{n-1} + \cdots + a_1 s^{n-1})$$

makes it clear r divides a_0 . \square

Remark 4.1.11: In particular, if $a_n = 1$, then all roots of f in K are in R . A ring is said to be normal if whenever $t \in K$ is a root of a monic polynomial in $R[x]$, then $t \in R$. Thus the above shows that UFDs are normal.

1.2 Problems

1. (Bézout bound) Let $f(x, y), g(x, y) \in \mathbb{C}[x, y]$. Prove that either f, g have a constant nonzero factor, or they have finitely many zeros (x, y) in common. (Hard: They have at most $\deg(f)\deg(g)$ common zeros.)
2. For a field K , let $K(x)$ be the field of rational functions, that is,

$$K(x) = \left\{ \frac{p}{q} \mid p, q \in K[x] \right\}.$$

Let f and g be rational functions such that $f(g(x)) = x$. Prove that f and g are both in the form $\frac{ax+b}{cx+d}$ with $ad \neq bc$.

In many ways, polynomials are similar to integers. Like integers, polynomials admit division with remainder, existence of greatest common divisors, and unique factorization.

2 Main Theorems

In this section K will stand for \mathbb{C} (the complex numbers), \mathbb{R} (the real numbers), \mathbb{Q} (the rational numbers), or $\mathbb{Z}/p\mathbb{Z}$ (the integers modulo p), while R will stand for any one of the before sets or \mathbb{Z} (the integers). Note that the sets we label with K all have multiplicative inverses, i.e. are *fields*.

Our first result is that when we divide polynomials, we can be assured to get a remainder with degree smaller than our divisor.

Theorem 4.2.1 (Division with remainder). *If $f, g \in K[x]$, then there exist polynomials $q, r \in K[x]$ such that $\deg r < \deg g$ and*

$$f = qg + r.$$

If $f, g \in \mathbb{Z}[x]$ and g is monic, then there exist $q, r \in \mathbb{Z}[x]$ such that $\deg r < \deg g$ and

$$f = qg + r.$$

Proof. This is the division algorithm familiar from high school algebra class. Namely, if f has leading term ax^n and g has leading term bx^m with $n \geq m$, then $f - \frac{a}{b}x^{n-m}g$ has degree less than f . Thus we can keep subtracting multiples of g from f until the result has degree less than $\deg g$.

If g is monic, then $b = 1$ so at each stage we subtracted an integer polynomial multiple of g , and both the quotient q and the remainder r will have integer coefficients. \square

Theorem 4.2.2 (Bézout). *Given $f, g \in R[x]$, there exists a polynomial h , called the **greatest common divisor** and denoted $\gcd(f, g)$, such that the following hold:*

1. h divides both f and g .

2. If p divides both f and g then p divides h .

Let $f, g \in K[x]$. There exist polynomials $u, v \in K[x]$ so that $uf + vg = \gcd(f, g)$.

(Note that h is only determined up to a unit. We'll "sweep this under the rug" and allow any choice of h up to that constant.)

To calculate the gcd, we often use the Euclidean algorithm. Given polynomials f and g , for any polynomial q we have

$$\gcd(f, g) = \gcd(g, f - qg).$$

Supposing $\deg f \geq \deg g$, take q so that $f - qg = r$ has degree less than g , as in the division algorithm; this reduces the degree of f . Repeating this process decreases the degrees of the polynomials; we eventually get to $\gcd(h, 0)$ in which case the answer is seen to be h .

Theorem 4.2.3 (Unique factorization). *Every polynomial in $R[x]$ factors uniquely in $R[x]$, up to constants. In fact, every polynomial in $R[x_1, \dots, x_n]$ factors uniquely in $R[x_1, \dots, x_n]$, up to constants.*

We give two more useful results.

Theorem 4.2.4 (Chinese Remainder Theorem). *If polynomials $Q_1, \dots, Q_n \in K[x]$ are pairwise relatively prime, then the system $P \equiv R_i \pmod{Q_i}, 1 \leq i \leq n$ has a unique solution modulo $Q_1 \cdots Q_n$.*

Theorem 4.2.5 (Rational Roots Theorem). *Suppose $f(x) = a_n x^n + \cdots + a_0$ is a polynomial with integer coefficients and with $a_n \neq 0$. Then all rational roots of f are in the form*

$$\frac{\text{factor of } a_0}{\text{factor of } a_n}.$$

In particular, if $a_n = \pm 1$, then all rational roots of f are integers.

Here's a cute application of Bézout's Theorem:

Example 4.2.6: Let f, g be polynomials with integer coefficients and with no common factor. Prove that $\gcd(f(n), g(n)), n \in \mathbb{Z}$ can only attain a finite number of values.

Solution. By Bézout's Theorem, we have $u(x)f(x) + v(x)g(x) = 1$ for some $u, v \in \mathbb{Q}[x]$ and nonzero. Clearing denominators of u and v , we get $u'(x)f(x) + v'(x)g(x) = k$ for some $u', v' \in \mathbb{Z}[x]$ and nonzero $k \in \mathbb{Z}$. Hence $\gcd(f(n), g(n)) \mid k$.

2.1 Problems

- [1] Show by example we cannot always carry out division with remainder in $\mathbb{Z}[x]$ and that Bézout's Theorem does not hold for $\mathbb{Z}[x]$.

2. [1] Compute the greatest common divisors in $\mathbb{Z}[x]$:

(a) $\gcd(x^6 - x^5 - x^2 + 1, x^3 - 2x^2 + 2x - 1)$.

(b) $\gcd(x^{12} - 1, x^8 + 1)$.

3. Find the greatest common divisor in $\mathbb{Z}[x]$:

(a) [2] $\gcd(x^n - 1, x^m - 1)$.

(b) [2.5] $\gcd(x^n + 1, x^m + 1)$.

Are your answers the same if we work in $(\mathbb{Z}/p\mathbb{Z})[x]$?

4. [1.5] Let $n > 0$ be an integer. Find the remainder upon division of $x^n + x^{n-1} + \cdots + 1$ by:

(a) $x^2 + 1$.

(b) $x^2 + x + 1$.

(c) $x^2 - x + 1$.

5. [2.5] Let f, g be relatively prime polynomials with integer coefficients. Prove that there exist nonzero polynomials u, v with integer coefficients such that $uf + vg = k$ where k is a nonzero integer.

Suppose that $u_1f + v_1g = k_0$ and u_1, v_1 are integer polynomials with $u_1 = \sum_{i=0}^m a_i x^i, v_1 = \sum_{i=0}^n b_i x^i, \deg(u_1) < \deg(g), \gcd(a_0, \dots, a_m, b_0, \dots, b_n) = 1$. Prove that $k_0 \mid k$.

6. [3] Let $f : \mathbb{Q} \rightarrow \mathbb{Q}$ satisfy $f(f(f(x))) + 2f(f(x)) + f(x) = 4x$. and $f(f(\cdots f(x))) = x$ where f is taken 2009 times. Prove that $f(x) = x$.

7. [3] (BAMO 2004) Find all polynomials f with integer coefficients taking irrationals to irrationals.

8. [5] (USAMO 1997/3) Prove that for any integer n , there exists a unique polynomial Q with coefficients in $\{0, 1, \dots, 9\}$ such that $Q(-2) = Q(-5) = n$.

9. [2] For how many integers n is $\frac{n^3+1000}{n-10}$ an integer?

10. [2] Suppose that f and g are integer polynomials such that $f(n)/g(n)$ is an integer for infinitely many $n \in \mathbb{Z}$. Show that as polynomials, $g(x)$ divides $f(x)$.

11. [5] (IMO 2002/3) Find all pairs of integers $m > 2, n > 2$ such that there are infinitely many positive integers a for which $a^n + a^2 - 1$ divides $a^m + a - 1$.

3 Arithmetic Properties

In this section we concentrate on polynomials with integer coefficients. The following is a simple but very useful idea.

Theorem 4.3.1. *If P has integer coefficients, then $a - b \mid P(a) - P(b)$ for all integers a, b .*

Proof. Let $m = a - b$. Then $a \equiv b \pmod{m}$. Let $P = c_n x^n + \cdots + c_1 x + c_0$. Then

$$c_n a^n + \cdots + c_1 a + c_0 \equiv c_n b^n + \cdots + c_1 b + c_0 \pmod{m}$$

giving $P(a) \equiv P(b) \pmod{m}$, as needed. □

Here is a typical application. Note the use of the extremal principle.

Example 4.3.2 (USAMO 1974/1): $P(x)$ is a polynomial with integral coefficients. If a, b, c are integers so that $P(a) = b, P(b) = c, P(c) = a$, prove that $a = b = c$.

Proof. If not, then no two are equal. Without loss of generality, assume that c is between a and b . Then

$$|P(a) - P(b)| = |c - b| < |b - a|.$$

However, $b - a \mid P(b) - P(a)$, a contradiction. □

Example 4.3.3: Let P be a nonconstant polynomial with integer coefficients. Prove that there is an integer x so that $P(x)$ is composite.

Proof. Take n so that $P(n)$ is nonzero. Suppose it is prime. For all $k \in \mathbb{Z}$, we have $P(n) \mid P(n + kP(n)) - P(n)$, and hence $P(n) \mid P(n + kP(n))$. If $P(x)$ is not composite for any integer x , then $P(n + kP(n))$ is $\pm P(n)$ or 0 for all $k \in \mathbb{Z}$. P attains one of these values infinitely many times, so must be constant, a contradiction. □

One question we could ask is what values a polynomial can take modulo a given integer m as x ranges over the residues modulo m . (From Theorem 4.3.1 we know that the value modulo m depends only on x modulo m .) We know by the Lagrange Interpolation formula that we can manufacture a polynomial taking arbitrary values at a given set of points if we're allowed to divide—so it works for \mathbb{R}, \mathbb{Q} , and even $\mathbb{Z}/p\mathbb{Z}$. However Lagrange Interpolation will not work modulo m for m composite because in general we cannot divide modulo m (for example, 2 has no inverse modulo 4). For instance, Theorem 4.3.1 already tells us that given $P(x)$, $P(x + p)$ cannot be any residue modulo p^2 ; it can only be those residues that are congruent to x modulo p .

Example 4.3.4 (TST 2007/6): For a polynomial $P(x)$ with integer coefficients, $r(2i - 1)$ (for $i = 1, 2, 3, \dots, 512$) is the remainder obtained when $P(2i - 1)$ is divided by 1024. The sequence

$$(r(1), r(3), \dots, r(1023))$$

is called the *remainder sequence* of $P(x)$. A remainder sequence is called *complete* if it is a permutation of $(1, 3, 5, \dots, 1023)$. Prove that there are no more than 2^{35} different complete remainder sequences.

Solution. Step 1

For $i \in \mathbb{N}$, let

$$P_i(x) = \prod_{k=1}^i (x - (2k - 1)).$$

(Define $P_0(x) = 1$.) By Problem 7, any polynomial with integer coefficients can be written in the form $\sum_{0 \leq i \leq n} c_i P_i(x)$.

Step 2

Let $a_i = \sum_{k=0}^{\infty} \lfloor \frac{i}{2^k} \rfloor$. We claim that $2^{a_i} \mid P_i(x)$ for all $i \in \mathbb{N}$ and all odd x . For a prime p and $n \in \mathbb{Z}$, denote by $v_p(n)$ the exponent of the highest power of p dividing n (by convention $v_p(0) = \infty$). For given odd x let $f(\alpha)$ be the number of values of k ($0 \leq k \leq i - 1$) where $2^\alpha \mid x - 1 - 2k$. Then

$$v_2(P_i(x)) = \sum_{k=0}^{i-1} v_2(x - 1 - 2k) = \sum_{\alpha=1}^{\infty} f(\alpha)$$

since each k with $2^\alpha \parallel x - 1 - 2k$ is counted α times in either sum.

Since any set of $2^{\alpha-1}$ consecutive even integers has one divisible by 2^α , any set of i consecutive even integers has at least $\lfloor \frac{i}{2^{\alpha-1}} \rfloor$ integers divisible by 2^α . Hence $f(\alpha) \geq \lfloor \frac{i}{2^{\alpha-1}} \rfloor$, and $v_2(P_i(x)) \geq \sum_{\alpha=0}^{\infty} \lfloor \frac{i}{2^\alpha} \rfloor$ as desired.

Note $a_0 = 0, a_1 = 1, a_2 = 3, a_3 = 4, a_4 = 7, a_5 = 8$, and $a_i \geq 10$ for $i \geq 6$.

Step 3

Next, we claim that if $P(x) = \sum_{0 \leq i \leq n} c_i P_i(x)$ has a complete remainder sequence then c_1 is odd. (c_0 obviously needs to be odd.) We have $4 \mid P(4k + i) - P(i)$ for any integer i ; hence $r(4k + 1) \equiv r(1) \pmod{4}$ and $r(4k + 3) \equiv r(3) \pmod{4}$ for each k . In order for the remainder sequence to be complete, we need $r(1) \not\equiv r(3) \pmod{4}$. But noting that $a_i \geq 2$ and $P_i(x) \equiv 0 \pmod{4}$ for odd x and $i \geq 2$, we have $P(3) - P(1) \equiv c_1(P_1(3) - P_1(1)) \equiv 2c_1 \pmod{4}$. Hence c_1 is odd.

Step 4

Since for any odd x , $P_i(x)$ is divisible by 2^{a_i} , if we mod out c_i by 2^{10-a_i} , and delete the terms with P_i for $i \geq 6$ (where $a_i \geq 10$), we get a polynomial with the same remainder sequence as P_i . If $P(x)$ gives a complete remainder sequence, then c_0 is odd, so there are 2^9 choices for it; c_1 is odd, so there are at most 2^8 choices for $c_1 \pmod{2^9}$ ($a_1 = 1$); for $2 \leq i \leq 5$ there are at most 2^{10-a_i} choices for $c_i \pmod{2^{10-a_i}}$. Hence the number of complete remainder sequences is at most

$$2^9 \cdot 2^8 \cdot \prod_{i=2}^5 2^{10-a_i} = 2^9 \cdot 2^8 \cdot 2^7 \cdot 2^6 \cdot 2^3 \cdot 2^2 = 2^{35}.$$

□

Rather than ask about polynomials with integer coefficients, we could ask about polynomials with integer values, that is P such that $P(n)$ is an integer whenever n is an integer. It turns out that there is a nice description of such polynomials, as the following example shows.

Theorem 4.3.5. *Let $f(x) \in \mathbb{C}[x]$. Then the following are equivalent:*

- a. For every $x \in \mathbb{Z}$, $f(x) \in \mathbb{Z}$.
- b. For $n + 1$ consecutive integers x , where n is the degree of f , $f(x) \in \mathbb{Z}$.
- c. There are $a_0, a_1, \dots, a_n \in \mathbb{Z}$ with

$$f(x) = a_n \binom{x}{n} + a_{n-1} \binom{x}{n-1} + \cdots + a_0 \binom{x}{0}.$$

Here $\binom{x}{n}$ is defined as

$$\frac{x^n}{n!} = \frac{x(x-1)\cdots(x-(n-1))}{n!}$$

Proof. The assertions (a) \Rightarrow (b) and (c) \Rightarrow (a) are clear ($\binom{x}{i}$ are integers for all integers x and nonnegative integers i , by combinatorial argument).

Suppose (b) holds. First assume that $f(x)$ takes on integer values at $0, 1, \dots, n$. We inductively build the sequence a_0, a_1, \dots so that the polynomial

$$P_m(x) = a_m \binom{x}{m} + a_{m-1} \binom{x}{m-1} + \cdots + a_0 \binom{x}{0}$$

matches the value of $f(x)$ at $x = 0, \dots, m$. Define $a_0 = f(0)$; once a_0, \dots, a_m have been defined, let

$$a_{m+1} = f(m+1) - P_m(m+1).$$

Noting that $\binom{x}{m+1}$ equals 1 at $x = m+1$ and 0 for $0 \leq x \leq m$, this gives $P_{m+1}(x) = f(x)$ for $x = 0, 1, \dots, m+1$. Now $P_n(x)$ is a degree n polynomial that agrees with $f(x)$ at $x = 0, 1, \dots, n$, so they must be the same polynomial.

Now if f takes on integer values for any $n+1$ consecutive values $m, \dots, m+n$, then by the argument above on $f(x-m)$, $f(x)$ takes on integer values for all x ; in particular, for $x = 0, 1, \dots, n$. Use the above argument to get the desired representation in (c). □

The key idea here in both examples is that once we know that $P(x) = R(x)$ at some points x_1, \dots, x_n , then we can write

$$P(x) = R(x) + (x - x_1) \cdots (x - x_n)Q(x). \tag{D.1}$$

When we're working over \mathbb{Q} or \mathbb{R} , (D.1) doesn't put a restriction on other values of P , but when we're working over \mathbb{Z} or $\mathbb{Z}/m\mathbb{Z}$, then it does. For instance, if we're working over \mathbb{Z} and x_1, \dots, x_n are integers, then we know $P(x)$ and $R(x)$ have to differ by a multiple of $(x - x_1) \cdots (x - x_n)$.

3.1 Problems

1. [1] Suppose P is a polynomial with integer coefficients such that $P(0)$ and $P(1)$ are both odd. Show that P has no integer root.
2. [2] (Schur) Let P be a nonconstant polynomial with integer coefficients. Prove that the set of primes dividing $P(n)$ for some integer n is infinite.
3. [2] Polynomial $P(x)$ has integer coefficients, and satisfies $P(2) = 18$ and $P(3) = 20$. Find all possible integer roots of $P(x) = 0$.
4. [3] (Putnam 2008) Let p be prime. Let $h(x)$ be a polynomial with integer coefficients such that $h(0), h(1), \dots, h(p^2 - 1)$ are distinct modulo p^2 . Show that $h(0), h(1), \dots, h(p^3 - 1)$ are distinct modulo p^3 .
5. [4] (IMO 2006/5) Let $P(x)$ be a polynomial of degree $n > 1$ with integer coefficients and let k be a positive integer. Consider the polynomial

$$Q(x) = \underbrace{P(P(\dots P(P(x))))}_{k \text{ times}}.$$

Prove that there are at most n integers such that $Q(t) = t$.

6. [4] (MOSP 2001) Let f be a polynomial with rational coefficients such that $f(n) \in \mathbb{Z}$ for all $n \in \mathbb{Z}$. Prove that for any integers m, n , the number

$$\text{lcm}[1, 2, \dots, \deg(f)] \cdot \frac{f(m) - f(n)}{m - n}$$

is an integer.

7. [2] (Helpful for the next few problems) Let $f(x) \in R[x]$, and let p_0, p_1, \dots be a sequence of polynomials whose leading coefficients u_0, u_1, \dots are units (i.e. invertible), and $\deg(p_i) = i$. Show that f can be uniquely written in the form

$$f(x) = a_n p_n(x) + \dots + a_1 p_1(x) + a_0 p_0(x).$$

In particular, this is true for $p_i(x) = x^i = x(x - 1) \cdots (x - i + 1)$.

8. [2.5] How many polynomials of degree at most 5 with integer coefficients satisfy $0 \leq P(x) < 120$ for $x = 0, 1, 2, 3, 4, 5$?
9. [4] (USAMO 1995/4) Suppose q_0, q_1, q_2, \dots is an infinite sequence of integers satisfying the following two conditions:

- (a) $m - n$ divides $q_m - q_n$ for $m > n \geq 0$,
- (b) there is a polynomial P such that $|q_n| < P(n)$ for all n .

Prove that there is a polynomial Q such that $q_n = Q(n)$ for each n .

10. [5] (TST 2008/9) Let n be a positive integer. Given an integer coefficient polynomial $f(x)$ define its *signature modulo n* to be the ordered sequence $f(1), \dots, f(n)$ modulo n . Of the n^n such n -term sequences of integers modulo n , how many are the signature of some polynomial $f(x)$ if n is a positive integer not divisible by the cube of a prime? (Easier variant: if n is not divisible by the square of a prime)
11. [5] (variant of TST 2005/3) For a positive integer n , let S denote the set of polynomials $P(x)$ of degree n with positive integer coefficients not exceeding $n!$. A polynomial $P(x)$ in set S is called *fine* if for any positive integer k , the sequence $P(1), P(2), P(3), \dots$ contains infinitely many integers relatively prime to k . Prove that the proportion of fine polynomials is at most

$$\prod_{\text{prime } p \leq n} \left(1 - \frac{1}{p^p}\right).$$

(Original statement: Prove that between 71% and 75% of the polynomials in the set S are fine.)

12. [5] Suppose $f(x)$ is a polynomial of degree d taking integer values such that

$$m - n \mid f(m) - f(n)$$

for all pairs of integers (m, n) satisfying $0 \leq m, n \leq d$. Is it necessarily true that

$$m - n \mid f(m) - f(n)$$

for all pairs of integers (m, n) ?

4 Polynomials in Number Theory

We give an interesting application of polynomials to number theory. Recall the following.

Theorem 4.4.1 (Vieta's Theorem). *Let r_1, \dots, r_n be the roots of $\sum_{i=0}^n a_i x^i$, and let*

$$s_j = \sum_{1 \leq i_1 < \dots < i_j \leq n} r_{i_1} \cdots r_{i_j}.$$

Then $s_j = (-1)^j \frac{a_{n-j}}{a_n}$.

Theorem 4.4.2 (Wolstenholme). *Prove that $\binom{pa}{pb} \equiv \binom{a}{b} \pmod{p^3}$ for prime $p \geq 5$.*

Proof. By Fermat's Little Theorem, $x^{p-1} \equiv 1 \pmod{p}$. Thus in $\mathbb{Z}/p\mathbb{Z}$,

$$x^{p-1} - 1 \equiv \prod_{i=1}^{p-1} (x - i) \pmod{p}. \tag{D.2}$$

Write $(x - 1)^{p-1} = \sum_{i=0}^{p-1} a_i x^i$. Then matching coefficients on both sides of (D.2) gives

$$a_i \equiv 0 \pmod{p} \text{ for all } 1 \leq i < p - 1. \tag{D.3}$$

Since $p \geq 5$, letting $x = p$ gives

$$(p - 1)! = (x - 1)^{p-1} = p^{p-1} + \left(\sum_{i=2}^{p-2} a_i p^i \right) + a_1 p + (p - 1)!$$

since $(-1)(-2) \cdots (-p + 1) = (-1)^{p-1}(p - 1)! = (p - 1)!$. Subtracting $(p - 1)!$ on both sides,

$$0 = p^{p-1} + \left(\sum_{j=2}^{p-2} a_j p^j \right) + a_1 p.$$

Using (D.3), $p^3 \mid a_i p^i$ for $2 \leq i < p - 1$. Hence, since $p \geq 5$, $p^3 \mid p^{p-1} + \sum_{i=2}^{p-2} a_i p^i$. Since p^3 divides the LHS, $p^3 \mid a_1 p$ and $p^2 \mid a_1$. Now $p^3 \mid (kp)^{p-1} + \left(\sum_{i=2}^{p-2} a_i (kp)^i \right)$ as well and we get

$$\begin{aligned} (kp - 1)^{p-1} &= (x - 1)^{p-1} \Big|_{x=pk} \\ &= (kp)^{p-1} + \left(\sum_{j=2}^{p-1} a_j (kp)^j \right) + a_1 kp + (p - 1)! \\ &\equiv (p - 1)! \pmod{p^3}. \end{aligned} \tag{D.4}$$

Now,

$$\begin{aligned} \binom{pa}{pb} &= \frac{(pa)^{pb}}{(pb)!} \\ &= \frac{\prod_{i=a-b+1}^a [(pi)(pi - 1)^{p-1}]}{\prod_{i=1}^b [(pi)(pi - 1)^{p-1}]} \\ &= \frac{a^b}{b!} \left[\prod_{i=1}^b \frac{[p(i + a - b) - 1]^{p-1}}{(pi - 1)^{p-1}} \right] \end{aligned} \tag{D.5}$$

By (D.4), $[p(i + a - b) - 1]^{p-1} \equiv (pi - 1)^{p-1} \pmod{p^3}$. Hence (D.5) becomes $\binom{a}{b}$ modulo p^3 , as needed. \square

Theorem 4.4.3 (Lucas's Theorem). *Suppose that the base p expansions of m and n are*

$$\begin{aligned} m &= (m_k \dots m_1 m_0)_p, \\ n &= (n_k \dots n_1 n_0)_p. \end{aligned}$$

Then

$$\binom{m}{n} \equiv \binom{m_k}{n_k} \cdots \binom{m_1}{n_1} \binom{m_0}{n_0}.$$

Proof. We have the identity

$$(1 + X)^m = (1 + X)^{m_k p^k} \cdots (1 + X)^{m_1 p} (1 + X)^{m_0}.$$

Now take both sides modulo p and use the fact that $(1 + X)^{p^n} \equiv 1 + X^{p^n} \pmod{p}$ to obtain

$$(1 + X)^m \equiv (1 + X^{p^k})^{m_k} \cdots (1 + X^p)^{m_1} (1 + X)^{m_0} \pmod{p}.$$

Now match the coefficients of X^n on each side. The coefficient on the left hand side is $\binom{m}{n}$. For the right hand side, note the only way to get the term X^n is by choosing $X^{n_j p^j}$ from the term $(1 + X^{p^j})^{m_j}$, simply by uniqueness of base p representation; the coefficient of $X^{n_j p^j}$ is $\binom{m_j}{n_j}$. Hence the coefficient of X^n on the right hand side is $\binom{m_k}{n_k} \cdots \binom{m_1}{n_1} \binom{m_0}{n_0}$. Equating the coefficients gives the desired result. \square

Corollary 4.4.4. *Let n be a positive integer, and let $B(n)$ be the number of 1's in the binary expansion of n . Then the number of odd entries in the n th row of Pascal's triangle is $2^{B(n)}$.*

4.1 Problems

- [3] Prove that for prime $p \geq 5$,

$$p^2 | (p-1)! \left(1 + \frac{1}{2} + \cdots + \frac{1}{p-1} \right).$$

- [3.5] (APMO 2006/3) Prove that for prime $p \geq 5$, $\binom{p^2}{p} \equiv p \pmod{p^5}$.
- [3.5] (ISL 2005/N3) Let a, b, c, d, e, f be positive integers. Suppose that the sum $S = a + b + c + d + e + f$ divides both $abc + def$ and $ab + bc + ca - de - ef - fd$. Prove that S is composite.
- [5] (China TST 2009/3) Prove that for any odd prime p , the number of positive integers n satisfying $p \mid n! + 1$ is less than or equal to $cp^{\frac{2}{3}}$, where c is a constant independent of p .²
- [4-5] (TST 2002/2) Let p be a prime number greater than 5. For any positive integer x , define

$$f_p(x) = \sum_{k=1}^{p-1} \frac{1}{(px + k)^2}.$$

Prove that for all positive integers x and y the numerator of $f_p(x) - f_p(y)$, when written in lowest terms, is divisible by p^3 .

²Hint: A polynomial of degree n over a field (such as $\mathbb{Z}/p\mathbb{Z}$) can have at most n zeros.

(MOSP 2007/2.2) Let d be a positive integer. Integers t_1, t_2, \dots, t_d and real numbers a_1, \dots, a_d are given such that

$$a_1 t_1^j + a_2 t_2^j + \dots + a_d t_d^j$$

is an integer for all integers j with $0 \leq j < d$. Prove that

$$a_1 t_1^d + a_2 t_2^d + \dots + a_d t_d^d$$

is also an integer.

5 Resultant

Definition 4.5.1: Let R be a UFD, and let $A(x) = a_m x^m + \dots + a_0$ and $B(x) = b_n x^n + \dots + b_0$ be in $R[x]$. Define

$$M(A, B) = \begin{bmatrix} a_0 & \cdots & \cdots & \cdots & a_m & 0 & 0 \\ 0 & \ddots & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & a_0 & \cdots & \cdots & \cdots & a_m \\ b_0 & \cdots & \cdots & b_n & 0 & 0 & 0 \\ 0 & \ddots & \ddots & \ddots & \ddots & 0 & 0 \\ 0 & 0 & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & b_0 & \cdots & \cdots & b_n \end{bmatrix}$$

where the first n rows contain the a_j and the last m rows contain the b_j . The **resultant** of A and B is

$$\text{Res}(A, B) = \det(M(A, B)).$$

Note that $\text{Res}(A, B)$ is homogeneous of degree n in a_0, \dots, a_m and homogeneous of degree m in b_0, \dots, b_n . The main use of the resultant is that considering it as a function of $a_0, \dots, a_m, b_0, \dots, b_n$, it tells us when A and B have a common factor.

For homogeneous polynomials, we write $A(x, y) = a_m X^m + \dots + a_0 Y^m$ and $B(x, y) = b_n X^n + \dots + b_0 Y^n$ and define the resultant the same way.

Proposition 4.5.2:

1. $\text{Res}(A, B) = 0$ if and only if A and B have a common factor, i.e. have a common zero in \overline{K} .
2. If $a_m b_n \neq 0$, and $A = a_m \prod_{j=1}^m (X - \alpha_j)$, $B = b_n \prod_{k=1}^n (X - \beta_k)$, then

$$\text{Res}(A, B) = a_m^n b_n^m \prod_{j=1}^m \prod_{k=1}^n (\alpha_j - \beta_k).$$

3. There exist polynomials $F, G \in R[a_0, \dots, b_n][X]$ such that

$$FA + GB = \text{Res}(A, B).$$

Proof.

1. A and B have a common factor in $K[X]$ if and only if there exist polynomials nonzero polynomials $C(x) = c_{m-1}X^{m-1} + \cdots + c_0$ and $D(x) = d_{n-1}X^{n-1} + \cdots + d_0$ in $K[X]$ such that $AC = BD$ and $\deg C \leq m - 1$ and $\deg(D) \leq n - 1$. Multiplying out $AC = BD$ and treating it as a system of linear equations in the c_j and d_j , we get that the determinant of the coefficient matrix is $\pm \text{Res}(A, B)$. Thus there is a nonzero solution if and only if $\text{Res}(A, B) = 0$.
2. First assume $a_m = b_n = 1$. We consider the coefficients a_k, b_k and $\text{Res}(A, B)$ as functions of the roots, $\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_n$. By Vieta's formulas, a_{m-k} is homogeneous of degree k in $\alpha_1, \dots, \alpha_m$ and b_{n-k} is homogeneous of degree k in β_1, \dots, β_n . The "big formula" for the determinant says that $\text{Res}(A, B)$ is the sum of entries of the form $\prod_{k=1}^{m+n} M_{k, \sigma(k)}$, where $M = M(A, B)$.
 - For $1 \leq k \leq n$, $M_{k, \sigma(k)}$ is either a polynomial of degree $\sigma(k) - k$ in the α_j or zero, and
 - For $1 \leq k \leq m$, $M_{n+k, \sigma(n+k)}$ is a polynomial of degree $\sigma(n+k) - k$ in the β_j .

Hence if $\prod_{k=1}^{m+n} M_{k, \sigma(k)} \neq 0$, then it has degree

$$\sum_{k=1}^n (\sigma(k) - k) + \sum_{k=1}^m (\sigma(m+k) - k) = \sum_{k=1}^{m+n} \sigma(m+n) - \sum_{k=1}^n k - \sum_{k=1}^m k = mn.$$

Now when $\alpha_j = \beta_k$, then by part 1, $\text{Res}(A, B) = 0$. Hence $\alpha_j - \beta_k$ divides $\text{Res}(A, B)$. By comparing degrees, we must have

$$\text{Res}(A, B) = C \prod_{j=1}^m \prod_{k=1}^n (\alpha_j - \beta_k).$$

To compute C , note that there is exactly one term in the determinant that gives $a_m^n = (\alpha_1 \cdots \alpha_m)^n$, so $C = 1$. By scaling, the desired result holds for $a_m, b_n \neq 0$.

3. We have

$$M(A, B) \begin{pmatrix} X^{m+n-1} \\ \vdots \\ 1 \end{pmatrix} = \begin{pmatrix} X^{n-1}A \\ \vdots \\ A \\ X^{m-1}B \\ \vdots \\ B \end{pmatrix}.$$

Let C denote the cofactor matrix of $M(A, B)$. Multiplying by C^T on both sides gives

$$\text{Res}(A, B) \begin{pmatrix} X^{m+n-1} \\ \vdots \\ 1 \end{pmatrix} = C^T \begin{pmatrix} X^{n-1}A \\ \vdots \\ A \\ X^{m-1}B \\ \vdots \\ B \end{pmatrix}.$$

Let the coefficients of F, G be given by the bottom row of C^T . Then multiplying out the matrices and looking at the bottommost entry gives the desired conclusion. \square

Appendix E

Galois Theory

1 Galois groups and Galois extensions

Definition 5.1.1: The **Galois group** of an extension L/K , denoted

$$\text{Gal}(L/K) = G(L/K)$$

is the group of field automorphisms of L fixing K .

Definition 5.1.2: A **Galois extension** of K is a normal, separable extension.

Theorem 5.1.3. *Suppose L/K is a finite field extension. L/K is a Galois extension if and only if*

$$|G(L/K)| = [L : K].$$

2 Fixed fields

Definition 5.2.1: Let H be a group of automorphisms of a field K . The **fixed field** of H , K^H , is the set of elements of K fixed by every group element.

$$K^H = \{\alpha \in K : \sigma(\alpha) = \alpha \text{ for every } \sigma \in H\}.$$

The following relationship between H and K^H will be instrumental in proving the Fundamental Theorem of Galois Theory.

Theorem 5.2.2 (Fixed field theorem). 1. $[K : K^H] = |H|$: *The degree of K over K^H is the order of the group.*

2. $H = G(K/K^H)$: *K is a Galois extension of K^H with Galois group H .*

Proof.

□

3 Splitting fields

Definition 5.3.1: A **splitting field** of $f \in K[X]$ over K is an extension L/K such that

1. f splits completely in K : $f = (X - \alpha_1) \cdots (X - \alpha_n)$, $\alpha_k \in K$.
2. $L = K(\alpha_1, \dots, \alpha_n)$.

A splitting field is a finite extension, and every finite extension is contained in a splitting field.

The following shows that the splitting property of a splitting field is in a sense independent of the polynomial chosen. This will help us relate splitting fields to Galois extensions.

Definition 5.3.2: A field extension L/K is **normal** if every polynomial $g(X) \in K[X]$ with one root in K splits completely in K .

Theorem 5.3.3 (Splitting theorem). *The normal extensions L/K are exactly the splitting fields of polynomials in $K[X]$.*

Proof. Suppose $g(X)$ has the root $\beta \in K$. Then $p_1(\alpha_1, \dots, \alpha_n) = \beta$ for some $p_1 \in K[X_1, \dots, X_n]$. Let p_1, \dots, p_k be the orbit of p_1 under the symmetric group. Then $\prod_{i=1}^k (X - p_i(\alpha_1, \dots, \alpha_n)) \in K[X]$ by symmetry so it is divisible by $g(X)$, the irreducible polynomial of β . \square

The order of $G = G(L/K)$ divides $[L : K]$, since

$$[L : K] = \underbrace{[L : L^G]}_{|G|} [L^G : K].$$

Theorem 5.3.4 (Characteristic properties of Galois extensions). *For a finite extension L/K , the following are equivalent.*

1. L/K is a Galois extension.
2. $L^{G(L/K)} = L$.
3. L is a splitting field over K .

Proof. (1) \iff (2): By the Fixed Field Theorem, $|G| = [L : L^G]$.

(1) \iff (3): Let γ_1 be a primitive element for L with irreducible polynomial f . Let $\gamma_1, \dots, \gamma_r$ be the roots of f in L . There is a unique K -automorphism σ_i sending $\gamma_1 \mapsto \gamma_i$ for each i and these make up the group $G(L/K)$. Thus the order of $G(L/K)$ is equal to the number of conjugates of γ_1 in L . Hence we get the following chain of equivalences.

1. L/K Galois
2. $|G| = [L : L^G]$

3. f splits completely in K
4. K is a splitting field.

□

Proposition 5.3.5 (Properties of the Galois group): If L/K is a Galois extension, and $g \in K[X]$ splits completely in L with roots β_1, \dots, β_r , then

1. G operates on the set of roots β_i .
2. G operates faithfully if L is a splitting field of g over K .
3. G operates transitively if g is irreducible over K .
4. If L is the splitting field of irreducible g , then G embeds as a transitive subgroup of S_r .

4 Fundamental theorem of Galois theory

Theorem 5.4.1 (Fundamental theorem of Galois theory). *Let L/K be a finite Galois extension and let $G = G(L/K)$. Then there is a bijection between subgroups of G and intermediate fields, defined by*

$$\begin{aligned} H &\mapsto K^H \\ G(L/K') &\leftrightarrow K'. \end{aligned}$$

Moreover, letting $K' = K^H$, K'/K is a Galois extension iff H is a normal subgroup of G . If so, then $G(K'/K) \cong G/H$. [Diagram here.]

Proof. Let γ_1 be a primitive element for K'/K and g the irreducible polynomial for γ_1 over K . Let the roots of g in K be $\gamma_1, \dots, \gamma_r$. For $\sigma \in G$, $\sigma(\gamma_1) = \gamma_i$, the stabilizer of γ_i is $\sigma H \sigma^{-1}$. Thus $\sigma H \sigma^{-1} = H$ if and only if $\gamma_i \in K' = K^H$. H is normal iff all $\gamma_i \in L$ iff K'/K Galois. Restricting σ to L gives a homomorphism $\varphi : G \rightarrow G(K'/K)$ with kernel H . □

Definition 5.4.2: A **normal basis** of a Galois extension L/K is a basis in the form

$$\{\sigma(\beta) : \sigma \in G(L/K)\}$$

for some $\beta \in L$.

Theorem 5.4.3 (Normal basis theorem). *Every Galois extension has a normal basis.*

Proof. Write $G(L/K) = \{\sigma_1, \dots, \sigma_m\}$. Consider two cases.

Case 1: K is infinite. We show the following.

Lemma 5.4.4. *If $f \in K[X_1, \dots, X_m]$ is such that $f(\sigma_1\alpha, \dots, \sigma_m\alpha) = 0$ for all $\alpha \in E$, then $f = 0$.*

Proof. Let $X = (X_1, \dots, X_m)^T$; we write $f(X)$ for $f(X_1, \dots, X_m)$. Let $Y = (Y_1, \dots, Y_m)$, and define

$$g(Y) = g \begin{pmatrix} \sigma_1\alpha_1 & \cdots & \sigma_1\alpha_m \\ \vdots & \ddots & \vdots \\ \sigma_m\alpha_1 & \cdots & \sigma_m\alpha_m \end{pmatrix} X.$$

Then by assumption $g(\alpha) = 0$ for each $\alpha \in K$. Since K is infinite g is the zero polynomial. Note the matrix $\begin{pmatrix} \sigma_1\alpha_1 & \cdots & \sigma_1\alpha_m \\ \vdots & \ddots & \vdots \\ \sigma_m\alpha_1 & \cdots & \sigma_m\alpha_m \end{pmatrix}$ is invertible (this is corollary of indep. of char -). Hence f must also be the zero polynomial. \square

Let A be the matrix with X_k in entry (i, j) if $\sigma_i \circ \sigma_j = \sigma_k$. Let

$$f(X_1, \dots, X_m) = \det(A).$$

Note $f(1, 0, \dots, 0)$ is the determinant of a permutation matrix (since given any g, h in a group, there is exactly one element k and one element l so that $lg = hk = h$), so equals ± 1 . This shows f is not the zero polynomial. Therefore, by Lemma 5.4.4, there exists $\alpha \in K$ such that $f(\sigma_1\alpha, \dots, \sigma_m\alpha) \neq 0$. Suppose that $a_1, \dots, a_m \in K$ and

$$\sum_{k=1}^m a_k \sigma_k(\alpha) = 0.$$

Then

$$\sum_{k=1}^m a_k \sigma_i \sigma_k(\alpha) = 0$$

for all i . Think of this as a system in the a_i . The matrix corresponding to this system is $\det(A) \neq 0$, so all the $a_i = 0$. This shows that $\sigma_k(\alpha)$ are linearly independent.

Case 2: K is a finite field. Then the Galois group is cyclic (Theorem ??); say $G = \langle \sigma \rangle$. By independence of characters, $I, \sigma, \dots, \sigma^{n-1}$ are linearly independent so the minimal polynomial of σ is $X^n - 1$. Consider L as a $K[\sigma] \cong K[X]/(X^n - 1)$ -module. By the structure theorem for modules, we have

$$L \cong K[X]/(p_1) \oplus \cdots \oplus K[X]/(p_m)$$

for some polynomials (p_1) dividing $X^n - 1$ with $p_1 \mid \cdots \mid p_m$. Since the minimal polynomial of σ is $X^n - 1$, we must have $p_m = X^n - 1$. But $[L : K] = n$ so $m = 1$ and $L \cong K[X]$.¹ This means there exists an element α such that $\alpha, \sigma\alpha, \dots, \sigma^{n-1}\alpha$ generate L over K . \square

¹Compare this to the proof of primitive elements in finite fields.

5 Cubic and quartic equations

6 Quintic equations

Theorem 5.6.1 (Quintic impossibility theorem).

7 Inverse limits and profinite groups

To study infinite Galois groups, it is fruitful to view them as the “limit” of finite Galois groups. Thus we first introduce the notion of an inverse limit. This gives infinite Galois groups the structure of a *profinite group*, and its topology becomes important.

7.1 Limits

We will eventually care about limits not just for abelian groups but also topological groups, modules, and so forth. To take care of all this in one fell swoop, we introduce a bit of abstraction, via category theory.

Definition 5.7.1: A **category** \mathcal{C} is a collection of **objects** and **morphisms** (or maps). Each morphism φ has a source and target object A and B ; let $\text{Hom}_{\mathcal{C}}(A, B)$ be the set of morphisms from A to B . There is a composition law

$$\begin{aligned}\text{Hom}_{\mathcal{C}}(A, B) \times \text{Hom}_{\mathcal{C}}(B, C) &\rightarrow \text{Hom}_{\mathcal{C}}(A, C) \\ (\alpha, \beta) &\mapsto \beta \circ \alpha\end{aligned}$$

satisfying the following:

1. For each object B there exists an **identity morphism** $1_B \in \text{Hom}_{\mathcal{C}}(B, B)$ such that $1_B \circ \alpha = \alpha$ for any $\alpha \in \text{Hom}_{\mathcal{C}}(A, B)$ and $\beta \circ 1_B = \beta$ for any $\beta \in \text{Hom}_{\mathcal{C}}(B, C)$.
2. Composition is associative:

$$\gamma \circ (\beta \circ \alpha) = (\gamma \circ \beta) \circ \alpha$$

for any $\alpha \in \text{Hom}_{\mathcal{C}}(A, B)$, $\beta \in \text{Hom}_{\mathcal{C}}(B, C)$, and $\gamma \in \text{Hom}_{\mathcal{C}}(C, D)$.

A morphism $\alpha \in \text{Hom}_{\mathcal{C}}(A, B)$ is an **isomorphism** if there exists $\beta \in \text{Hom}_{\mathcal{C}}(B, A)$ such that $\beta \circ \alpha = 1_A$ and $\alpha \circ \beta = 1_B$.

Example 5.7.2:² We can often think of the objects as sets, possibly endowed with extra structure, and morphisms as maps between them preserving the structure.


²We have to be careful about the word “sets”... See [Mac71] for all the stuff we’re sweeping under the rug.

1. ((Sets)) Objects: sets. Morphisms: functions.
2. ((Rings)) Objects: rings. Morphisms: ring homomorphisms.
3. ((R -mod)), where R is a ring. Objects: R -modules. Morphisms: ring homomorphisms.
4. ((Groups)) Objects: groups. Morphisms: group homomorphisms.
 - (a) ((Ab Groups)) Objects: abelian groups. Morphisms: group homomorphisms.
5. ((Top)) Objects: topological spaces. Morphisms: Continuous maps.
6. ((Top Groups)) Objects: topological groups.³ Morphisms: Continuous homomorphisms.

However, objects in categories do not have to be sets. For instance, any poset S can be turned into a category, by letting the elements be the objects, and declaring a morphism φ_j^i whenever $i, j \in S$ and $i \preceq j$.

Definition 5.7.3: Let \mathcal{C} be a category. Let $\{A_i\}$ and $\{\varphi_j^i\}$ be a set of objects in \mathcal{C} and homomorphisms between them.⁴ We say that $(\{A_i\}, \{\varphi_j^i\})$ form a **inverse** (or projective) **system** if the following two conditions are satisfied.

1. For every $A_i \neq A_j$ there exists A_k such that there are morphisms $\varphi_i^k : A_k \rightarrow A_i$ and $\varphi_j^k : A_k \rightarrow A_j$.



2. For every pair of maps $\varphi_k^j : A_j \rightarrow A_k$ and $\varphi_k'^j : A_j \rightarrow A_k$ there exists a map⁵ $\alpha_j^i : A_i \rightarrow A_j$ such that $\varphi_k^j \circ \varphi_j^i = \varphi_k'^j \circ \varphi_j^i$.

In our applications there will only ever be one map $A_i \rightarrow A_j$, so the second condition is empty.

Finally, we define the notion of inverse limit.

³Groups endowed with a topology such that multiplication is continuous on $G \times G$ and taking the inverse is continuous.

⁴We're allowed to have different maps between A_i and A_j ; however in our examples we usually won't.

⁵called the *equalizer*

Definition 5.7.4: Let $\{A_i\}$ and $\{\varphi_j^i\}$ be a set of objects in \mathcal{C} and homomorphisms between them. Suppose that $\{\varphi_j^i\}$ is closed under composition.⁶ We say a sequence of maps $\alpha_i : A_i \rightarrow A$ is **compatible** if for every map $\varphi_j^i : A_i \rightarrow A_j$ in our set of maps,

$$\alpha_j = \varphi_j^i \circ \alpha_i.$$

The **inverse limit**

$$A = \varprojlim A_i$$

is the unique object in \mathcal{C} (up to isomorphism) with compatible maps α_i , satisfying the following universal mapping property (UMP): For every object B with compatible maps β_i , there is a map $\varphi : B \rightarrow A$ such that $\beta_i = \alpha_i \circ \varphi$ for every i , i.e. the following commutes:

$$\begin{array}{ccc}
 & B & \\
 \beta_i \swarrow & \downarrow \varphi & \searrow \beta_j \\
 A_i & A & A_j \\
 \alpha_i \swarrow & \varphi_j^i & \searrow \alpha_j \\
 & A_i & \rightarrow A_j
 \end{array} \tag{E.1}$$

This is a very abstract definition, but we will be able to construct A explicitly in the cases we care about. Uniqueness follows from the UMP; the inverse limit exists for all inverse systems if and only if \mathcal{C} has *products* and *equalizers*. (See 18.705 notes.)

Theorem 5.7.5. *Suppose \mathcal{C} is $((\text{Sets}))$, $((\text{Groups}))$, $((R\text{-mod}))$, or $((R\text{-alg}))$. If $(\{A_i\}, \{\varphi_j^i\})$ is an inverse system, then $\varprojlim A_i$ can be realized as the set of all sequences*

$$\{(a_i) : a_i \in A_i, \varphi_j^i(a_i) = a_j \text{ for all } \varphi_j^i\},$$

with the natural module or algebra structure, as applicable.

Proof. Just verify that the UMP is satisfied. □

Example 5.7.6: The ring of p -adic integers

$$\mathbb{Z}_p = \varprojlim \mathbb{Z}/p^n\mathbb{Z}$$

is defined the inverse limit of $(\{\mathbb{Z}/p^n\mathbb{Z}\}_{n \in \mathbb{Z}}, \varphi_m^n)$ where $\varphi_m^n : \mathbb{Z}/p^n\mathbb{Z} \rightarrow \mathbb{Z}/p^m\mathbb{Z}$, for $n \geq m$, are the natural projection maps. An element of \mathbb{Z}_p can be thought of as a number modulo arbitrarily high powers of p . We have an injective map $\mathbb{Z} \hookrightarrow \mathbb{Z}_p$, but there are elements of \mathbb{Z}_p not in \mathbb{Z} (think this through).

We will explore p -adics in depth in Chapter 7.

⁶Equivalently, the A_i and φ_j^i are indexed by a category, i.e. there is a functor from a small category into \mathcal{C} .

Example 5.7.7: Define

$$\widehat{\mathbb{Z}} = \varprojlim \mathbb{Z}/n\mathbb{Z}$$

as the inverse limit of $(\{\mathbb{Z}/n\mathbb{Z}\}_{n \in \mathbb{Z}}, \varphi_m^n)$, where the maps $\varphi_m^n : \mathbb{Z}/n\mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$ with $m \mid n$ are given by projection.

In the next section we will interpret these limits not just as limit of groups, but of topological groups.

7.2 Profinite groups

We assume knowledge of topology (continuous maps, compactness, separation axioms, connectedness, product topology, Tychonoff's theorem).

Definition 5.7.8: A **profinite group** is an inverse limit $\varprojlim_{i \in I} G_i$ of finite discrete topological groups G_i .

Suppose that $\varphi_i : G \rightarrow G_i$ are all surjective. The **order** $\#G$ of G is the formal product

$$\prod_p p^{\max_{i \in I} v_p(|G_i|)}.$$

In other words it is the “least common multiple” of the $|G_i|$.

We know that if we only consider the G_i as *groups*, then by Theorem 5.7.5, the inverse limit can be described as the set of tuples $(g_i)_{i \in I}$ such that $\varphi_j^i(g_i) = g_j$ for every transition map $\varphi_j^i : G_i \rightarrow G_j$. But we need to show that the inverse limit is well-defined when the G_i are *topological groups*. We give a topology on the inverse limit of groups, $\varprojlim_{i \in I} G_i$, so that it satisfies the UMP for the inverse limit of topological groups. (In the category of topological groups, homomorphisms must be continuous.)

Proposition 5.7.9: Give

$$G = \varprojlim_{\text{groups}} G_i$$

the following topology: Equip each finite group G_i with the discrete topology and $\prod_{i \in I} G_i$ with the product topology. Then $G = \varprojlim_{i \in I} G_i$ is the closed subspace of $\prod_{i \in I} G_i$ of compatible sequences; give it the subspace topology.

Then

$$G = \varprojlim_{\text{top. group}} G_i.$$

Proof. We show that G satisfies the UMP.

First, note that the maps $\varphi_i : G \rightarrow G_i$ are continuous. Indeed for any open $U_i \in G_i$, letting π_i be the projection map $\prod_{i \in I} G_i$, $\pi_i^{-1}(U_i)$ is open. Hence $\varphi_i^{-1}(U_i) = \pi_i^{-1}(U_i) \cap G$ is open in G .

Now let H be a topological group with compatible maps $\beta_i : H \rightarrow G_i$. In order for E.1 to commute, we must define

$$\varphi(h) = (\beta_i(h))_i.$$

This is a continuous map $\beta_i : H \rightarrow \prod_i G_i$ because it is the product of continuous maps; it is also a homomorphism. Its image is in $G \subseteq \prod_i G_i$ because the β_i are compatible. Since G is given the subspace topology, β is continuous, as desired. \square

The following characterizes the topology of profinite groups.

Proposition 5.7.10: A topological group G is profinite iff it is compact, Hausdorff, and totally disconnected.

Proof. First suppose $G = \varprojlim_{i \in I} G_i$ is profinite.

1. $\prod_{i \in I} G_i$ is compact by Tychonoff's Theorem (an arbitrary product of compact spaces is compact) so the closed subspace G is compact.
2. Given $g = (g_i)$ and $h = (h_i)$, suppose $g_i \neq h_i$. Partition G_i into two sets A and B containing g_i and h_i , respectively. Then $\alpha_i^{-1}(g_i)$ and $\alpha_j^{-1}(g_j)$ are disjoint clopen (open and closed) sets containing g and h , respectively. This shows that G is Hausdorff and totally disconnected.

The converse is left as an exercise (we won't need it). \square

Profinite groups can be constructed from arbitrary abelian groups as follows.

Definition 5.7.11: Let G be a group. Define the **profinite completion** of G to be

$$\widehat{G} = \varprojlim_{\substack{N \text{ normal of} \\ \text{finite index}}} G/N$$

with the natural projection maps.

Example 5.7.12: This agrees with our definition of $\widehat{\mathbb{Z}}$ in Example 5.7.7. In the profinite topology of $\widehat{\mathbb{Z}}$, the subsets $n\widehat{\mathbb{Z}}$ form a neighborhood base of 0.

8 Infinite Galois theory

Let Ω/K be an infinite Galois extension. We equip $G(\Omega/K)$ with a topology by interpreting it as a profinite group, as follows.

Proposition 5.8.1:

$$G(\Omega/K) = \varprojlim_{L/K \text{ finite}} G(L/K),$$

where the limit is with respect to the quotient maps $G(M/K) \rightarrow G(L/K)$.

Proof. Identify the right side with compatible elements of $\prod_{L/K} G(L/K)$ and send $\sigma \in G(\Omega/K)$ to $(\sigma|_L)_L$. This is a bijection because any element of Ω is in a finite extension over K , so specifying a map $\Omega \rightarrow \Omega$ is the same as specifying a compatible sequence of maps $L \rightarrow L$ for every finite Galois extension. \square

Now we give $G(\Omega/K)$ the profinite topology. Equivalently, it is the topology such that a neighborhood base of 1 is

$$G(S) = \{\sigma \in G(\Omega/K) : \sigma s = s \text{ for all } s \in S\}, \quad S \text{ finite.}$$

We next show surjectivity of the quotient map.

Proposition 5.8.2: Every homomorphism $\sigma : L \rightarrow \Omega$ extends to a homomorphism $\Omega \rightarrow \Omega$. If L/K is Galois, then the restriction map $G(\Omega/K) \twoheadrightarrow G(L/K)$ is surjective.

Proof. Use Zorn's lemma, as follows. Define a poset P whose elements are pairs (M, φ_M) , where M is a field with $L \subseteq M \subseteq \Omega$ and φ_M is a homomorphism $M \rightarrow \Omega$. Introduce a partial ordering by saying

$$(M, \varphi_M) \preceq (N, \varphi_N)$$

if $M \subseteq N$ and $\varphi_N|_M = \varphi_M$. If (M_i, φ_{M_i}) is a chain (totally ordered subset), then it has a maximal element in P , namely,

$$\left(\bigcup_i M_i, \varphi \right)$$

where φ is defined as $\varphi(x) = \varphi_i(x)$ if $x \in M_i$. Thus by Zorn's lemma P has a maximal element (M, φ_M) .

For any element $\alpha \in \Omega$, by finite Galois theory (ref) we can extend (M, φ_M) to $(M(\alpha), \varphi_{M(\alpha)})$. By maximality of M , $M = M(\alpha)$, i.e. $M = \Omega$.

The second part follows directly. \square

The following is the analogue of the fixed field theorem. Note that topology now plays a role.

Theorem 5.8.3 (Fixed field theorem, infinite extensions). *Suppose Ω/K is Galois and $G = G(\Omega/K)$.*

1. $G(\Omega/L)$ is closed, and $\Omega^{G(\Omega/L)} = L$.
2. For every subgroup $H \subseteq G$, $G(\Omega/\Omega^H) = \overline{H}$.

Proof. 1. The sets $G(S)$ are open of finite index, hence closed. Hence $G(\Omega/L) = \bigcap_{\text{finite } S \subseteq L} G(S)$ is closed.

For the second part, note that for every finite Galois extension M/L , we know

$$\Omega^{G(M/L)} \cap M = L.$$

Since this is true for every such M/L , and $G(\Omega/L) \twoheadrightarrow G(M/L)$ is surjective, the result follows.

2. It is clear that $G(\Omega/\Omega^H) \supseteq H$. By part 1, $G(\Omega/\Omega^H)$ is closed, so it contains \overline{H} .
FINISH...

□

Theorem 5.8.4 (Fundamental theorem of infinite Galois theory). *There is a bijection between closed subgroups of G and intermediate fields L with $K \subseteq L \subseteq \Omega$.*

$$\begin{aligned} H &\mapsto \Omega^H \\ G(\Omega/L) &\leftrightarrow L. \end{aligned}$$

We have the following.

1. This map is inclusion-reversing.
2. H is open if and only if $[\Omega^H : K] < \infty$. Then $[G : H] = [\Omega^H : K]$.
3. $\sigma H \sigma^{-1}$ corresponds to σM , so H is normal if and only if Ω^H/K is Galois. Then $G(\Omega^H/K) \cong G/H$.

Note given closed, open iff of finite index.

Example 5.8.5: We have

$$G(\overline{\mathbb{F}_p}/\mathbb{F}_p) = \varprojlim G(\Omega/\mathbb{F}_p) = \varprojlim \mathbb{Z}/n\mathbb{Z} = \widehat{\mathbb{Z}}.$$

Example 5.8.6: Let

$$\mathbb{Q}(\zeta_\infty) := \mathbb{Q}(\{\zeta_n : n \in \mathbb{N}\}).$$

Then

$$G(\mathbb{Q}(\zeta_\infty)/\mathbb{Q}) = \varprojlim_{n \in \mathbb{N}} G(\mathbb{Q}(\zeta_n)/\mathbb{Q}) = \varprojlim (\mathbb{Z}/n\mathbb{Z})^\times = \widehat{\mathbb{Z}}^\times$$

(Note that, by the Kronecker-Weber Theorem 11.11.7.2, $\mathbb{Q}(\zeta_\infty) = \mathbb{Q}^{\text{ab}}$.)

Appendix F

Arithmetic over Finite Fields

Our main goal in this chapter is to find a way to find the number of solutions for equations over finite fields. One problem we will look at in detail is, for a fixed b , how many solutions are there to

$$b = y_1^d + \cdots + y_n^d$$

over a finite field? We encapsulate the number of representations as a sum of n d th powers in a sum of orthonormal functions on \mathbb{F}_q called the additive characters χ . We consider the product

$$\left(\sum_{y \in \mathbb{F}_q} \chi(y^d) \right)^n = \sum_{y_1, \dots, y_n \in \mathbb{F}_q} \chi(y_1^d + \cdots + y_n^d). \quad (\text{F.1})$$

(The additive characters have the nice property that $\chi(a + b) = \chi(a)\chi(b)$.) Note (F.1) is true for all characters. To extract out the coefficient of $\chi(b)$, we multiply by $\overline{\chi(b)}$, average over all distinct characters χ , and take advantage of orthonormality to get

$$r_{d,n}(b) = \frac{1}{q} \sum_{\chi} \left\{ \left(\sum_{y \in \mathbb{F}_q} \chi(y^d) \right)^n \overline{\chi(b)} \right\}. \quad (\text{F.2})$$

In the next section we will give define and give properties of characters that help us estimate (F.2).

1 Characters

To evaluate (F.2) it would be helpful if $\chi(y^d) = \chi(y)^d$. However, this cannot hold as we defined χ so that it would preserve additive structure, not multiplicative structure. Thus to evaluate (F.2) we would like to rewrite it as a sum of functions ψ such that $\psi(ab) = \psi(a)\psi(b)$, and such that the set of ψ are orthonormal. Thus we will need both the concepts of additive and multiplicative characters. We make this precise below.

Definition 6.1.1: Let G be an abelian group. A **character** of G is a homomorphism from G to \mathbb{C}^\times . A character is trivial if it is identically 1. We denote the trivial character by χ_0 or ψ_0 .

Definition 6.1.2: Let R be a given finite ring. An additive character $\chi : R^+ \rightarrow \mathbb{C}$ is a character χ with R considered as an additive group. A multiplicative character $\psi : R^\times \rightarrow \mathbb{C}$ is a character with R^\times , the units of R , considered as a multiplicative group.

The two cases we will be working with are $R = \mathbb{Z}/N\mathbb{Z}$ (Section 1.1), and $R = \mathbb{F}_q$ (Section 1.2). We extend multiplicative characters ψ to R by defining $\psi(x) = 0$ for $x \in R \setminus R^\times$, except we follow the convention of setting $\psi_0(0) = 1$ when $R = \mathbb{F}_q$. Note that in any case the extended ψ still preserves multiplication.

We proceed to give an explicit description of characters for abelian groups. First, recall the following theorem.

Theorem 6.1.3 (Structure Theorem for Abelian Groups). *Let G be a finite abelian group. Then there exist positive integers m_1, \dots, m_k so that*

$$G \cong \mathbb{Z}/m_1\mathbb{Z} \times \cdots \times \mathbb{Z}/m_k\mathbb{Z}.$$

Theorem 6.1.4 (Characters of abelian groups). *The group $G = \mathbb{Z}/m_1\mathbb{Z} \times \cdots \times \mathbb{Z}/m_k\mathbb{Z}$ has $|G|$ characters and each is given by an element $(r_1, \dots, r_k) \in \mathbb{Z}/m_1\mathbb{Z} \times \cdots \times \mathbb{Z}/m_k\mathbb{Z}$:*

$$\chi_{r_1, \dots, r_k}(n_1, \dots, n_k) = \prod_{j=1}^k e^{\frac{2\pi i r_j n_j}{m_j}}.$$

Moreover the set of characters \widehat{G} form a multiplicative group isomorphic to G .¹

Proof. Use Theorem 6.1.3. It is easy to check that $\chi = \chi_{r_1, \dots, r_k}$ is a homomorphism. Let e_j be the element in G with 1 in the j th coordinate and 0's elsewhere. Since $\chi(e_j)^{m_j} = 1$, we must have $\chi(e_j) = e^{\frac{2\pi i r_j}{m_j}}$ for some r_j . Each element of G can be expressed as a combination of the e_j , so this shows all characters are in the above form.

This shows that $(r_1, \dots, r_k) \mapsto \chi_{r_1, \dots, r_k}$ is surjective and hence an isomorphism. \square

Corollary 6.1.5. *Every finite abelian group G has $|G|$ characters.*

Theorem 6.1.6 (Orthogonality relations). *Let G be a finite abelian group and $\chi_j, 1 \leq j \leq n$ be all characters of G . Then*

$$1. \text{ (Row orthogonality) } \langle \chi_j, \chi_k \rangle := \frac{1}{|G|} \sum_{g \in G} \chi_j(g) \overline{\chi_k(g)} = \begin{cases} 0, & j \neq k \\ 1, & j = k \end{cases}$$

$$2. \text{ (Column orthogonality) } \sum_{j=1}^n \chi_j(g) \overline{\chi_j(h)} = \begin{cases} 0, & g \neq h \\ |G|, & g = h \end{cases}$$

¹This is a noncanonical isomorphism.

Proof. Write G as $\mathbb{Z}/m_1\mathbb{Z} \times \cdots \times \mathbb{Z}/m_k\mathbb{Z}$. Let (r_1, \dots, r_k) and (s_1, \dots, s_k) be in G . Then

$$\langle \chi_{r_1, \dots, r_k}, \chi_{s_1, \dots, s_k} \rangle = \sum_{(p_1, \dots, p_k) \in G} \prod_{j=1}^k e^{\frac{2\pi i(r_j - s_j)p_j}{m_j}} \quad (\text{F.3})$$

$$= \sum_{(p_1, \dots, p_{k-1}) \in G} \left[\left(\prod_{j=1}^{k-1} e^{\frac{2\pi i(r_j - s_j)p_j}{m_j}} \right) \sum_{p_k=0}^{m_k-1} e^{2\pi i(r_k - s_k)p_k} \right]. \quad (\text{F.4})$$

If $(r_1, \dots, r_k) = (s_1, \dots, s_k)$ then (F.3) evaluates to $|G|$. Otherwise, we may assume without loss of generality that $r_k \neq s_k$; then the inner sum in (F.4) evaluates to 0 by writing it as a geometric series.

The proof for column orthogonality is similar. \square

The most useful case of row orthogonality is when we set $\chi_k = \chi_0$:

Corollary 6.1.7. *If χ is a character of G and $\chi \neq \chi_0$ then*

$$\sum_{g \in G} \chi(g) = 0.$$

Having established the basic properties of characters of abelian groups, we now turn to the specific cases $\mathbb{Z}/N\mathbb{Z}$ and \mathbb{F}_q .

1.1 Dirichlet characters

For our applications, it is helpful to think of consider characters on $\mathbb{Z}/N\mathbb{Z}$ as functions on \mathbb{Z} . From Theorem 6.1.4, the additive characters are simply given by

$$\chi_a(g) = e^{\frac{2\pi i a g}{N}}.$$

Next we consider multiplicative characters.

Definition 6.1.8: A **Dirichlet character** of level N is a function $\chi : \mathbb{Z} \rightarrow \mathbb{C}$ that induces a group homomorphism

$$\chi : (\mathbb{Z}/N\mathbb{Z})^\times \rightarrow \mathbb{C},$$

and such that $\chi(n) = 0$ for any n sharing a common factor with N . In other words, it induces a multiplicative character $\mathbb{Z}/N\mathbb{Z} \rightarrow \mathbb{C}$.

We say χ is **principal** if $\chi(n) = 1$ for all $(\mathbb{Z}/N\mathbb{Z})^\times$, and **primitive** if χ does not induce a group homomorphism $(\mathbb{Z}/M\mathbb{Z})^\times \rightarrow \mathbb{C}$ for any $M < N$.

We say χ is **even** or **odd** if $\chi(-1) = 1$ or $\chi(-1) = -1$, respectively; we say χ is **real** when $\text{im}(\chi) \subset \mathbb{R}$ and say it is **nonreal** otherwise.

Any character can be written uniquely as a product of a primitive character χ_1 of level $M \mid N$ and the principal character of level N :

$$\chi = \chi_1 \chi_0.$$

1.2 Characters on finite fields

We give the additive and multiplicative characters on \mathbb{F}_q explicitly. We know that \mathbb{F}_q^\times is cyclic; let ξ be a generator.

Theorem 6.1.9 (Multiplicative characters of \mathbb{F}_q). *The multiplicative characters of \mathbb{F}_q are given by*

$$\psi_j(\xi^n) = e^{\frac{2\pi i j n}{q-1}}$$

for $0 \leq j < q - 1$.

Proof. By identifying $\xi \in \mathbb{F}_q^\times$ with $1 \in \mathbb{Z}/(q-1)\mathbb{Z}$, this follows directly from Theorem 6.1.4. \square

Describing the additive characters takes slightly more creativity, since it is inconvenient to decompose \mathbb{F}_q^+ into cyclic groups.

Theorem 6.1.10 (Additive characters of \mathbb{F}_q). *Suppose $q = p^r$ with p prime. The additive characters of \mathbb{F}_q are given by*

$$\chi_a(g) = e^{\frac{2\pi i}{p} \operatorname{tr}(ag)} \tag{F.5}$$

for $a \in \mathbb{F}_q$ where²

$$\operatorname{tr}(g) = g + g^p + \cdots + g^{p^{r-1}}.$$

Proof. The automorphisms of \mathbb{F}_q fixing \mathbb{F}_p are generated by the Frobenius automorphism σ sending g to g^p . Since $\operatorname{tr}(g)$ is fixed under this operation, it must be in the ground field \mathbb{F}_p . This makes (F.5) well-defined since only the value of $\operatorname{tr}(ag)$ modulo p matters in (F.5). The fact that χ_a is a homomorphism comes directly from the fact that σ is a homomorphism.

Since $\chi_1(ag) = \chi_a(g)$, if $\chi_a = \chi_b$ then $\chi_1(ag) = \chi_1(bg)$ and $\chi_1((a-b)g) = 0$. However, χ_1 is not trivial (identically equal to 1) since there are at most p^{r-1} values of g such that $g + \cdots + g^{p^{r-1}} = 0$. Thus $a = b$. This shows all characters in our list are distinct. Since we have found $|G|$ characters we have found all of them. \square

Remark 6.1.11: In general, a n -dimensional complex representation of a group G is a homomorphism ρ from G into $GL_n(\mathbb{C})$, and the character χ of a representation is defined by $\chi(g) = \operatorname{tr}(\rho(g))$. This coincides with Definition 6.1.1 for abelian G , if we just consider 1-dimensional representations, since ρ is multiplication by a constant and χ is just that constant.

The general case of Corollary 6.1.5 is replaced by the following: every finite group has a number of irreducible characters equal to the number of conjugacy classes. The orthogonality relations hold when we consider just irreducible characters, and with $|G|$ replaced by the size of the centralizer of g in the equation for column orthogonality.

²For the general definition of trace see Definition 1.1.2.1.

2 Gauss Sums

To relate additive characters to multiplicative characters, we need to evaluate sums in the form

$$G(\psi, \chi) = \sum_{y \in \mathbb{F}_q^\times} \psi(y)\chi(y). \quad (\text{F.6})$$

where ψ is a multiplicative character and χ is an additive character.

Suppose we wanted to write an additive character on \mathbb{F}_q in terms of multiplicative characters. By row orthogonality, $\frac{1}{q-1} \sum_{\psi \in \widehat{\mathbb{F}_q^\times}} \psi(y)\overline{\psi(g)}$ equals 1 if $y = g$ and is 0 otherwise. This allows us to introduce multiplicative characters as follows: for $y \in \mathbb{F}_q^\times$,

$$\begin{aligned} \chi(y) &= \frac{1}{q-1} \sum_{g \in \mathbb{F}_q^\times} \chi(g) \sum_{\psi \in \widehat{\mathbb{F}_q^\times}} \psi(y)\overline{\psi(g)} \\ &= \frac{1}{q-1} \sum_{\psi \in \widehat{\mathbb{F}_q^\times}} \psi(y) \sum_{g \in \mathbb{F}_q^\times} \overline{\psi(g)}\chi(g) \\ &= \frac{1}{q-1} \sum_{\psi \in \widehat{\mathbb{F}_q^\times}} G(\overline{\psi}, \chi)\psi(y). \end{aligned} \quad (\text{F.7})$$

The Gauss sums are the coefficients of the expansion of χ in terms of multiplicative characters. The next theorem tells us how to calculate Gauss sums.

Theorem 6.2.1 (Magnitude of Gauss sums). *Let ψ_0 and χ_0 denote the trivial multiplicative and additive characters on \mathbb{F}_q , respectively. Then for multiplicative and additive characters ψ and χ on \mathbb{F}_q , we have*

$$G(\psi, \chi) = \begin{cases} q-1, & \psi = \psi_0, \chi = \chi_0 \\ -1, & \psi = \psi_0, \chi \neq \chi_0 \\ 0, & \psi \neq \psi_0, \chi = \chi_0 \end{cases}$$

and

$$|G(\psi, \chi)| = \sqrt{q}, \quad \psi \neq \psi_0, \chi \neq \chi_0.$$

If ψ is a nontrivial multiplicative character and χ is a primitive additive character on $\mathbb{Z}/N\mathbb{Z}$, then

$$|G(\psi, \chi)| = \sqrt{N}.$$

Proof. The first case is trivial. For the second case,

$$G(\psi_0, \chi) = \sum_{y \in \mathbb{F}_q^\times} \chi(y) = \left(\sum_{y \in \mathbb{F}_q} \chi(y) \right) - 1 = -1$$

by Corollary 6.1.7. The third case directly from Corollary 6.1.7 with ψ .

Now we consider the case when ψ is nontrivial, and either $\chi \neq \chi_0$ (in the case $R = \mathbb{F}_q$) or χ is primitive (in the case $R = \mathbb{Z}/N\mathbb{Z}$), respectively. We have

$$\begin{aligned}
 |G(\psi, \chi)|^2 &= \sum_{g_1, g_2 \in R^\times} \overline{\psi(g_1)} \psi(g_2) \overline{\chi(g_1)} \chi(g_2) \\
 &= \sum_{g_1, g_2 \in R^\times} \psi(g_1^{-1} g_2) \chi(g_2 - g_1) \\
 &= \sum_{h \in R^\times} \sum_{g_1 \in R^\times} \psi(h) \chi(g_1(h-1)) && \text{setting } h = g_1^{-1} g_2 \\
 &= \sum_{h \in R^\times} \psi(h) \left[\left(\sum_{g_1 \in R} \chi(g_1(h-1)) \right) - \sum_{y \in R \setminus R^\times} \chi(y) \right] \\
 &= \sum_{h \in R^\times} \psi(h) \left(\sum_{g_1 \in R} \chi(g_1(h-1)) \right) && \text{by Corollary 6.1.7 with } \psi
 \end{aligned}$$

Now we note the following: when $h = 1$ all terms in the inner sum are 1, so it equals q or N , respectively. When $h \neq 1$, consider two cases.

1. $R = \mathbb{F}_q$: As g_1 ranges over \mathbb{F}_q , $g_1(h-1)$ ranges over \mathbb{F}_q .
2. $R = \mathbb{Z}/N\mathbb{Z}$: As g_1 ranges over $\mathbb{Z}/N\mathbb{Z}$, $g_1(h-1)$ ranges over a subgroup $H \subseteq \mathbb{Z}/N\mathbb{Z}$, hitting each element $\frac{N}{|H|}$ times. Since χ is primitive, $\chi|_H$ is nontrivial.

In either case, Corollary 6.1.7 gives the inner sum to be 0. Hence $|G(\psi, \chi)|^2$ evaluates to $\psi(1)q = q$ or $\psi(1)N = N$, respectively. \square

We will need the following fact later on.

Proposition 6.2.2 (Gauss sum dependence on additive character): Let $R = \mathbb{F}_q$ or $\mathbb{Z}/N\mathbb{Z}$. For $a \in R^\times$ and $b \in R$,

$$G(\psi, \chi_{ab}) = \overline{\psi(a)} G(\psi, \chi_b).$$

Proof. Using the fact that $\chi_c(g) = \chi_1(cg)$,

$$\begin{aligned}
 G(\psi, \chi_{ab}) &= \sum_{y \in R^\times} \psi(y) \chi_{ab}(y) \\
 &= \sum_{y \in R^\times} \psi(y) \chi_b(ay) \\
 &= \sum_{y \in R^\times} \psi(a^{-1}y) \chi_b(y) && \text{replacing } y \rightarrow a^{-1}y \\
 &= \psi(a)^{-1} \sum_{y \in R^\times} \psi(y) \chi_b(y) \\
 &= \overline{\psi(a)} G(\psi, \chi_b) && \square
 \end{aligned}$$

3 Enumerating Solutions

We return to our original problem. Rather than just work with sums of d th powers, we work with diagonal equations

$$a_1 y_1^{d_1} + \dots + a_n y_n^{d_n} = b \tag{F.8}$$

where $a_i \in \mathbb{F}_q^\times$ and $d_i \in \mathbb{N}$. First, note that because of the following lemma, we can restrict to case where $d_i | q - 1$.

Lemma 6.3.1. *The multisets $\{y^d | y \in \mathbb{F}_q\}$ and $\{y^{\gcd(d, q-1)} | y \in \mathbb{F}_q\}$ are equal.*

Proof. Let ξ be a generator for \mathbb{F}_q^\times , and write $d = k \gcd(d, q - 1)$ where $\gcd(k, q - 1) = 1$. Then removing the one occurrence of 0 in the two sets, we get $\{\xi^{jd} | 0 \leq j < q - 1\}$ and $\{\xi^{j \gcd(d, q-1)} | 0 \leq j < q - 1\}$. The lemma follows from the fact that as multisets,

$$\{jd \pmod{q-1} | 0 \leq j < q-1\} = \{j \gcd(d, q-1) \pmod{q-1} | 0 \leq j < q-1\}.$$

Indeed, each multiple of $\gcd(d, q - 1)$ appears $\frac{q-1}{\gcd(d, q-1)}$ times on both sides. □

As (F.8) always has the trivial solution when $b = 0$, we just need to estimate the number of solutions to (F.8) when $b \neq 0$.

Theorem 6.3.2. *[?, 6.37] Fix $b \neq 0, d_i | q - 1$ and let N be the number of solutions to (F.8) when $b \neq 0$ is fixed. Then*

$$|N - q^{n-1}| \leq [(d_1 - 1) \cdots (d_n - 1) - (1 - q^{-\frac{1}{2}})M(d_1, \dots, d_n)]q^{\frac{n-1}{2}}$$

where $M(d_1, \dots, d_n)$ is the number of n -tuples in the set

$$S := \left\{ (j_1, \dots, j_n) \in \mathbb{Z}^n \mid 1 \leq j_i \leq d_i - 1 \text{ and } \sum_{i=1}^n \frac{j_i}{d_i} \in \mathbb{Z} \right\}.$$

Note that we would expect N to be close to q^{n-1} , because there are q^n possible choices for (y_1, \dots, y_n) and q possible values for their sum.

Proof. We use the idea mentioned in the introduction. We have

$$N = \frac{1}{q} \sum_{y_1, \dots, y_n \in \mathbb{F}_q, \chi \in \widehat{\mathbb{F}_q^+}} \chi(a_1 y_1^{d_1} + \dots + a_n y_n^{d_n}) \bar{\chi}(b) = \frac{1}{q} \sum_{y_1, \dots, y_n \in \mathbb{F}_q, \chi \in \widehat{\mathbb{F}_q^+}} \chi(a_1 y_1^{d_1}) \cdots \chi(a_n y_n^{d_n}) \bar{\chi}(b)$$

since by row orthogonality the inner sum is 1 if $a_1 y_1^{d_1} + \dots + a_n y_n^{d_n} = b$ and 0 otherwise. Note that χ_0 contributes q^n to the sum. Taking it out and factoring the remaining terms gives

$$N = q^{n-1} + \frac{1}{q} \sum_{\chi \in \widehat{\mathbb{F}_q^+}, \chi \neq \chi_0} \left(\bar{\chi}(b) \prod_{j=1}^n \sum_{y_j \in \mathbb{F}_q} \chi(a_j y_j^{d_j}) \right) \tag{F.9}$$

We write the sums of additive characters as sums of multiplicative characters using the following lemma.

Lemma 6.3.3. *Let χ be a nontrivial additive character and λ a multiplicative character of order d dividing $q - 1$. Then*

$$\sum_{y \in \mathbb{F}_q} \chi(ay^d) = \sum_{j=1}^{d-1} \bar{\lambda}(a)^j G(\lambda^j, \chi).$$

Proof. Note that λ exists since the group of multiplicative characters is isomorphic to $\mathbb{Z}/(q-1)\mathbb{Z}$ by Theorem 6.1.4. Suppose $\chi = \chi_c$. We write χ as a sum of multiplicative characters using (F.7), get the Gauss sum to be independent of a by using Proposition 6.2.2, and take out the exponent as we were hoping to do:

$$\begin{aligned} \sum_{y \in \mathbb{F}_q} \chi(ay^d) &= \sum_{y \in \mathbb{F}_q} \chi_{ac}(y^d) \\ &= 1 + \sum_{y \in \mathbb{F}_q^\times} \chi_{ac}(y^d) \\ &= 1 + \frac{1}{q-1} \sum_{\psi \in \widehat{\mathbb{F}_q^\times}} \sum_{y \in \mathbb{F}_q^\times} G(\bar{\psi}, \chi_{ac}) \psi(y^d) \\ &= 1 + \frac{1}{q-1} \sum_{\psi \in \widehat{\mathbb{F}_q^\times}} \bar{\psi}(a) G(\bar{\psi}, \chi_c) \sum_{y \in \mathbb{F}_q} \psi(y)^d \end{aligned} \tag{F.10}$$

$$= 1 + \sum_{j=0}^{d-1} \bar{\lambda}(a)^j G(\lambda^j, \chi) \tag{F.11}$$

$$= \sum_{j=1}^{d-1} \bar{\lambda}(a)^j G(\lambda^j, \chi) \tag{F.12}$$

Note (F.11) follows since by Corollary 6.1.7, $\sum_{y \in \mathbb{F}_q^\times} \psi(y)^d = 0$ unless ψ^d is the trivial character, which is true iff ψ is a power of λ . In that case, the inner sum in (F.10) is $q - 1$. In (F.12) we used $G(\psi_0, \chi) = -1$ (Theorem 6.2.1). \square

Using Lemma 6.3.3 and letting λ_j be the multiplicative character with $\lambda_j(\xi^t) = e^{\frac{2\pi it}{d_j}}$ we rewrite (F.9) as

$$\begin{aligned} N - q^{n-1} &= \frac{1}{q} \sum_{\chi \in \widehat{\mathbb{F}_q^+}, \chi \neq \chi_0} \left(\bar{\chi}(b) \prod_{j=1}^n \sum_{k=1}^{d_j-1} \bar{\lambda}_j(a_j)^k G(\lambda_j^k, \chi) \right) \\ &= \frac{1}{q} \sum_{\chi \in \widehat{\mathbb{F}_q^+}, \chi \neq \chi_0} \sum_{(k_1, \dots, k_n), 1 \leq k_i \leq d_i-1} \bar{\chi}(b) \bar{\lambda}_1^{k_1}(a_1) \cdots \bar{\lambda}_n^{k_n}(a_n) G(\lambda_1^{k_1}, \chi) \cdots G(\lambda_n^{k_n}, \chi) \\ &= \frac{1}{q} \sum_{c \in \mathbb{F}_q^\times} \sum_{(k_1, \dots, k_n), 1 \leq k_i \leq d_i-1} \bar{\chi}_c(b) \bar{\lambda}_1^{-k_1}(a_1) \cdots \bar{\lambda}_n^{-k_n}(a_n) G(\lambda_1^{k_1}, \chi_c) \cdots G(\lambda_n^{k_n}, \chi_c) \\ &= \frac{1}{q} \sum_{(k_1, \dots, k_n), 1 \leq k_i \leq d_i-1} G(\lambda_1^{k_1}, \chi_{a_1}) \cdots G(\lambda_n^{k_n}, \chi_{a_n}) \sum_{c \in \mathbb{F}_q^\times} \bar{\chi}_b(c) \bar{\lambda}_1^{-k_1}(c) \cdots \bar{\lambda}_n^{-k_n}(c) \end{aligned} \tag{F.13}$$

$$= \frac{1}{q} \sum_{(k_1, \dots, k_n), 1 \leq k_i \leq d_i - 1} G(\lambda_1^{k_1}, \chi_{a_1}) \cdots G(\lambda_n^{k_n}, \chi_{a_n}) G(\overline{\lambda_1}^{-k_1} \cdots \overline{\lambda_n}^{-k_n}, \overline{\chi_b}) \tag{F.14}$$

where in (F.13) we used Proposition 6.2.2 twice, to get

$$\overline{\lambda_j}^{k_j}(a_j) G(\lambda_j^{k_j}, \chi_c) = \overline{\lambda_j}^{k_j}(c) \overline{\lambda_j}^{-k_j}(a_j) G(\lambda_j^{k_j}, \chi_1) = \overline{\lambda_j}^{k_j}(c) G(\lambda_j^{k_j}, \chi_{a_j}).$$

Now we apply Theorem 6.2.1 to get that $|G(\lambda_i^{k_i}, \chi_{a_i})| = \sqrt{q}$. Note

$$(\overline{\lambda_1}^{-k_1} \cdots \overline{\lambda_n}^{-k_n})(\xi^t) = e^{(2\pi i) \left(\frac{k_1}{d_1} + \cdots + \frac{k_n}{d_n} \right) t}$$

is the trivial character iff $(k_1, \dots, k_n) \in S$. Hence $|G(\overline{\lambda_1}^{-k_1} \cdots \overline{\lambda_n}^{-k_n}, \overline{\chi_b})| = 1$ if $(k_1, \dots, k_n) \in S$ and \sqrt{q} otherwise. Using this and the triangle inequality, (F.14) becomes

$$|N - q^{n-1}| \leq \frac{1}{q} [q^{\frac{n}{2}} |S| + q^{\frac{n+1}{2}} ((d_1 - 1) \cdots (d_n - 1) - |S|)],$$

proving the theorem. □

4 Applications to Waring’s Problem

Now we derive Small’s bound for Waring’s constant $g(d, q)$, the minimum n such that (F.8) has a solution with $d_1 = \cdots = d_n = d$ for all b . By Lemma 6.3.1, $g(d, q) = g(\gcd(d, q - 1), q)$, so it suffices to consider the case $d|q - 1$.

First, note that sufficient condition for Waring’s constant to exist is that the set $\{y^d | y \in \mathbb{F}_q\}$ is not contained in a proper subfield of \mathbb{F}_q . Since this set is generated multiplicatively by ξ^d , and any subfield is multiplicatively generated by $\xi^{\frac{p^r - 1}{p^k - 1}}$ for some $k|d$, writing $q = p^r$ with p prime we need

$$\frac{p^r - 1}{p^k - 1} \nmid d \quad \text{for every proper divisor } k \text{ of } r. \tag{F.15}$$

Apply Theorem 6.3.2 (dropping the term with $M(d_1, \dots, d_n)$) to get

$$N \geq q^{n-1} - (d - 1)^n q^{\frac{n-1}{2}} \tag{F.16}$$

This is positive when

$$q^{\frac{n-1}{2}} > (d - 1)^n \iff \frac{n}{2} (\ln q - 2 \ln(d - 1)) > \frac{\ln q}{2} \tag{F.17}$$

Thus we obtain the following bound for $g(d, q)$:

Theorem 6.4.1. *Suppose $d|q - 1$ and $q > (d - 1)^2$. Then*

$$g(d, q) \leq \left\lceil \frac{\ln q}{\ln q - 2 \ln(d - 1)} + 1 \right\rceil.$$

Note that in particular, (F.17) for $n = 2$ allows us to make the “inverse” statement that if $q > (d - 1)^4$, then the equation $y_1^d + y_2^d = b$ has a solution for any $b \in \mathbb{F}_q$. That is, for any d , in any sufficiently large finite field every element can be written as a sum of 2 d th powers.

Bibliography

- [AD08] T. Andreescu and G. Dospinescu. *Problems from the Book*. XYZ Press, 2008.
- [Ahl79] L. Ahlfors. *Complex Analysis*. McGraw-Hill, 1979.
- [Apo94] T. Apostol. *Modular forms and Dirichlet series*. Number 110 in GTM. Springer, 2nd edition, 1994.
- [Bha04] M. Bhargava. Higher composition laws i: A new view on gauss composition, and quadratic generalizations. *Annals Math.*, 159(1):217–250, Jan. 2004.
- [BS66] Z. Borevich and I. Shafarevich. *Number Theory*. Academic Press, 1966.
- [CE56] H. Cartan and S Eilenberg. *Homological Algebra*, volume 19. Princeton Math Series, 1956.
- [CF69] J. Cassels and A. Frohlich, editors. *Algebraic Number Theory*. Academic Press, 1969.
- [Coh03] P. Cohn. *Further Algebra and Applications*. Springer, 2003.
- [Cox89] D. Cox. *Primes of the form $x^2 + ny^2$* . John Wiley & Sons, Inc., 1989.
- [Dal11] C. Dalawat. Splitting primes, March 2011.
- [Eti] P. Etingof. Introduction to representation theory.
- [GBGL10] T. Gowers, J. Barrow-Green, and I. Leader. *The Princeton Companion to Mathematics*. Princeton University Press, 2010.
- [Lan94] S. Lang. *Algebraic Number Theory*. Number 110 in GTM. Springer, 1994.
- [Mac71] S. MacLane. *Categories for the Working Mathematician*. Springer, 1971.
- [Mil08] J. Milne. *Class Field Theory*. 4.00 edition, 2008.
- [Neu99] J. Neukirch. *Algebraic Number Theory*. Springer, 1999.
- [Rot09] J. Rotman. *An Introduction to Homological Algebra*. Springer, 2009.

- [SD72] H. P. F. Swinnerton-Dyer. On l -adic representations and congruences for coefficients of modular forms. 1972.
- [Ser73] J.-P. Serre. *A Course in Arithmetic*. Number 7 in GTM. Springer, 1973.
- [Ser79] J.-P. Serre. *Local Fields*. Number 67 in GTM. Springer, 1979.
- [Ser03] J.-P. Serre. On a theorem of Jordan. *Bull. Amer. Math. Soc.*, 40(4):429–440, 2003.
- [Sil86] J. Silverman. *The Arithmetic of Elliptic Curves*. Number 106 in GTM. Springer, 1986.
- [Sil94] J. Silverman. *Advanced Topics in the Arithmetic of Elliptic Curves*. Number 151 in GTM. Springer, 1994.
- [Was82] L. Washington. *Introduction to Cyclotomic Fields*. Number 83 in GTM. Springer, 1982.
- [Wei] J. Weinstein. Analytic class number formula.

Index

- abstract Galois group, 235
- adeles, 157
- Artin's conjecture, 309
- automorphic form, 307

- bar resolution, 180
- biquadratic reciprocity, 285
- Brauer group, 216
- Brauer-Hasse-Noether theorem, 267

- central simple algebra, 216
- centralizer, 217
- chain map, 171
- change of group, 192
- Chebotarev density theorem, 296, 297
- class formation, 235
- class group, 33
- cohomological functor, 177
- cohomology, 172
- cohomology of lattices, 260
- cohomology of units, 226
- coinduced module, 183
- Comparison theorem, 172
- complete resolution, 187
- complex, 171
- complex multiplication, 317
- conductor, 156
- congruence subgroup, 154
- corestriction, 192
- cubic reciprocity, 289
- cup product, 189
- cuspidal form, 312
- Cyclotomic polynomials, 73

- decomposition group, 27
- Dedekind domain, 18

- degree equation, 22
- density, 297
- derivation, 181
- derived functors, 173
- descent, 212
- dimension shifting, 188
- Dirichlet's S-unit theorem, 71
- Dirichlet's theorem for number fields, 297
- Dirichlet's unit theorem, 67
- discrete valuations, 17
- double centralizer theorem, 217

- Euler's theorem, 430
- existence theorem, 249
- Ext, 174

- factor set, 181
- Fermat's little theorem, 430
- first inequality, 256
- fixed field theorem, 473
- formation, 236
- Frobenius element, 147
- Fundamental theorem of Galois theory, 475
- fundamental unit, 237

- Galois cohomology, 207
- Galois extension, 473
- Galois group, 473
- Galois representation, 308
- Gauss composition, 58
- Gauss's lemma, 445, 455
- global reciprocity, 155
- Größencharacter, 310
- Größencharacter of elliptic curve, 339
- group cohomology, 179
- group homology, 182

- group ring, 179
- Hasse norm theorem, 290
- Hasse-Minkowski, 291
- Hecke character, 309
- Hensel's lemma, 96
- Herbrand quotient, 201
- Hilbert class field, 303
- Hilbert class field of imaginary quadratic field, 326
- Hilbert symbol, 246
- Hilbert's Theorem 90, 207
- homology, 172
- homotopy, 172
- idele class group, 159
- ideles, 157
- induced module, 183
- inertia group, 27
- inflation, 192
- inflation-restriction exact sequence, 197
- injective object, 170
- integral, 3
- invariant map, 228
- Krasner's lemma, 101
- Kronecker-Weber theorem, 166
- Kummer theory, 208
- L-function, 308
- L-series of elliptic curve, 342
- Langlands program, 307
- Legendre symbol, 280, 443
- local reciprocity, 151
- long exact sequence, 176
- Lucas's Theorem, 467
- main theorem of complex multiplication, 335
- maximal abelian extension of imaginary quadratic field, 330
- Minkowski's Theorem, 35
- Mittag-Leffler, 358
- modular form, 312
- modularity theorem, 313
- modulus, 153
- narrow class group, 58
- Noether-Skolem theorem, 217
- nonabelian cohomology, 204
- norm group, 152
- norm limitation theorem, 240
- normal basis, 475
- normal extension, 474
- normality, 4
- numerical norm, 34
- order, 428
- pointed set, 204
- prime number theorem, 367
- product formula, 115
- product formula for Hilbert symbols, 283
- projective object, 170
- proper ideal, 56
- quadratic reciprocity, 284, 444
- quadratic residue, 443
- quintic equations, 477
- ramification index, 22
- rational roots theorem, 5
- ray class group, 153
- reciprocity law, 238
- reciprocity laws, 280
- representation, 217
- residue class degree, 22
- resolutions, 170
- restriction, 192
- resultant, 469
- Riemann hypothesis, 383
- ring class field, 304
- roots, linear independence of, 211
- Sato-Tate conjecture, 314
- second inequality, 263
- Serre's conjecture, 312
- Shapiro's lemma, 184
- splitting field, 473
- Stirling's approximation, 360

strong reciprocity, 283

Tate groups, 185

Tate module, 313

Tate's Theorem, 202

tensor, 214

topological G-module, 203

Tor, 175

torsor, 321

transfer, 198

units in number fields, 67

von Mangoldt's formula, 378

von Mangoldt's Theorem, 376

weak reciprocity, 280

Wedderburn's structure theorem, 217

Weil group, 152

Wilson's theorem, 439

Wolstenholme's Theorem, 466

zeta function, 370